

Human Extinction Learning Is Accelerated by an Angiotensin Antagonist via Ventromedial Prefrontal Cortex and Its Connections With Basolateral Amygdala

Feng Zhou, Yayuan Geng, Fei Xin, Jialin Li, Pan Feng, Congcong Liu, Weihua Zhao, Tingyong Feng, Adam J. Guastella, Richard P. Ebstein, Keith M. Kendrick, and Benjamin Becker

ABSTRACT

BACKGROUND: Deficient extinction learning and threat adaptation in the ventromedial prefrontal cortex (vmPFC)-amygdala circuitry strongly impede the efficacy of exposure-based interventions in anxiety disorders. Recent animal models suggest a regulatory role of the renin-angiotensin system in both these processes. Against this background, the present randomized placebo-controlled pharmacologic functional magnetic resonance imaging experiment aimed at determining the extinction enhancing potential of the angiotensin II type 1 receptor antagonist losartan (LT) in humans.

METHODS: Seventy healthy male subjects underwent Pavlovian threat conditioning and received single-dose LT (50 mg) or placebo administration before extinction. Psychophysiological threat reactivity (skin conductance response) and neural activity during extinction served as primary outcomes. Psychophysiological interaction, voxelwise mediation, and novel multivariate pattern classification analyses were used to determine the underlying neural mechanisms.

RESULTS: LT significantly accelerated the decline of the psychophysiological threat response during within-session extinction learning. On the neural level, the acceleration was accompanied and critically mediated by threat-specific enhancement of vmPFC activation. Furthermore, LT enhanced vmPFC-basolateral amygdala coupling and attenuated the neural threat expression, particularly in the vmPFC, during early extinction.

CONCLUSIONS: Overall the results indicate that LT facilitates within-session threat memory extinction by augmenting threat-specific encoding in the vmPFC and its regulatory control over the amygdala. The findings document a pivotal role of angiotensin regulation of extinction learning in humans and suggest that adjunct LT administration has the potential to facilitate the efficacy of exposure-based interventions in anxiety disorders.

Keywords: Angiotensin, Anxiety disorders, Extinction, fMRI, Functional magnetic resonance imaging, Losartan, Ventromedial prefrontal cortex, vmPFC

<https://doi.org/10.1016/j.biopsych.2019.07.007>

Extinction learning refers to the attenuation of a previously learned defensive response when the threat-predictive stimulus is repeatedly encountered in the absence of adverse consequences. Exposure-based interventions capitalize on extinction learning mechanisms to reduce excessive fear in patients with anxiety-related disorders, and they are considered an efficient therapy in these illnesses (1). A significant number of patients, however, do not adequately respond to exposure therapy (2), and impaired extinction processes are considered one key mechanism underpinning the lack of therapeutic efficacy (3). Anxiety-related disorders are highly prevalent and associated with significant psychosocial impairments and societal costs (4). As such, innovative strategies to improve the efficacy or shorten the duration of exposure therapies are urgently needed.

Whereas the behavioral and neural pathomechanisms underlying anxiety disorders are becoming increasingly understood, the translation into efficacious clinical interventions remains inadequate. Notably, the neural mechanisms mediating extinction are extremely well conserved over the course of evolutionary time; hence, the identification in animal models of receptor targets sensitive to pharmacologic modulation that facilitate neural plasticity in pathways supporting extinction learning can feasibly augment the efficacy of exposure-based interventions (5).

Animal models and human neuroimaging research have demonstrated a crucial role of the infralimbic cortex, which is homologous to the human ventromedial prefrontal cortex (vmPFC) and its interactions with the amygdala in extinction learning (6–12). The vmPFC is critically engaged in the reduction of threat expression during extinction (3,12,13), and it governs the

SEE COMMENTARY ON PAGE 874

amygdala inhibition of the conditioned threat response (8,9). Translational models suggest that dysfunctions in amygdala-prefrontal neuroplasticity contribute to extinction failure in anxiety disorders (14). Converging evidence from clinical research suggests that anxiety disorders are characterized by deficient extinction, hypoactivation within the vmPFC, and attenuated vmPFC-amygdala functional connectivity (6,8,10,13,15).

Intriguingly, recent evidence suggests that the renin-angiotensin (RA) system, primarily known for its role as a blood pressure and renal absorption regulator, represents a promising target to facilitate extinction (5). Central angiotensin receptors are densely expressed in limbic and prefrontal brain regions and are critical to changes in neuroplasticity and extinction (16–18). Initial studies in rodents have demonstrated the potential of pharmacologic modulation of RA signaling toward facilitating extinction using the selective competitive angiotensin II type 1 (AT1) antagonist losartan (LT) (19,20). LT is an approved treatment for high blood pressure, and it has an excellent safety record (21,22). Notably, increasing evidence suggests an association between the RA system and anxiety-related disorders [e.g., Terock *et al.* (23)], and initial clinical observations suggested unexpected beneficial effects of LT on memory and anxiety symptomatology (5,21) [see also a recent study demonstrating effects of LT on aversive learning in healthy participants in Pulcu *et al.* (24)]. Accumulating evidence suggests that LT modulation of central RA signaling represents a promising target to enhance extinction learning.

Against this background, we conducted a preregistered randomized placebo (PLC)-controlled pharmacologic experiment to determine whether targeting the RA system can facilitate extinction in humans. To uncover the underlying neural mechanisms, functional magnetic resonance imaging (fMRI) and psychophysiological threat responses (skin conductance response [SCR]) were acquired simultaneously. The specific goal was to determine the potential of LT (50 mg, single oral dose) as a therapeutic candidate for the clinical augmentation of extinction learning. Previous translational research leads us to expect that LT would accelerate attenuation of the psychophysiological threat responses and that enhanced extinction would be mediated by two neural processes: 1) increased activation in the vmPFC and attenuation of its threat expression in the context of 2) stronger functional interaction of the vmPFC with the amygdala.

METHODS AND MATERIALS

Participants and Experimental Protocols

Seventy healthy male subjects underwent a validated Pavlovian threat acquisition and extinction procedure with simultaneous fMRI and SCR acquisition. To reduce variance related to differences in extinction-related neural activity between men and women (25,26) and sex differences in the responses to RA blockade (27), only male participants were included in the present proof-of-concept study [for a similar approach, see Eckstein *et al.* (28)]. Participants were asked to abstain from caffeinated drinks on the day of the assessment (e.g., coffee, tea, energy beverages). Because of technical issues (SCR recording failure, $n = 3$) or absence of threat acquisition ($n = 8$), data from 11 subjects were excluded,

leading to 30 LT- and 29 PLC-treated subjects for the evaluation of the primary study hypotheses. For exclusion criteria and a description of the study sample, see [Supplemental Methods](#).

The experiment consisted of 3 sequential stages: 1) acquisition, 2) treatment administration, and 3) extinction. Twenty minutes after acquisition, participants were administered either a single dose (50 mg, by mouth) of LT or PLC, packed in identical capsules. Capsules were dispensed by an independent researcher according to a computer-generated randomization list ($n = 2$ groups). Consistent with the pharmacodynamic profile of LT [e.g., following oral administration peak plasma levels are reached after 90 minutes, and the terminal elimination half-life of LT proximately ranges from 1.5 to 2.5 hours (29,30–32)], extinction was started 90 minutes after treatment. Although previous studies reported no effects of single-dose LT on cardiovascular activity or mood (18,29,33), these indices were monitored to additionally control primarily for unspecific effects of treatment. The experimental timeline is provided in [Figure 1A](#) (for the pharmacodynamic profile and assessment of confounders see [Supplemental Methods](#)).

The study was approved by the local institutional ethics committee; it adhered to the Declaration of Helsinki and was a preregistered trial ([ClinicalTrials.gov](https://clinicaltrials.gov), registration number: NCT03396523; URL: <https://clinicaltrials.gov/ct2/show/NCT03396523>). The experiments were conducted at the Brain Imaging Center of the Southwest University, Chongqing, China, between January and October 2018.

Experimental Paradigm

During the acquisition stage, participants were repeatedly presented with two different colored squares—the conditioned stimulus (CS). One CS (CS+, 4 seconds) was pseudorandomly paired with a mild electric shock (unconditioned stimulus [US], 2 ms) with 43% contingency, whereas the other CS (CS–, 4 seconds) was never paired with a US. Acquisition was followed by extinction, in which the same cues were presented without US ([Supplemental Methods](#)). To enhance threat memory acquisition and to increase the statistical power to determine treatment effects, both learning phases included 2 subsequent runs of the task [for a similar approach, see Feng *et al.* (34)]. The extinction procedure encompassed 2 runs, with each run including 10 CS+ trials without the US, intermixed with 10 presentations of the CS–. No trial-type was repeated more than twice in a row during either acquisition or extinction. Before each run, subjects were informed that “the experimental runs are independent and you may or may not receive the electric shock”; thus, subjects were unable to predict the presence or absence of the US at the beginning of the respective run.

Skin Conductance Response Analysis

Following previous studies (28,35–37), both the CS+ and CS– stimuli of each run were divided into early (first half) and late (last half) to determine learning-related changes during extinction. Consistent with previous studies examining novel strategies to facilitate extinction (35,36), psychophysiological threat responses were defined as baseline-corrected CS+ by subtracting the mean responses to the CS– (see [Supplemental Methods](#)). Treatment effects were determined using a phase




Figure 1. Experimental timeline and losartan (LT) effects on psychophysiological threat responses. **(A)** Experimental timeline and schematic synopsis of functional magnetic resonance imaging (fMRI) tasks. **(B)** Psychophysiological threat responses (conditioned stimulus [CS]⁺ > CS⁻) during acquisition demonstrating successful CS discrimination with enhanced skin conductance response (SCR) to the CS⁺ relative to the CS⁻ in both groups. **(C)** Mean SCR for CS⁻ presentations during acquisition. **(D)** Psychophysiological threat responses (CS⁺ > CS⁻) during extinction learning. **(E)** Mean SCR for CS⁻ presentations during extinction learning. Early run, first trials (1–5); late run, last trials (6–10) in each run. [†]*p* < .05, 1-tailed. BP, blood pressure; HR, heart rate; ISI, interstimulus interval; PANAS, Positive and Negative Affect Schedule; PLC, placebo; STAI, State-Trait Anxiety Inventory; US, unconditioned stimulus.

(early, late) × run (run1, run2) × treatment (LT, PLC) 3-way mixed analysis of variance (ANOVA) with psychophysiological threat responses as the dependent variable. The additional analysis of the CS⁻ served primarily to exclude that LT had an effect on the skin conductance response per se.

MRI Acquisition and Analysis

MRI data were acquired using a Siemens TRIO 3T system with a 12-channel head coil (Siemens, Erlangen, Germany). Functional time-series were processed using SPM12 (<https://www.fil.ion.ucl.ac.uk/spm/software/spm12/>). On the first level, the CS⁺ and CS⁻ stimuli were modeled, and condition-specific regressors for early (first half) and late (second half) of extinction were defined (see Supplemental Methods).

Whole-Brain Analyses

Effects of LT on extinction were assessed using a whole-brain phase (early, late) × run (run1, run2) × treatment (LT, PLC) 3-way mixed ANOVA with the CS⁺ > CS⁻ contrasts as the

dependent variable. Significant interaction effects were further disentangled by 2 independent post hoc approaches to warrant both high regional specificity (whole-brain voxelwise post hoc *t* tests) and high robustness (leave-one-subject-out cross-validation [LOSO-CV] procedure) (38). Group-level analyses (including LOSO-CV) were conducted using FSL Randomise (FMRIB Software Library; <http://fsl.fmrib.ox.ac.uk/fsl/fslwiki/>) with permutation-based inferences (10,000 permutations). Significant clusters were determined using a height threshold of *p* < .001 (2-tailed) and an extent threshold of *p* < .05 (2-tailed) with cluster-based familywise error (FWE) correction.

Region of Interest: vmPFC Contributions During Early Extinction

Because of the critical role of the vmPFC in extinction (8,12,37,39,40) and our a priori regional hypothesis, we explored LT effects on stimulus-specific vmPFC activation during early extinction. To this end, activity estimates were extracted from an

Table 1. Participant Demographics and Control Measures

Measure	Time	LT Group (n = 30)	PLC Group (n = 29)	t_{57}	p Value
Age, Years		20.50 (1.80)	20.86 (1.68)	-0.80	.43
Body Mass Index, kg/m ²		22.33 (2.51)	21.34 (2.52)	1.51	.14
Systolic Blood Pressure	Before drug administration	110.03 (6.42)	107.59 (5.49)	1.57	.12
	Before extinction	111.17 (6.07)	108.79 (6.18)	1.49	.14
	After extinction	111.00 (5.69)	109.72 (5.76)	0.86	.40
Diastolic Blood Pressure	Before drug administration	71.70 (7.86)	71.10 (5.95)	0.33	.74
	Before extinction	72.70 (7.51)	70.17 (4.88)	1.53	.13
	After extinction	72.50 (8.57)	72.34 (4.15)	0.09	.93
Heart Rate	Before drug administration	76.93 (9.27)	76.03 (10.09)	0.36	.72
	Before extinction	75.33 (10.06)	74.17 (10.04)	0.44	.66
	After extinction	77.27 (9.44)	76.76 (10.93)	0.19	.85
STAI State Anxiety Score	Before extinction	41.90 (8.88)	41.24 (10.51)	0.26	.80
	After extinction	37.23 (8.48)	38.14 (8.16)	-0.42	.68
PANAS Negative Affect Score	Before extinction	16.17 (6.95)	16.34 (9.08)	-0.08	.93
	After extinction	14.43 (6.46)	14.59 (7.60)	-0.08	.93
PANAS Positive Affect Score	Before extinction	22.73 (7.55)	23.07 (6.83)	-0.18	.86
	After extinction	23.70 (7.67)	22.34 (6.02)	0.75	.45

Values are presented as mean (SD).

LT, losartan; PANAS, Positive and Negative Affect Schedule; PLC, placebo; STAI, Spielberger State-Trait Anxiety Inventory.

anatomically defined vmPFC region of interest (Supplemental Methods) and subjected to a stimulus (CS+, CS-) × run (run1, run2) × treatment (LT, PLC) 3-way mixed ANOVA.

Neural Threat Expression: Multi-Voxel Pattern Analysis

Following Reddan *et al.* (39), a neural pattern of threat was developed to differentiate CS+ versus CS- (trained and tested on the acquisition data) and subsequently applied to early extinction activation to determine treatment effects on the neural threat expression (Supplemental Methods).

Voxelwise Mediation Analyses

To determine whether treatment effects on vmPFC activation (CS+ > CS-) during early extinction critically contributed to the accelerated attenuation of the psychophysiological threat responses (SCR, CS+ > CS-), voxelwise mediation analyses were conducted (Mediation Toolbox: <https://github.com/canlab/MediationToolbox>) (41,42). Mediation effects were inferred using bootstrapping (10,000 replacements) and false discovery rate (FDR) correction.

Network-Level Effects: Functional Connectivity Analysis

Given that both human and animal studies strongly implicate vmPFC-mediated inhibition of the amygdala as a key extinction mechanism (9–12), a functional connectivity analysis (43) was used to determine treatment effects on the vmPFC-amygdala coupling during early extinction. We hypothesized that LT-induced extinction enhancement would be accompanied by stronger functional interaction between these regions (1-sided).

RESULTS

Participants

Consistent with previous studies (18,29,33), no effects of drug or placebo on cardiovascular and affective indices were observed, which, together with the chance level guesses for treatment, argues against unspecific confounding effects of treatment (Table 1). During the pretreatment acquisition phase, both groups exhibited successful threat acquisition on the psychophysiological (Figure 1B, C) and neural (Supplemental Results and Supplemental Figure S1) levels. Importantly, 2-sample *t* tests did not reveal between-group activation differences during this stage.

Reduced Psychophysiological Threat Responses During Early Extinction

A mixed ANOVA model using the psychophysiological threat response during extinction learning as a dependent variable demonstrated a significant main effect of run ($F_{1,57} = 17.063$; $p < .001$; partial $\eta^2 = .230$; η^2 indicates effect size in terms of eta squared) reflecting decreased psychophysiological threat responses in run2 compared to run1 and a marginally significant main effect of phase ($F_{1,57} = 3.387$; $p = .071$; partial $\eta^2 = .056$) reflecting decreased psychophysiological threat responses during late extinction. Together, these results demonstrated successful extinction learning. Moreover, a significant treatment and phase interaction effect ($F_{1,57} = 5.017$, $p = .029$, partial $\eta^2 = .081$) was observed, with post hoc 2-sample *t* tests indicating that relative to the PLC group, the LT group exhibited decreased psychophysiological threat responses during early extinction learning across both runs ($t_{57} = -2.179$; $p = .034$, $d = -0.567$; d indicates effect size in terms of Cohen's *d*), suggesting accelerated extinction learning. Exploratory run-specific




Figure 2. Losartan (LT) treatment effects on brain activity (conditioned stimulus [CS]⁺ > CS⁻) during extinction learning. **(A)** Ventromedial prefrontal cortex (vmPFC) activity showed significant treatment by phase interaction effect. **(B)** LT specifically increased vmPFC activity during early extinction learning. **(C)** Mean vmPFC activity (CS⁺ > CS⁻) extracted from the regions of interest depicted in **(D)** showed that LT increased vmPFC activity during early, but not late, extinction learning. **(D)** Overlay of all 59 leave-one-subject-out cross-validation regions of interest. All regions of interest were created leaving out one subject at the group-level statistic (cluster-level familywise error [FWE]-corrected). Statistical images were thresholded at $p < .05$ (2-tailed), cluster-level FWE-corrected with a cluster-forming threshold of $p < .001$ (2-tailed). Examples of unthresholded patterns are presented in the inserts; small squares indicate voxel statistical weight; red-outlined squares indicate significance at $p_{\text{clusterFWE}} < .05$. The filled curve indicates the null-hypothesis distribution of the difference of means (Δ), and the 95% confidence interval of Δ is illustrated by the black line. ** $p < .01$. L, left; n.s., not significant; PLC, placebo; R, right.

analyses confirmed that LT treatment enhanced early extinction learning during both initial and repeated extinction learning (run1 $t_{57} = 1.805$, $p = .038$, $d = 0.470$; run2 $t_{57} = 2.012$, $p = .024$, $d = 0.525$; 2-sample t tests comparing the treatment groups, 1-tailed; **Figure 1D**). No further significant main or interaction effects were observed on the psychophysiological threat responses (all $p > .4$), nor were any effects of treatment observed on the safety signal (CS⁻; all $p > .15$; **Figure 1E**), confirming the threat-specific effects of LT and the absence of unspecific treatment effects on the SCR psychophysiological signal.

To determine whether LT generally reduced the SCR to threat signals rather than facilitating extinction learning, we also explored between-group differences in the SCR reactivity to the first CS⁺ occurrence in each run. The lack of significant

differences in the SCR reactivity to the first CS⁺ trials (run1 $t_{57} = -1.305$, $p = .197$, $d = -0.338$; run2 $t_{57} = -0.8975$, $p = .3732$, $d = -0.233$) confirmed specific effects on extinction learning.

Increased Threat-Specific vmPFC Engagement During Early Extinction

Whole-brain analysis revealed a significant treatment and phase interaction effect on extinction-related neural activity (CS⁺ > CS⁻) in the vmPFC (peak Montreal Neurological Institute (MNI) $x, y, z = -3, 27, -12$; $F_{1,57} = 23.582$; $p_{\text{clusterFWE}} = .011$; $k = 187$; **Figure 2A**). Regional specificity of treatment effects was examined with voxelwise whole-brain comparisons demonstrating increased vmPFC activity following LT relative to PLC during early (peak MNI $x, y, z = 6, 48, -3$; $t_{57} = 4.505$;




Figure 3. Losartan (LT) treatment specifically increased ventromedial prefrontal cortex (vmPFC) activity to threat stimulus (conditioned stimulus [CS]⁺) during early extinction learning. **(A)** LT increased vmPFC activity to CS⁺ but not CS⁻ in the early extinction phase in both runs. * $p < .05$ and *** $p < .001$; error bars represent standard errors. **(B)** Single-trial analysis confirmed that LT increased vmPFC activity to CS⁺ in early trials during extinction learning. * $q < .05$, false discovery rate corrected. Data are represented as group mean \pm SEM. **(C)** vmPFC region of interest. n.s., not significant; PLC, placebo.

$p_{\text{clusterFWE}} = .013$; 2-tailed $k = 139$) but not late extinction (Figure 2B). Post hoc analyses with the LOSO-CV procedure further demonstrated that LT—relative to PLC—increased vmPFC activation during early ($t_{57} = 3.417$; $p = .001$; $d = 0.890$) but not late ($t_{57} = -1.556$; $p = .125$; $d = -0.405$) extinction (Figures 2C and 2D display an overlay of all leave-one-subject-out regions of interest). Note that the vmPFC exhibited robust early extinction learning–induced treatment and phase interaction effect.

Further exploration of vmPFC contributions during early extinction by means of extraction of beta estimates revealed significant main effects of stimulus type and run, and a significant treatment and stimulus type interaction effect (see Supplemental Results). Exploratory post hoc 2-sample t tests demonstrated that LT enhanced threat-specific vmPFC reactivity during both the initial (run1 $t_{57} = 2.141$; $p = .037$; $d = 0.557$) and the repeated (run2 $t_{57} = 2.126$; $p = .038$; $d = 0.554$) extinction in the absence of effects on the safety signal (CS⁻, $p > .29$; Figure 3A).

Consistent with our hypothesis on accelerated extinction by LT, an exploratory single trial analysis (Supplemental Methods) revealed that LT specifically increased vmPFC activation during initial trials of re-exposure to the threat stimulus ($q < .05$, FDR corrected; Figure 3B).

Reduced Neural Threat Expression During Early Extinction

The threat-predictive pattern reliably evoked neural threat reactivity during acquisition [comparable to Reddan *et al.* (39)] (see Supplemental Results and Supplemental Figures S2A, S2B). Applying the threat-predictive pattern to early extinction activation (CS⁺ > CS⁻) using a LOSO-CV procedure demonstrated that first, in the entire sample, higher neural threat expression was associated with stronger psychophysiological threat reactivity ($r_{57} = 0.571$; $p < .001$) and confirmed functional relevance of the neural expression of threat (39). Second, relative to PLC, LT significantly decreased the

magnitude of the threat-predictive pattern expression ($t_{57} = -2.091$; $p = .041$; $d = -0.544$, 2-sample t test), confirming attenuated neural threat expression during early extinction. Based on our a priori regional hypothesis, and the key role of the vmPFC in extinction (3,8,13,35), a vmPFC-focused partial threat expression analysis was conducted. In concordance with the whole-brain results, LT significantly attenuated the vmPFC partial threat pattern expression (CS⁺ > CS⁻) during early extinction ($t_{57} = -3.410$; $p = .001$; $d = -0.888$; Supplemental Results and Supplemental Figure S2C).

vmPFC Activation Drives LT-Induced Accelerated Extinction

The voxelwise mediation analysis aimed at further determining the relationship between treatment, psychophysiological threat attenuation, and the underlying neural basis. Conjunction effects (paths a, b, and $a \times b$) were observed in a vmPFC cluster (peak MNI $x, y, z = -3, 45, -15$; $z = -3.692$; $q < .05$; FDR-small volume corrected in the anatomically defined vmPFC, $k = 138$), demonstrating that LT increased vmPFC activation (path a), while activation in this region was associated with stronger suppression of psychophysiological threat independent of treatment (path b). Importantly, the $a \times b$ mediation effect reached significance, indicating that vmPFC activation critically mediated the effects of LT on extinction acceleration (Figure 4A; for details, see Supplemental Results). To test the robustness and to visualize the mediation effect, an independent vmPFC-focused mediation analysis was conducted, which confirmed the critical contribution of the vmPFC (Figure 4B, $a \times b$ effect, bootstrapped $p = .016$; each path is shown in Figure 4C).

Enhanced vmPFC-Amygdala Coupling

During early extinction, LT induced enhanced functional coupling between the vmPFC and the right amygdala [peak MNI $x, y, z = 24, -3, -24$; $t_{57} = 3.557$; $q < .05$; 1-tailed; FDR-small volume corrected in the amygdala, $k = 13$; location




Figure 4. Ventromedial prefrontal cortex (vmPFC) activity mediated losartan (LT) treatment effect on accelerated extinction learning. **(A)** Sagittal slice showing regions whose activity increased response to the LT treatment in yellow (path a), regions whose activity significant negative correlated with psychophysiological threat responses while controlling for the treatment effect in green (path b), and regions whose activity showed significant mediation ($a \times b$) effect in blue. All images were thresholded at $q < .05$, false discovery rate corrected within the vmPFC mask. **(B)** Mediation path diagram with the brain activity in the vmPFC region of interest. **(C)** Examples of each path in the mediation path diagram. * $p < .05$; ** $p < .01$; *** $p < .001$. The filled curve indicates the null-hypothesis distribution of the difference of means (Δ), and the 95% confidence interval of Δ is illustrated by the black line. CS, conditioned stimulus; PLC, placebo; SCR, skin conductance response.

cyto-architecturally (44) mapped to the basolateral amygdala [BLA]; Figure 5A]. Subsequent examination of stimulus-specific connectivity estimates from the vmPFC-bilateral BLA pathway confirmed specific effects of LT on threat signal (CS+) processing ($t_{57} = 2.147$; $p = .036$; $d = 0.559$; Figure 5B).

DISCUSSION

The present study demonstrated that LT treatment accelerated the attenuation of a previously acquired psychophysiological threat response, indicating its potential to facilitate threat extinction learning in humans. During early extinction, the acceleration was critically mediated by enhanced threat-signal-specific vmPFC activity in the context of stronger functional coupling of the vmPFC with the BLA. These findings were further paralleled by a pattern classification approach showing that LT treatment accelerated attenuation of the neural threat expression, particularly in the vmPFC. Overall, the present findings provide first evidence for an important contribution of the RA system to fear extinction in humans and the potential of LT to accelerate extinction through effects on the vmPFC and its inhibitory connections with the BLA.

In humans, successful extinction is accompanied by decreased psychophysiological threat reactivity and concomitantly increased vmPFC activation in response to the threat signal (37,40,45). In the present study, LT treatment reduced the psychophysiological threat responses and selectively enhanced vmPFC activation in response to the threat signal (CS+) during early extinction, indicating its potential to accelerate extinction learning in humans. Moreover, the acceleration effects were found not only during the initial extinction learning (i.e., run1), but also in the following “new” learning process (the instruction before run2 allowed us to treat the design as two factors—run [run1, run2] and phase [early and late within each run]—instead of treating phases 1 to 4 across the entire extinction as a single factor; see Supplemental Results). The findings resemble previously observed LT-enhanced extinction learning in rodents (19,20) and further confirm the important contribution of the vmPFC to successful extinction. Exploring the stimulus-specific effects of LT treatment on vmPFC activation revealed that reactivity to the safety signal (CS-) remained unaffected. Decoding the temporal pattern of LT effects further suggested that LT specifically attenuated vmPFC reactivity during early re-exposure toward the previously




Figure 5. Losartan treatment effect on ventromedial prefrontal cortex (vmPFC)-amygdala functional coupling. **(A)** Sagittal slice showing that losartan (LT) increased functional connectivity between vmPFC and basolateral amygdala (BLA). The image was thresholded at $q < .05$, false discovery rate corrected within the amygdala mask. **(B)** Extracted generalized form of context-dependent psychophysiological interaction (gPPI) parameter estimates from the bilateral BLA showed that LT treatment specifically enhanced vmPFC-bilateral BLA functional connectivity during fear-associated stimulus presentation. The filled curve indicates the null hypothesis distribution of the difference of means (Δ), and the 95% confidence interval of Δ is illustrated by the black line. * $p < .05$. CS, conditioned stimulus; n.s., not significant; PLC, placebo.

conditioned threat signal. Nonpharmacologic stimulation of the vmPFC homologous infralimbic cortex accelerates extinction learning in rodents (46–48), whereas inactivation or lesioning of this region critically impedes threat reduction during extinction (8,13). Compatible with the present findings, previous studies demonstrated that nonpharmacologic stimulation of the vmPFC can enhance early extinction learning in humans (49), although nonspecific effects on CS– reactivity have also been reported (50).

Converging evidence from different research focuses suggests that the vmPFC, or the homologous infralimbic cortex in rodents, critically contributes to the reduction of threat expression during extinction learning (3,8,12,13) and regulates amygdala output to inhibit the conditioned threat response (8,9). Consonant with the proposed contribution of the vmPFC to extinction learning, LT-attenuated psychophysiological threat responses during early extinction were accompanied by an attenuated neural threat expression, particularly in the anatomically defined vmPFC. Compatible with previous studies that showed critical contributions of the vmPFC to extinction enhancement (46–48) and associations between activity in this region and psychophysiological threat reactivity during extinction (51), an additional mediation analysis revealed that higher vmPFC activation was associated with stronger suppression of the psychophysiological threat response. Notably, LT-facilitated suppression of the psychophysiological threat response crucially involved enhanced vmPFC activation (for convergent mediation effects of vmPFC threat expression, see [Supplemental Methods](#) and [Supplemental Results](#)), further emphasizing the key role of this region in extinction enhancement.

On the network level, LT-accelerated threat reduction during early extinction was paralleled by stronger functional communication between the vmPFC and the amygdala, specifically the basolateral subregion. Previous lesion studies in humans demonstrated a critical role of the BLA in threat processing (52) and of the vmPFC in inhibiting amygdala threat responses by exerting top-down control over this region (53). Animal models have further confirmed the importance of pathway-specific neuroplastic changes in the vmPFC-amygdala circuitry during

extinction memory formation (9–11) and suggest that vmPFC inputs to the amygdala instruct threat memory formation and/or gate the expression of conditioned threat (9) during early extinction (11). The present findings of CS+–specific increase in vmPFC-BLA functional connectivity following LT-treatment likely reflect an important modulatory role of angiotensin signaling on vmPFC regulation of the amygdala. Previously, animal models demonstrated that stimulation of vmPFC inputs to the amygdala promotes the formation of extinction memories (9). We suggest the notion that enhanced transmission in this pathway might reflect a core mechanism underlying angiotensin regulation of extinction learning. Angiotensin receptors are densely expressed in limbic and prefrontal regions critically engaged in extinction (16–18) and are considered to modulate learning-related neuroplasticity. LT is a selective competitive antagonist of the AT1 receptor, and it increases availability of angiotensin II–converted angiotensin IV—an agonist at the AT4 receptor subtype. The AT4 system is thought to play a role in neuroplasticity and learning and memory (16,17,33,54), a mechanism that is suggested to likely contribute to LT-induced extinction enhancement.

Consistent with previous animal models demonstrating the potential of LT to enhance extinction in rodents (19,20), the present study successfully demonstrated the potential of a single low-dose administration of LT to facilitate extinction learning in humans. In the context of recent findings suggesting a direct association between extinction-related vmPFC functioning and exposure therapy success (55), the current results indicate that LT represents a highly promising candidate to augment the efficacy of exposure-based interventions in therapeutic settings. On the neural level, the effects of LT were mediated by circuits consistently involved in anxiety disorders, with exaggerated threat reactivity and deficient extinction being associated with decreased vmPFC activation and dysfunction in the vmPFC-BLA circuit (6,8,10,13). Importantly, dysregulation in this circuitry normalizes during the course of successful treatment (56), suggesting that they represent treatment-responsive—rather than stable—markers and consequently promising targets for therapeutic interventions. A previous human neuroimaging study reported that LT enhances early

amygdala discrimination of threat and safety signals in highly anxious individuals (18). Both vmPFC dependent extinction learning and amygdala threat discrimination are considered core therapeutic mechanisms of exposure-based interventions, suggesting that LT facilitates effects of exposure therapy via synergistic effects on both mechanisms.

The present study observed effects of LT on within-session extinction learning. Although accumulating evidence suggests an association between extinction learning and associated vmPFC activation with better therapeutic response to exposure-based interventions (55,57,58), extinction retention (e.g., extinction recall) deficits have been reported frequently in anxiety-related disorders (59–61). To further explore the clinical efficacy of LT to enhance extinction, future studies should examine effects of LT on extinction retention after longer time intervals. In addition, despite these initial promising results, subsequent studies need to 1) determine the generalization of the effects to female subjects [for instance, see also a recent study demonstrating that LT enhances fear extinction in female rats in an estradiol-dependent manner (62)] and 2) evaluate its potential to enhance exposure-based interventions in clinical trials. Moreover, pharmacologic blockade of the AT1 receptors in animal models has been shown to inhibit the hypothalamic-pituitary-adrenal stress response via both central and peripheral AT1 receptors (63). Thus, we cannot rule out that effects on peripheral AT1 receptors might have contributed to the observed effects.

Overall, the present results indicate an important regulatory role of the RA system in fear extinction learning in humans. This role is mediated by modulatory effects on vmPFC threat processing and its interaction with the amygdala. From a clinical perspective, adjunct LT treatment could represent an innovative strategy to enhance the efficacy of exposure-based interventions.

ACKNOWLEDGMENTS AND DISCLOSURES

This work was supported by grants from National Natural Science Foundation of China (Grant Nos. 91632117 [to BB] and 31530032 [to KK]); Fundamental Research Funds for the Central Universities (Grant No. ZYGX2015Z002 [to BB]); and Science, Innovation and Technology Department of the Sichuan Province (Grant No. 2018JY0001 [to BB]).

FZ and BB designed the study, analyzed the data, and wrote the manuscript. FZ, YG, FX, JL, PF, CL, and WZ conducted the experiment. PF, TF, AG, RE, and KK revised the manuscript draft.

We thank Marianne Cumella Reddan for assistance with the threat expression analysis.

Unthresholded group-level statistical maps are available on NeuroVault (<https://neurovault.org/collections/4722/>) and code that supports the findings of this study is available from the corresponding author upon reasonable request.

A preprint of the manuscript has been archived on the [biorxiv.org](https://doi.org/10.1101/512657) repository (doi: <https://doi.org/10.1101/512657>).

The authors report no biomedical financial interests or potential conflicts of interest.

The present study was pre-registered at [ClinicalTrials.gov](https://clinicaltrials.gov) (trial name: Losartan and Emotion Processing; registration number: NCT03396523; URL: <https://clinicaltrials.gov/ct2/show/NCT03396523>).

ARTICLE INFORMATION

From the Clinical Hospital of Chengdu Brain Science Institute (FZ, YG, FX, JL, CL, WZ, KMK, BB) and Ministry of Education (MOE) Key Laboratory for Neuroinformation, University of Electronic Science and Technology of

China, Chengdu; China Center for Behavior Economics and Finance (RPE), Southwestern University of Finance and Economics, Chengdu; Faculty of Psychology (PF, TF), Southwest University, Chongqing; and Key Laboratory of Cognition and Personality (PF, TF), Ministry of Education, Southwest University, Chongqing, China; Autism Clinic for Translational Research (AJG) and Youth Mental Health Unit (AJG), Brain and Mind Centre, Central Clinical School, Faculty of Medicine, University of Sydney, Camperdown, Australia.

Address correspondence to Benjamin Becker, Ph.D., Center for Information in Medicine, University of Electronic Science and Technology of China, Chengdu 611731, China; E-mail: ben_becker@gmx.de.

Received Apr 25, 2019; revised Jun 27, 2019; accepted Jul 10, 2019.

Supplementary material cited in this article is available online at <https://doi.org/10.1016/j.biopsych.2019.07.007>.

REFERENCES

- Taylor S, Thordarson DS, Maxfield L, Fedoroff IC, Lovell K, Ogradniczuk J (2003): Comparative efficacy, speed, and adverse effects of three PTSD treatments: Exposure therapy, EMDR, and relaxation training. *J Consult Clin Psychol* 71:330.
- Loerinc AG, Meuret AE, Twohig MP, Rosenfield D, Bluett EJ, Craske MG (2015): Response rates for CBT for anxiety disorders: Need for standardized criteria. *Clin Psychol Rev* 42:72–82.
- Quirk GJ, Mueller D (2008): Neural mechanisms of extinction learning and retrieval. *Neuropsychopharmacology* 33:56.
- Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE (2005): Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 62:593–602.
- Morrison FG, Ressler KJ (2014): From the neurobiology of extinction to improved clinical treatments. *Depress Anxiety* 31:279–290.
- Milad MR, Quirk GJ (2012): Fear extinction as a model for translational neuroscience: ten years of progress. *Annu Rev Psychol* 63:129–151.
- Milad MR, Quirk GJ (2002): Neurons in medial prefrontal cortex signal memory for fear extinction. *Nature* 420:70.
- Giustino TF, Maren S (2015): The role of the medial prefrontal cortex in the conditioning and extinction of fear. *Front Behav Neurosci* 9:298.
- Bukalo O, Pinard CR, Silverstein S, Brehm C, Hartley ND, Whittle N, et al. (2015): Prefrontal inputs to the amygdala instruct fear extinction memory formation. *Sci Adv* 1:e1500251.
- Dejean C, Courtin J, Rozeske RR, Bonnet MC, Dousset V, Michelet T, et al. (2015): Neuronal circuits for fear expression and recovery: recent advances and potential therapeutic strategies. *Biol Psychiatry* 78:298–306.
- Delgado MR, Beer JS, Fellows LK, Huettel SA, Platt ML, Quirk GJ, et al. (2016): Viewpoints: Dialogues on the functional role of the ventromedial prefrontal cortex. *Nature Neurosci* 19:1545.
- Schiller D, Delgado MR (2010): Overlapping neural systems mediating extinction, reversal and regulation of fear. *Trends Cogn Sci* 14:268–276.
- Sotres-Bayon F, Cain CK, LeDoux JE (2006): Brain mechanisms of fear extinction: Historical perspectives on the contribution of prefrontal cortex. *Biol Psychiatry* 60:329–336.
- Fragale JE, Khariv V, Gregor DM, Smith IM, Jiao X, Elkabes S, et al. (2016): Dysfunction in amygdala-prefrontal plasticity and extinction-resistant avoidance: A model for anxiety disorder vulnerability. *Exp Neurol* 275:59–68.
- Sylvester C, Corbetta M, Raichle M, Rodebaugh T, Schlaggar B, Sheline Y, et al. (2012): Functional network dysfunction in anxiety and anxiety disorders. *Trends Neurosci* 35:527–535.
- Chai SY, Bastias MA, Clune EF, Matsacos DJ, Mustafa T, Lee JH, et al. (2000): Distribution of angiotensin IV binding sites (AT4 receptor) in the human forebrain, midbrain and pons as visualised by in vitro receptor autoradiography. *J Chemical Neuroanat* 20:339–348.
- Wright JW, Harding JW (2013): The brain renin-angiotensin system: A diversity of functions and implications for CNS diseases. *Pflugers Arch* 465:133–151.
- Reinecke A, Browning M, Breteler JK, Kappellmann N, Ressler KJ, Harmer CJ, et al. (2018): Angiotensin regulation of amygdala response

Losartan Enhances Fear Extinction in Humans via vmPFC

- to threat in high-trait anxious individuals. *Biol Psychiatry Cogn Neurosci Neuroimaging* 3:826–835.
19. Marvar PJ, Goodman J, Fuchs S, Choi DC, Banerjee S, Ressler KJ (2014): Angiotensin type 1 receptor inhibition enhances the extinction of fear memory. *Biol Psychiatry* 75:864–872.
 20. do Nascimento Lazaroni TL, Bastos CP, Moraes MFD, Santos RS, Pereira GS (2016): Angiotensin-(1-7)/Mas axis modulates fear memory and extinction in mice. *Neurobiol Learn Mem* 127:27–33.
 21. Khoury NM, Marvar PJ, Gillespie CF, Wingo A, Schwartz A, Bradley B, *et al.* (2012): The renin-angiotensin pathway in posttraumatic stress disorder: angiotensin-converting enzyme inhibitors and angiotensin receptor blockers are associated with fewer traumatic stress symptoms. *J Clin Psychiatry* 73:849–855.
 22. Goldberg AI, Dunlay MC, Sweet CS (1995): Safety and tolerability of losartan potassium, an angiotensin II receptor antagonist, compared with hydrochlorothiazide, atenolol, felodipine ER, and angiotensin-converting enzyme inhibitors for the treatment of systemic hypertension. *Am J Cardiol* 75:793–795.
 23. Terock J, Hannemann A, Janowitz D, Freyberger HJ, Felix SB, Dörr M, *et al.* (2019): Associations of trauma exposure and post-traumatic stress disorder with the activity of the renin-angiotensin-aldosterone-system in the general population. *Psychol Med* 49:843–851.
 24. Pulcu E, Shkreli L, Holst CG, Woud ML, Craske MG, Browning M, Reinecke A (2019): The effects of the angiotensin II receptor antagonist losartan on appetitive versus aversive learning: A randomized controlled trial. *Biol Psychiatry* 86:397–404.
 25. Merz CJ, Tabbert K, Schweckendiek J, Klucken T, Vaitl D, Stark R, *et al.* (2011): Neuronal correlates of extinction learning are modulated by sex hormones. *Soc Cogn Affect Neurosci* 7:819–830.
 26. Hwang MJ, Zsido RG, Song H, Pace-Schott EF, Miller KK, Lebron-Milad K, *et al.* (2015): Contribution of estradiol levels and hormonal contraceptives to sex differences within the fear network during fear conditioning and extinction. *BMC Psychiatry* 15:295.
 27. Miller JA, Cherney DZ, Duncan JA, Lai V, Burns KD, Kennedy CR, *et al.* (2006): Gender differences in the renal response to renin-angiotensin system blockade. *J Am Soc Nephrol* 17:2554–2560.
 28. Eckstein M, Becker B, Scheele D, Scholz C, Preckel K, Schlaepfer TE, *et al.* (2015): Oxytocin facilitates the extinction of conditioned fear in humans. *Biol Psychiatry* 78:194–202.
 29. Ohtawa M, Takayama F, Saitoh K, Yoshinaga T, Nakashima M (1993): Pharmacokinetics and biochemical efficacy after single and multiple oral administration of losartan, an orally active nonpeptide angiotensin II receptor antagonist, in humans. *Br J Clin Pharmacol* 35:290–297.
 30. Sica DA, Gehr TW, Ghosh S (2005): Clinical pharmacokinetics of losartan. *Clin Pharmacokinet* 44:797–814.
 31. Lo M-W, Goldberg MR, McCrear JB, Lu H, Furtek CI, Bjornsson TD (1995): Pharmacokinetics of losartan, an angiotensin II receptor antagonist, and its active metabolite EXP3174 in humans. *Clin Pharmacol Ther* 58:641–649.
 32. Samyuktha M, Vasanth P, Suresh K, Ramesh T, Ramesh M (2013): Formulation and evaluation of gastroretentive floating tablets of losartan potassium. *Int J Biopharm* 4:18–26.
 33. Mechaieil R, Gard P, Jackson A, Rusted J (2011): Cognitive enhancement following acute losartan in normotensive young adults. *Psychopharmacology* 217:51–60.
 34. Feng P, Becker B, Feng T, Zheng Y (2018): Alter spontaneous activity in amygdala and vmPFC during fear consolidation following 24 h sleep deprivation. *NeuroImage* 172:461–469.
 35. Schiller D, Monfils M-H, Raio CM, Johnson DC, LeDoux JE, Phelps EA (2010): Preventing the return of fear in humans using reconsolidation update mechanisms. *Nature* 463:49.
 36. Koizumi A, Amano K, Cortese A, Shibata K, Yoshida W, Seymour B, *et al.* (2016): Fear reduction without fear through reinforcement of neural activity that bypasses conscious exposure [published online ahead of print Nov 21]. *Nat Hum Behav*.
 37. Schiller D, Kanen JW, LeDoux JE, Monfils M-H, Phelps EA (2013): Extinction during reconsolidation of threat memory diminishes prefrontal cortex involvement. *Proc Natl Acad Sci U S A* 110:20040–20045.
 38. Esterman M, Tamber-Rosenau BJ, Chiu Y-C, Yantis S (2010): Avoiding non-independence in fMRI data analysis: Leave one subject out. *NeuroImage* 50:572–576.
 39. Reddan MC, Wager TD, Schiller D (2018): Attenuating neural threat expression with imagination. *Neuron* 100:994–1005.e1004.
 40. Milad MR, Wright CI, Orr SP, Pitman RK, Quirk GJ, Rauch SL (2007): Recall of fear extinction in humans activates the ventromedial prefrontal cortex and hippocampus in concert. *Biol Psychiatry* 62:446–454.
 41. Wager TD, Davidson ML, Hughes BL, Lindquist MA, Ochsner KN (2008): Prefrontal-subcortical pathways mediating successful emotion regulation. *Neuron* 59:1037–1050.
 42. Wager TD, van Ast VA, Hughes BL, Davidson ML, Lindquist MA, Ochsner KN (2009): Brain mediators of cardiovascular responses to social threat, part II: Prefrontal-subcortical pathways and relationship with anxiety. *NeuroImage* 47:836–851.
 43. McLaren DG, Ries ML, Xu G, Johnson SC (2012): A generalized form of context-dependent psychophysiological interactions (gPPI): A comparison to standard approaches. *NeuroImage* 61:1277–1286.
 44. Eickhoff SB, Stephan KE, Mohlberg H, Grefkes C, Fink GR, Amunts K, *et al.* (2005): A new SPM toolbox for combining probabilistic cytoarchitectonic maps and functional imaging data. *NeuroImage* 25:1325–1335.
 45. Phelps EA, Delgado MR, Nearing KI, LeDoux JE (2004): Extinction learning in humans: role of the amygdala and vmPFC. *Neuron* 43:897–905.
 46. Vidal-Gonzalez I, Vidal-Gonzalez B, Rauch SL, Quirk GJ (2006): Microstimulation reveals opposing influences of prelimbic and infralimbic cortex on the expression of conditioned fear. *Learn Mem* 13:728–733.
 47. Orsini CA, Maren S (2012): Neural and cellular mechanisms of fear and extinction memory formation. *Neurosci Biobehav Rev* 36:1773–1802.
 48. Do-Monte FH, Manzano-Nieves G, Quiñones-Laracuente K, Ramos-Molina L, Quirk GJ (2015): Revisiting the role of infralimbic cortex in fear extinction with optogenetics. *J Neurosci* 35:3607–3615.
 49. Guhn A, Dresler T, Andreatta M, Müller LD, Hahn T, Tupak SV, *et al.* (2014): Medial prefrontal cortex stimulation modulates the processing of conditioned fear. *Front Behav Neurosci* 8:44.
 50. Dittert N, Huettner S, Polak T, Herrmann MJH (2018): Augmentation of fear extinction by transcranial direct current stimulation (tDCS). *Front Behav Neurosci* 12:76.
 51. Gottfried JA, Dolan RJ (2004): Human orbitofrontal cortex mediates extinction learning while accessing conditioned representations of value. *Nat Neurosci* 7:1144.
 52. Becker B, Mihov Y, Scheele D, Kendrick KM, Feinstein JS, Matusch A, *et al.* (2012): Fear processing and social networking in the absence of a functional amygdala. *Biol Psychiatry* 72:70–77.
 53. Motzkin JC, Philippi CL, Wolf RC, Baskaya MK, Koenigs M (2015): Ventromedial prefrontal cortex is critical for the regulation of amygdala activity in humans. *Biol Psychiatry* 77:276–284.
 54. Braszko JJ, Walesiuk A, Wielgat P (2006): Cognitive effects attributed to angiotensin II may result from its conversion to angiotensin IV. *J Renin Angiotensin Aldosterone Syst* 7:168–174.
 55. Ball TM, Knapp SE, Paulus MP, Stein MB (2017): Brain activation during fear extinction predicts exposure success. *Depress Anxiety* 34:257–266.
 56. Milad MR, Rosenbaum BL, Simon NM (2014): Neuroscience of fear extinction: Implications for assessment and treatment of fear-based and anxiety related disorders. *Behav Res Ther* 62:17–23.
 57. Waters AM, Pine DS (2016): Evaluating differences in Pavlovian fear acquisition and extinction as predictors of outcome from cognitive behavioural therapy for anxious children. *J Child Psychol Psychiatry* 57:869–876.

58. Forcadell E, Torrents-Rodas D, Vervliet B, Leiva D, Tortella-Feliu M, Fullana MA (2017): Does fear extinction in the laboratory predict outcomes of exposure therapy? A treatment analog study. *Int J Psychophysiol* 121:63–71.
59. Marin M-F, Zsido RG, Song H, Lasko NB, Killgore WD, Rauch SL, *et al.* (2017): Skin conductance responses and neural activations during fear conditioning and extinction recall across anxiety disorders. *JAMA Psychiatry* 74:622–631.
60. Milad MR, Furtak SC, Greenberg JL, Keshaviah A, Im JJ, Falkenstein MJ, *et al.* (2013): Deficits in conditioned fear extinction in obsessive-compulsive disorder and neurobiological changes in the fear circuit. *JAMA Psychiatry* 70:608–618.
61. Milad MR, Pitman RK, Ellis CB, Gold AL, Shin LM, Lasko NB, *et al.* (2009): Neurobiological basis of failure to recall extinction memory in posttraumatic stress disorder. *Biol Psychiatry* 66:1075–1082.
62. Parrish JN, Bertholomey ML, Pang HW, Speth RC, Torregrossa MM (2019): Estradiol modulation of the renin-angiotensin system and the regulation of fear extinction. *Transl Psychiatry* 9:36.
63. Armando I, Carranza A, Nishimura Y, Hoe K-L, Barontini M, Terrón JA, *et al.* (2001): Peripheral administration of an angiotensin II AT1 receptor antagonist decreases the hypothalamic-pituitary-adrenal response to isolation stress. *Endocrinology* 142:3880–3889.