

# Impact of Body Mass Index on Outcomes in the Edoxaban Versus Warfarin Therapy Groups in Patients Underwent Cardioversion of Atrial Fibrillation (from ENSURE-AF)



Gregory Y.H. Lip, MD<sup>a,b,\*</sup>, Jose L. Merino, MD<sup>c</sup>, Maciej Banach, MD<sup>d</sup>, Joris R. de Groot, MD, PhD<sup>e</sup>, Lars S. Maier, MD<sup>f</sup>, Sakis Themistoclakis, MD<sup>g</sup>, Giuseppe Boriani, MD<sup>h</sup>, James Jin, PhD<sup>i</sup>, Michael Melino, MS, PhD<sup>i</sup>, Shannon M. Winters, MS<sup>i,1</sup>, and Andreas Goette, MD<sup>j,k</sup>

**In the Edoxaban versus warfarin in subjects Undergoing cardioversion of Atrial Fibrillation study (NCT 02072434), edoxaban showed similar efficacy and safety versus enoxaparin–warfarin in patients underwent electrical cardioversion of nonvalvular atrial fibrillation. In this ancillary analysis, we compared the primary efficacy (composite of stroke, systemic embolic event, myocardial infarction, cardiovascular death, and overall study period) and safety (composite of major and clinically relevant nonmajor bleeding, on-treatment) end points in relation to body mass index (BMI; <30 vs ≥30 kg/m<sup>2</sup>). We also compared cardioversion outcomes in relation to BMI. Of 2,199 patients enrolled, 1,095 were randomized to edoxaban and 1,104 to enoxaparin–warfarin. Mean age was 64 ± 10 and 64 ± 11 years and mean BMI 30.6 and 30.7 kg/m<sup>2</sup>, respectively. Cardiovascular and metabolic diseases were more prevalent in obese (n = 1067) than nonobese patients. Overall ischemic event rates were low; rates in the BMI <30 kg/m<sup>2</sup> subgroup were numerically lower than the ≥30 kg/m<sup>2</sup> subgroup, but not significantly different (odds ratio [OR], 0.74 [95% confidence interval 0.23, 2.24]). Composite major + clinically relevant nonmajor bleeding rates were low and numerically lower, but not significantly different (OR 0.88 [0.38, 2.04]), between the edoxaban and enoxaparin–warfarin arms and across weight categories. Successful cardioversion rate was higher in the BMI <30 versus ≥30 kg/m<sup>2</sup> subgroup (73.9% vs 69.9%; OR 1.22 [1.01 to 1.48]). In Edoxaban versus warfarin in subjects Undergoing cardioversion of Atrial Fibrillation, BMI did not significantly impact the relative efficacy and safety of edoxaban versus enoxaparin–warfarin. Nevertheless, the nonobese group had a higher rate of cardioversion success than the obese group. © 2018 Elsevier Inc. All rights reserved. (Am J Cardiol 2019;123:592–597)**

Obesity is a risk factor for all-cause and cardiovascular (CV) death in the general population; however, an inverse relation between being overweight or obese and

a better CV prognosis is observed, the so-called “obesity paradox.”<sup>1–4</sup> Our recent systematic review was suggestive of an obesity paradox in patients with atrial fibrillation (AF), particularly for all-cause and CV death outcomes.<sup>3</sup> In the nonvitamin K antagonist oral anticoagulant (NOAC) trials of stroke prevention in AF, an obesity paradox was also evident, with a treatment effect favoring NOACs over warfarin for both efficacy and safety that was significant only for normal-weight patients.<sup>3,5</sup> Nevertheless, there is uncertainty whether this obesity paradox is also evident for AF patients underwent rhythm control. Certainly, weight reduction is associated with better outcomes after rhythm control,<sup>6,7</sup> but limited prospective trial data are available. In the Edoxaban Versus Warfarin in Subjects Undergoing Cardioversion of Atrial Fibrillation (ENSURE-AF) trial, there were comparable low rates of major and clinically relevant nonmajor (CRNM) bleeding and thromboembolism when the oral factor Xa inhibitor edoxaban was compared with enoxaparin–warfarin.<sup>8</sup> This ancillary analysis from the ENSURE-AF trial compared clinical outcomes by body mass index (BMI, <30 vs ≥30 kg/m<sup>2</sup>).

<sup>a</sup>Liverpool Centre for Cardiovascular Science, University of Liverpool and Liverpool Heart & Chest Hospital, Liverpool, United Kingdom; <sup>b</sup>Aalborg Thrombosis Research Unit, Department of Clinical Medicine, Aalborg University, Aalborg, Denmark; <sup>c</sup>Hospital Universitario La Paz, Universidad Europea, Madrid, Spain; <sup>d</sup>Department of Hypertension, Medical University of Lodz, Lodz, Poland; <sup>e</sup>Amsterdam UMC, University of Amsterdam, Heart Center, Department of Cardiology, Amsterdam, The Netherlands; <sup>f</sup>Department of Internal Medicine II, University Hospital Regensburg, Regensburg, Germany; <sup>g</sup>Dell’Angelo Hospital, Venice-Mestre, Italy; <sup>h</sup>Cardiology Division, Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, Modena, Italy; <sup>i</sup>Daiichi Sankyo, Inc., Basking Ridge, New Jersey; <sup>j</sup>St. Vincenz-Hospital, Paderborn, Germany; and <sup>k</sup>Working Group: Molecular Electrophysiology, University Hospital Magdeburg, Magdeburg, Germany. Manuscript received September 1, 2018; revised manuscript received and accepted November 15, 2018.

Funding Source: The ENSURE-AF study was sponsored and funded by Daiichi Sankyo.

See page 597 for disclosure information.

<sup>1</sup>Employed at Daiichi Sankyo at the time the study was conducted.

\*Corresponding author: Tel: +44 151 794 9020.

E-mail address: [gregory.lip@liverpool.ac.uk](mailto:gregory.lip@liverpool.ac.uk) (G.Y.H. Lip).

## Methods

The design and principal results of the ENSURE-AF trial (NCT 02072434) have been published.<sup>8,9</sup> In brief, this was a multicenter, prospective, randomized, open, blinded end point trial in patients with nonvalvular AF underwent electrical cardioversion that compared edoxaban 60 mg once daily with enoxaparin–warfarin in 2,199 patients. Patients with an international normalized ratio (INR) <2.0 at randomization received enoxaparin bridging and daily warfarin until the INR was  $\geq 2.0$ . Those with INR  $\geq 2.0$  at the time of randomization did not require enoxaparin and were treated with warfarin alone; hence, edoxaban was compared with “optimized anticoagulation” with enoxaparin–warfarin.

The primary efficacy end point was the composite of stroke, systemic embolic event (SEE), myocardial infarction (MI), and CV death during the overall treatment period from randomization until end of study and the primary safety end point was the composite of major and CRNM bleeding during the on-treatment period (time of first dose to last dose of study drug taken). Successful cardioversion was confirmed by 12-lead electrocardiogram-documented sinus rhythm. The trial protocol was approved by ethics committees or institutional review boards. All patients provided written informed consent before participation in the study. This ancillary analysis compared the primary efficacy and safety end points with clinical outcomes by BMI (<30 vs  $\geq 30$  kg/m<sup>2</sup>).

Patients were followed for 28 days on study drug after cardioversion plus another 30 days to assess safety, which were analyzed in relation to body weight and BMI. For enoxaparin–warfarin patients, the clinical characteristics were summarized by BMIs of <30 and  $\geq 30$  kg/m<sup>2</sup>, with categorical variables as frequencies and percentages, and continuous variables as mean and standard deviation. Comparison of clinical characteristics for patients with BMI <30 and  $\geq 30$  kg/m<sup>2</sup> using the chi-square test for categorical variables and 1-way analysis of variance for continuous variables were provided.

The number and percent of patients with primary efficacy and safety outcomes were provided by treatment arm. Odds ratios (ORs) and 95% confidence intervals (CIs) are presented to assess the difference between treatment arms. We also explored outcomes in relation to BMI as a continuous variable. In addition, successful cardioversion in patients with BMI <30 kg/m<sup>2</sup> were compared with those with BMI  $\geq 30$  kg/m<sup>2</sup>. The number and percent of patients with successful cardioversion were provided by BMI category. ORs and 95% CIs are presented to assess the difference between BMI categories.

## Results

Of 2,199 patients enrolled, 1,095 were randomized to edoxaban and 1,104 to enoxaparin–warfarin. Mean  $\pm$  standard deviation age was 64.3  $\pm$  10 and 64.2  $\pm$  11 years and mean BMI 30.6 and 30.7 kg/m<sup>2</sup>, respectively. In all, 1,067 patients had a BMI of  $\geq 30$  kg/m<sup>2</sup>; in these obese patients, CV and metabolic diseases were more prevalent than in nonobese patients, as confirmed by the use of statins and

antihypertension medications (Table 1). Mean CHA<sub>2</sub>DS<sub>2</sub>-VASc (congestive heart failure, hypertension, age  $\geq 75$  years [2 points], diabetes mellitus, stroke [2 points], vascular disease, age 65–74 years, sex category) and HAS-BLED (hypertension, age, stroke, bleeding tendency/predisposition, labile INRs, elderly age/frailty, drugs such as concomitant aspirin/nonsteroidal anti-inflammatory drugs or alcohol excess) scores were significantly higher in the obese subgroup, suggesting they were at greater risk for stroke and bleeding. There were no relevant differences in time to therapeutic range and time in therapeutic range in relation to BMI <30 versus  $\geq 30$  kg/m<sup>2</sup>.

Rates of composite stroke/SEE, MI, or CV mortality rates were low and numerically lower for obese patients relative to nonobese patients, but were nonsignificant (OR 0.74 [0.23, 2.24]) even for both the edoxaban and enoxaparin–warfarin arms and across weight categories. Composite major and CRNM bleeding rates were low and numerically lower for obese patients relative to nonobese patients (OR 0.88 [0.38, 2.04]), as well as being nonsignificant in both the edoxaban and enoxaparin–warfarin arms and across weight categories. Major bleeding rates were numerically lower, but nonsignificant across weight categories (OR 0.32 [0.0, 1.8]).

Successful cardioversion was significantly more likely in those with BMI <30 kg/m<sup>2</sup> (OR 1.22 [1.01 to 1.48]; Table 2). Mean BMI was slightly lower in those with successful cardioversion compared with those with unsuccessful cardioversion (30.56 [standard deviation 5.71] vs 31.22 [5.45;  $p = 0.0472$ ]).

In a logistic regression analysis with the composite of major and CRNM bleeding as the response variable; and treatment, numerical BMI, and their interaction as independent variables; the  $p$  values for treatment, BMI, and interaction were 0.8852, 0.9016, and 0.9662, respectively (data not shown).

When comparing BMI <30 versus  $\geq 30$  kg/m<sup>2</sup>, composite ischemic events (stroke/SEE, MI, and CV mortality) were numerically lower, but given the low overall rates, this was nonsignificant (on-treatment analysis OR 0.74 [95% CI 0.23, 2.24]). In a logistic regression analysis with primary efficacy end point as the response variable; and treatment, numerical BMI, and their interaction as independent variables; the  $p$  values for treatment, BMI, and interaction were 0.0645, 0.2022, and 0.1034, respectively (data not shown).

Outcomes in relation to BMI as a continuous variable are shown in Figure 1. For major plus CRNM bleeding (on-treatment analysis), no relation was apparent between BMI and treatment with edoxaban or enoxaparin–warfarin. For stroke/SEE, MI, and CV mortality, there was a trend toward lower event rates with increasing BMI in the enoxaparin–warfarin group. For edoxaban, few events were seen at lower BMI values to show any trends, but no difference was observed when compared with enoxaparin–warfarin at higher BMI values. The proportion with successful cardioversion was higher in the BMI <30 kg/m<sup>2</sup> subgroup (827/1119; 73.9%) compared with the BMI  $\geq 30$  kg/m<sup>2</sup> subgroup (745/1,067; 69.9%; OR 1.22 [1.01 to 1.18]),  $p = 0.038$ . In a logistic regression analysis with successful cardioversion as the response variable; and treatment, numerical BMI, and

Table 1  
Baseline characteristics by BMI

Variable	BMI (kg/m <sup>2</sup> )								p value <sup>†</sup>
	Overall		<30			≥30			
	Edoxaban (N = 1095)	Enoxaparin– warfarin (N = 1104)	Edoxaban (n = 560)	Enoxaparin– warfarin (n = 559)	Total (n = 1119)	Edoxaban (n = 530)	Enoxaparin– warfarin (n = 537)	Total (n = 1067)	
Age, mean, SD	64.3 (10.3)	64.2 (10.8)	65.6 (11.1)	65.1 (11.3)	65.4 (11.2)	62.9 (9.3)	63.2 (10.1)	63.0 (9.7)	<0.0001
>65	509 (46.5%)	530 (48.0%)	293 (52.3%)	287 (51.3%)	580 (51.8%)	212 (40.0%)	239 (44.5%)	451 (42.3%)	
Weight, kg, mean (SD)	90.9 (18.3)	91.2 (19.0)	79.4 (11.3)	78.8 (11.4)	79.1 (11.3)	102.9 (16.5)	104.1 (16.6)	103.5 (16.6)	<0.0001
≤60	21 (1.9%)	33 (3.0%)	21 (3.8%)	33 (5.9%)	54 (4.8%)	0	0	0	
Anticoagulant experienced, n (%)									
Current* VKA user	513 (46.8)	558 (50.5)	257 (45.9)	298 (53.3)	555 (49.6)	254 (47.9)	259 (48.2)	513 (48.1)	0.4935
Current* NOAC user	157 (14.3)	148 (13.4)	78 (13.9)	70 (12.5)	148 (13.2)	77 (14.5)	78 (14.5)	155 (14.5)	0.3866
CrCl, mean (SD)	94.0 (35.7)	94.1 (34.7)	81.2 (29.2)	81.1 (27.9)	81.2 (28.6)	107.5 (37.1)	108.0 (35.9)	107.7 (36.5)	<0.0001
TtTR (days), mean (SD)	–	7.7 (5.1)	–	7.4 (4.9)	–	–	7.9 (5.4)	–	0.2428
TiTR (% of time), mean (SD)	–	70.8 (27.4%)	–	72.0 (26.9)	–	–	69.8 (28.0)	–	0.2153
TTR (% of time), mean (SD)	–	59.8 (30.6%)	–	60.5 (30.1)	–	–	59.3 (31.1)	–	0.5646
Heart failure	476 (43.5%)	484 (43.8%)	219 (39.1%)	239 (42.8%)	458 (40.9%)	256 (48.3%)	245 (45.6%)	501 (47.0%)	0.0051
Coronary artery disease	181 (16.5%)	197 (17.8%)	79 (14.1%)	93 (16.6%)	172 (15.4%)	102 (19.2%)	104 (19.4%)	206 (19.3%)	0.0174
Hypertension	850 (77.6%)	864 (78.3%)	399 (71.3%)	397 (71.0%)	796 (71.1%)	448 (84.5%)	464 (86.4%)	912 (85.5%)	<0.0001
Diabetes	218 (19.9%)	197 (17.8%)	80 (14.3%)	57 (10.2%)	137 (12.2%)	137 (25.8%)	139 (25.9%)	276 (25.9%)	<0.0001
Peripheral artery disease	40 (3.7%)	54 (4.9%)	25 (4.5%)	30 (5.4%)	55 (4.9%)	15 (2.8%)	24 (4.5%)	39 (3.7%)	0.1702
Valvular heart disease	250 (22.8%)	240 (21.7%)	133 (23.8%)	140 (25.0%)	273 (24.4%)	116 (21.9%)	100 (18.6%)	216 (20.2%)	0.0209
Intracranial haemorrhage	2 (0.2%)	3 (0.3%)	0	1 (0.2%)	1 (0.1%)	2 (0.4%)	2 (0.4%)	4 (0.4%)	0.2077
Ischemic stroke/transient ischaemic attack	68 (6.2%)	66 (6.0%)	41 (7.3%)	39 (7.0%)	80 (7.1%)	26 (4.9%)	27 (5.0%)	53 (5.0%)	0.0392
Myocardial infarction	69 (6.3%)	78 (7.1%)	37 (6.6%)	37 (6.6%)	74 (6.6%)	32 (6.0%)	41 (7.6%)	73 (6.8%)	0.8645
Life-threatening bleed	3 (0.3%)	3 (0.3%)	1 (0.2%)	1 (0.2%)	2 (<0.1%)	2 (0.4%)	2 (0.4%)	4 (0.4%)	0.4421
AF history									
Paroxysmal (≤7 days)	208 (19.0%)	207 (18.8%)	105 (18.8%)	115 (20.6%)	220 (19.7%)	103 (19.4%)	92 (17.2%)	195 (18.3%)	0.4450
Persistent (>7 days, <1 yr)	887 (81.0%)	890 (80.6%)	455 (81.3%)	444 (79.4%)	899 (80.3%)	427 (80.6%)	443 (82.8%)	870 (81.5%)	0.4450
CHA <sub>2</sub> DS <sub>2</sub> -VASc score, mean (SD)	2.6 (1.5)	2.6 (1.4)	2.6 (1.6)	2.5 (1.4)	2.5 (1.50)	2.7 (1.4)	2.7 (1.4)	2.7 (1.4)	0.0101
HAS-BLED Score, mean (SD)	0.9 (0.8)	0.9 (0.8)	0.9 (0.8)	0.9 (0.8)	0.9 (0.78)	0.8 (0.8)	0.9 (0.8)	0.9 (0.8)	0.0361
Drug therapies									
Aspirin	192 (17.5%)	221 (20.0%)	100 (17.9%)	105 (18.8%)	205 (18.3%)	92 (17.4%)	114 (21.2%)	206 (19.3%)	0.5841
Statins	429 (39.2%)	411 (37.2%)	211 (37.7%)	188 (33.6%)	399 (35.7%)	216 (40.8%)	220 (41.0%)	436 (40.9%)	0.0136
ACEI/ARB	692 (63.2%)	688 (62.3%)	322 (57.5%)	308 (55.1%)	630 (56.3%)	368 (69.4%)	376 (70.0%)	644 (60.4%)	<0.00010
Beta blockers	862 (78.7%)	847 (76.7%)	425 (75.9%)	434 (77.6%)	859 (76.8%)	434 (81.9%)	410 (76.4%)	844 (79.1%)	1975

ACEI = angiotensin-converting enzyme inhibitor; ARB = angiotensin receptor blocker; BMI = body mass index; CAD = coronary artery disease; CHA<sub>2</sub>DS<sub>2</sub>-VASc = congestive heart failure, hypertension, age ≥ 75 years (2 points), diabetes mellitus, stroke (2 points), vascular disease, age 65 to 74 years, sex category; CrCl = creatinine clearance; HAS-BLED = hypertension, age, stroke, bleeding tendency/predisposition, labile INRs, elderly age/frailty, drugs such as concomitant aspirin/nonsteroidal anti-inflammatory drugs or alcohol excess; HF = heart failure; ICH = intracerebral hemorrhage; MI = myocardial infarction; NOAC = nonvitamin K antagonist oral anticoagulant; PAD = peripheral arterial disease; SD = standard deviation; VHD = valvular heart disease; TIA = transient ischemic stroke; TiTR = time in therapeutic range (calculated from the first day with 2 ≤ INR ≤ 3); TTR = time in therapeutic range (calculated from day 8 of study drug); TtTR = time to achieve therapeutic range; VKA = vitamin K antagonist.

\* Current defined as using VKA or NOAC at randomization or within 30 days before randomization. Percentages are based on the numbers of anticoagulant experienced.

<sup>†</sup> Comparisons between total columns for BMI < 30 and BMI ≥ 30.

Table 2  
Event rates by BMI

	Overall		BMI < 30 kg/m <sup>2</sup>		BMI ≥ 30 kg/m <sup>2</sup>	
	Edoxaban	Enoxaparin–warfarin	Edoxaban	Enoxaparin–warfarin	Edoxaban	Enoxaparin–Warfarin
First stroke, SEE, MI, or CV mortality*						
N	1095	1104	560	559	530	537
n (%)	5 (0.5%)	11 (1.0%)	1 (0.2%)	6 (1.1%)	4 (0.8%)	5 (0.9%)
OR (95% CI)	0.46 (0.12, 1.43)		0.17 (0, 1.37)		0.81 (0.16, 3.78)	
Major or CRNM bleeding events <sup>†</sup>						
N	1067	1082	547	551	517	528
n (%)	16 (1.5%)	11 (1.0%)	8 (1.5%)	5 (0.9%)	8 (1.6%)	6 (1.1%)
OR (95% CI)	1.48 (0.64, 3.55)		1.62 (0.46, 6.34)		1.37 (0.41, 4.82)	
Major bleeding events <sup>†</sup>						
N	1067	1082	547	551	517	528
n (%)	3 (0.3%)	5 (0.5%)	1 (0.2%)	1 (0.2%)	2 (0.4%)	4 (0.8%)
OR (95% CI)	0.61 (0.09, 3.13)		1.01 (0.01, 79.21)		0.51 (0.05, 3.57)	
Successful cardioversion						
N	1095 <sup>‡</sup>	1104 <sup>§</sup>	560	559	530	537
n (%)	790 (72.2%)	788 (71.4%)	414 (73.9%)	413 (73.9%)	374 (70.6%)	372 (69.3%)
OR (95% CI)	1.04 (0.86, 1.26)		1.00 (0.76, 1.32)		1.06 (0.81, 1.39)	
			BMI <30 kg/m <sup>2</sup>		BMI ≥30 kg/m <sup>2</sup>	
First stroke, SEE, MI, or CV mortality*						
N			1119		1067	
n (%)			7 (0.6%)		9 (0.8%)	
OR (95% CI)			0.74 (0.23, 2.24)			
Major or CRNM bleeding events <sup>†</sup>						
N			1098		1045	
n (%)			13 (1.2%)		14 (1.3%)	
OR (95% CI)			0.88 (0.38, 2.04)			
Major bleeding events <sup>†</sup>						
N			1098		1045	
n (%)			2 (0.2%)		6 (0.6%)	
OR (95% CI)			0.32 (0.0, 1.8)			
Successful cardioversion <sup>‡</sup>						
N			1119		1067	
n (%)			827 (73.9%)		746 (69.9%)	
OR (95% CI)			1.22 (1.01, 1.48)			

BMI = body mass index; CI = confidence interval; CRNM = clinically relevant nonmajor; CV = cardiovascular; ITT = intention to treat; MI = myocardial infarction; OR = odds ratio; SEE = systemic embolic event.

\* ITT population, overall study period (28 days on study drug after cardioversion + 30 days follow-up).

<sup>†</sup> All treated patients, on-treatment period (time of first dose to last dose of study drug taken).

<sup>‡</sup> Data for BMI was not available for 5 patients.

<sup>§</sup> Data for BMI was not available for 8 patients.

their interaction as independent variables; the p values for treatment, BMI, and interaction are 0.7168, 0.2265, and 0.6644, respectively.

## Discussion

In this ancillary analysis from ENSURE-AF, the data suggest that obesity does not influence the rate of ischemic events after cardioversion regardless of the therapeutic strategy. The BMI <30 kg/m<sup>2</sup> group had a higher rate of cardioversion success than the BMI ≥30 kg/m<sup>2</sup> group; and edoxaban had comparable efficacy and safety to optimized usual anticoagulation with enoxaparin–warfarin, and were not significantly different in various BMI categories.

Our systematic review found that only obese patients were at lower risk for major bleeding compared with normal-weight patients.<sup>3</sup> In the present analysis from ENSURE-AF, no significant relation was evident between the primary bleeding outcome and BMI. The present patient population was at low bleeding risk, as evident by a mean HAS-BLED score of 0.9. While guidelines advocate focus on modifiable bleeding risk factors, recent evidence shows that the HAS-BLED score is a better assessment of the AF patient's potential bleeding risk compared with simply using modifiable bleeding risk factors.<sup>10–12</sup>

In a previous systematic review and meta-analysis, we found that there may be an obesity paradox in AF patients for all-cause and CV death outcomes.<sup>3</sup> An obesity paradox was also seen for stroke/SEEs, with a treatment effect




Figure 1. Event rate at 1 month of primary efficacy outcomes and BMI for (A) Stroke, SEE, MI, and CV and (B) major or CRNM bleeding. (a) Stroke, SEE, MI, and CV mortality, ITT analysis set—overall period; (b) Stroke, SEE, MI, and CV mortality, safety analysis set—on treatment period.

favoring NOACs over warfarin for both efficacy and safety that was significant only for normal-weight patients. In the present analysis from ENSURE-AF, obesity did not influence the rate of the composite efficacy events (stroke/SEE, MI, and CV mortality) after cardioversion regardless of treatment with NOAC or enoxaparin–warfarin. This is despite the ENSURE-AF trial including a relatively high-risk patient population for stroke (mean CHA<sub>2</sub>DS<sub>2</sub>VASc score 2.6), that was broadly comparable to the patient population in the ENGAGE-AF trial<sup>13</sup> (mean CHADS<sub>2</sub> score 2.8) and other NOAC stroke prevention trials.<sup>14</sup> Nonetheless, the follow-up duration in ENSURE-AF was shorter than that in the ENGAGE-AF trial.

As expected from previous studies,<sup>15,16</sup> cardioversion success was lower in obese patients. This may reflect associated co-morbidities or a greater body impedance relevant to ENSURE-AF since electrical cardioversion was the only modality used. Indeed, pharmacological cardioversion is perhaps more advocated in obese subjects with AF.

Although the ENSURE-AF trial is the largest study in AF pericardioversion to date, this study is limited by being a subgroup analysis of a selected clinical trial cohort, and the results may not be applicable to the general AF population. BMI measurement and categorization of obesity was based on baseline measurements, and changes in BMI over time were not considered. Also, the low overall event rates

and short follow-up period may have influenced outcome event rates, which may be underpowered.

In conclusion, edoxaban had efficacy and safety comparable with optimized standard anticoagulation with enoxaparin-warfarin; neither treatment group showed significant differences in various BMI categories. Obesity did not influence the rate of ischemic events after cardioversion regardless of the therapeutic strategy. Nevertheless, the BMI <30 kg/m<sup>2</sup> group had a higher rate of cardioversion success than the obese group.

## Disclosure

GYH Lip reports consultancy and speaker fees from Bayer, Bayer/Janssen, BMS/Pfizer, Biotronik, Medtronic, Boehringer Ingelheim, Microlife, Roche, and Daiichi Sankyo outside the submitted work; no fees received personally. JL Merino reports personal fees from Abbott, Bayer, Biotronik, Boston Scientific, Bristol-Myers Squibb, Cardione, Daiichi Sankyo LivaNova, Medtronic, Pfizer, and Sanofi outside the submitted work. M Banach has served as a consultant for Abbott Vascular, Akcea, Amgen, Esperion, Eli Lilly, Merck Sharp & Dohme, Resverlogix, and Sanofi-Aventis; a speaker for Abbott/Mylan, Abbott Vascular, Actavis, Akcea, Amgen, Biofarm, KRKA, Merck Sharp & Dohme, Sanofi-Aventis, and Valeant; and reports grants from Sanofi-Aventis and Valeant. JR de Groot served as a consultant for AtriCure, Boehringer Ingelheim, Bristol-Myers Squibb, Daiichi Sankyo, Medtronic, and Pfizer; as a speaker for BackBeat Medical and Daiichi Sankyo; and has received research grants from AtriCure and St. Jude Medical. LS Maier reports speaker fees from Bayer, BMS/Pfizer, Boehringer Ingelheim, and Daiichi Sankyo. S Themistoclakis has served as a consultant for Bayer, Boehringer Ingelheim, Bristol-Myers Squibb, Daiichi Sankyo, and Pfizer. G Boriani reported speaker's fees of small amounts from Biotronik, Boehringer, Boston, and Medtronic. J Jin and M Melino are employees of Daiichi Sankyo; SM Winters was an employee of Daiichi Sankyo at the time of writing. A Goette reports personal fees from AstraZeneca, Bayer, Berlin-Chemie, Boehringer Ingelheim, Bristol-Myers Squibb, Daiichi Sankyo, Medtronic, Pfizer, and Sanofi-Aventis.

## Acknowledgment

Editorial support was provided by AlphaBioCom, LLC (King of Prussia, Pennsylvania) and funded by Daiichi Sankyo, Inc. (Basking Ridge, New Jersey). The first draft was written by GYHL and all authors contributed to drafting and revisions for the finalized version.

1. Proietti M, Lane DA, Lip GY. Relation of nonvalvular atrial fibrillation to body mass index (from the SPORTIF Trials). *Am J Cardiol* 2016;118:72–78.
2. Boriani G, Laroche C, Diemberger I, Fantecchi E, Meeder J, Kurpesa M, Baluta MM, Proietti M, Tavazzi L, Maggioni AP, Lip GYH. Invest-

3. Proietti M, Guiducci E, Cheli P, Lip GY. Is There an obesity paradox for outcomes in atrial fibrillation? A systematic review and meta-analysis of non-vitamin k antagonist oral anticoagulant trials. *Stroke* 2017;48:857–866.
4. Alagiakrishnan K, Banach M, Ahmed A, Aronow WS. Complex relationship of obesity and obesity paradox in heart failure - higher risk of developing heart failure and better outcomes in established heart failure. *Ann Med* 2016;48:603–613.
5. De Caterina R, Lip GYH. The non-vitamin K antagonist oral anticoagulants (NOACs) and extremes of body weight—a systematic literature review. *Clin Res Cardiol* 2017;106:565–572.
6. Pathak RK, Middeldorp ME, Meredith M, Mehta AB, Mahajan R, Wong CX, Twomey D, Elliott AD, Kalman JM, Abhayaratna WP, Lau DH, Sanders P. Long-term effect of goal directed weight management in an atrial fibrillation cohort: a Long-term Follow-Up Study (LEGACY Study). *J Am Coll Cardiol* 2015 May 26;65(20):2159–2169.
7. Wong CX, Sullivan T, Sun MT, Mahajan R, Pathak RK, Middeldorp M, Twomey D, Ganesan AN, Rangnekar G, Roberts-Thomson KC, Lau DH, Sanders P. Obesity and the risk of incident, post-operative, and post-ablation atrial fibrillation: a meta-analysis of 626,603 individuals in 51 studies. *JACC: Clin Electrophysiol* 2015;1:139–152.
8. Goette A, Merino JL, Ezekowitz MD, Zamoryakhin D, Melino M, Jin J, Mercuri MF, Grosso MA, Fernandez V, Al-Saady N, Pelekh N, Merkely B, Zenin S, Kushnir M, Spinar J, Batushkin V, de Groot JR, Lip GY. investigators E-A. Edoxaban versus enoxaparin-warfarin in patients undergoing cardioversion of atrial fibrillation (ENSURE-AF): a randomised, open-label, phase 3b trial. *Lancet* 2016;388:1995–2003.
9. Lip GY, Merino J, Ezekowitz M, Ellenbogen K, Zamoryakhin D, Lanz H, Jin J, Al-Saadi N, Mercuri M, Goette A. A prospective evaluation of edoxaban compared to warfarin in subjects undergoing cardioversion of atrial fibrillation: the Edoxaban vs. warfarin in subjectS Undergoing cardioversion of Atrial Fibrillation (ENSURE-AF) study. *Am Heart J* 2015;169:597–604. e595.
10. Chao TF, Lip GYH, Lin YJ, Chang SL, Lo LW, Hu YF, Tuan TC, Liao JN, Chung FP, Chen TJ, Chen SA. Major bleeding and intracranial hemorrhage risk prediction in patients with atrial fibrillation: attention to modifiable bleeding risk factors or use of a bleeding risk stratification score? A nationwide cohort study. *Int J Cardiol* 2018;254:157–161.
11. Guo Y, Zhu H, Chen Y, Lip GYH. Comparing bleeding risk assessment focused on modifiable risk factors only versus validated bleeding risk scores in atrial fibrillation. *Am J Med* 2018;131:185–192.
12. Esteve-Pastor MA, Rivera-Caravaca JM, Shantsila A, Roldan V, Lip GYH, Marin F. Assessing bleeding risk in atrial fibrillation patients: comparing a bleeding risk score based only on modifiable bleeding risk factors against the HAS-BLED Score. The AMADEUS Trial. *Thromb Haemost* 2017;117:2261–2266.
13. Giugliano RP, Ruff CT, Braunwald E, Murphy SA, Wiviott SD, Halperin JL, Waldo AL, Ezekowitz MD, Weitz JI, Spinar J, Ruzyllo W, Ruda M, Koretsune Y, Betcher J, Shi M, Grip LT, Patel SP, Patel I, Hanyok JJ, Mercuri M, Antman EM. ENGAGE AF-TIMI 48 Investigators. Edoxaban versus warfarin in patients with atrial fibrillation. *N Engl J Med* 2013;369:2093–2104.
14. Ruff CT, Giugliano RP, Braunwald E, Hoffman EB, Deenadayalu N, Ezekowitz MD, Camm AJ, Weitz JI, Lewis BS, Parkhomenko A, Yamashita T, Antman EM. Comparison of the efficacy and safety of new oral anticoagulants with warfarin in patients with atrial fibrillation: a meta-analysis of randomised trials. *Lancet* 2014;383:955–962.
15. Blich M, Edoute Y. Electrical cardioversion for persistent or chronic atrial fibrillation: outcome and clinical factors predicting short and long term success rate. *Int J Cardiol* 2006;107:389–394.
16. Frick M, Frykman V, Jensen-Urstad M, Ostergren J, Rosenqvist M. Factors predicting success rate and recurrence of atrial fibrillation after first electrical cardioversion in patients with persistent atrial fibrillation. *Clin Cardiol* 2001;24:238–244.