

## Impact of Brief Nicotine Messaging on Nicotine-Related Beliefs in a U.S. Sample



Andrea C. Villanti, PhD, MPH,<sup>1</sup> Julia C. West, BA,<sup>1</sup> Darren Mays, PhD, MPH,<sup>2</sup> Eric C. Donny, PhD,<sup>3</sup> Joseph N. Cappella, PhD,<sup>4</sup> Andrew A. Strasser, PhD<sup>5</sup>

**Introduction:** The current study pilot tested the effect of a single, brief exposure to nicotine education messages on beliefs about nicotine, nicotine-replacement therapy (NRT), E-cigarettes, and cigarettes with reduced nicotine content (RNC).

**Methods:** Five hundred and twenty-one U.S. adults (aged  $\geq 18$  years) completed a 15-minute survey on Amazon Mechanical Turk in 2018. After completing items on sociodemographics, literacy, and cancer risk behaviors, participants were randomized in a 2:1:1 ratio to 1 of 3 conditions: nicotine education ( $n=263$ ), sun safety education (attention control,  $n=128$ ), or no message control ( $n=130$ ). All participants completed items regarding nicotine, NRT, E-cigarette, and RNC cigarette beliefs, as well as norms about nicotine use, behavioral control regarding cigarette/tobacco use, and intention to use cigarettes, NRT, E-cigarettes, and RNC cigarettes in the next 12 months. Analyses were conducted in 2019.

**Results:** Following exposure, nicotine education participants reported fewer false beliefs about nicotine ( $p<0.001$ ), NRT ( $p<0.001$ ), E-cigarettes ( $p<0.05$ ), and RNC cigarettes ( $p<0.05$ ) compared with the control conditions. Nicotine messaging doubled the probability of a correct response (false, 78.3% vs 36.8%) to *nicotine is a cause of cancer* and dramatically reduced the probability of responding *don't know* to this item (5.3% vs 26.0%). There was no impact of the intervention on beliefs about other substances within cigarette, norms, or behavioral intentions.

**Conclusions:** Findings from the current study support the hypothesis that a brief nicotine messaging intervention—similar to the messages likely to be seen on warning labels or in media campaigns—is likely to correct misperceptions of nicotine, NRT, E-cigarettes, and RNC cigarettes.

*Am J Prev Med 2019;57(4):e135–e142. © 2019 American Journal of Preventive Medicine. Published by Elsevier Inc. All rights reserved.*

## INTRODUCTION

Authoritative reviews of carcinogens in tobacco and tobacco smoke have not listed nicotine among the carcinogens,<sup>1–4</sup> and syntheses of evidence conclude that combustion compounds in tobacco smoke are the primary contributors to the cardiovascular risk of tobacco use.<sup>5,6</sup> However, population studies have quantified widespread misperceptions of nicotine<sup>7–9</sup> with some smokers equating the harms of using U.S. Food and Drug Administration–approved nicotine-replacement therapy (NRT) for smoking cessation with the harms of cigarette smoking.<sup>10–16</sup>

Findings from RCTs support that cigarettes with reduced nicotine content (RNC) can reduce the use of

cigarettes per day, and exposure to and dependence on nicotine, with minimal smoking compensation among

From the <sup>1</sup>Vermont Center on Behavior and Health, Department of Psychiatry, University of Vermont Larner College of Medicine, Burlington, Vermont; <sup>2</sup>Department of Oncology, Georgetown University Medical Center, Cancer Prevention and Control Program, Lombardi Comprehensive Cancer Center, Washington, District of Columbia; <sup>3</sup>Department of Physiology and Pharmacology, Wake Forest School of Medicine, Winston-Salem, North Carolina; <sup>4</sup>Annenberg School for Communication at the University of Pennsylvania, Philadelphia, Pennsylvania; and <sup>5</sup>Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania

Address correspondence to: Andrea C. Villanti, PhD, MPH, Vermont Center on Behavior and Health, University of Vermont, 1 South Prospect Street, MS 482, Burlington VT 05401. E-mail: [avillanti@uvm.edu](mailto:avillanti@uvm.edu).

0749-3797/\$36.00

<https://doi.org/10.1016/j.amepre.2019.05.015>

users.<sup>17,18</sup> The Food and Drug Administration is considering a nicotine reduction standard in cigarettes, but has not described how consumer education on nicotine would be used to support the intended effect of this policy on current tobacco users, nonusers, or the population overall.<sup>19</sup>

The goal of the current study was to pilot test the effect of a single, brief exposure to nicotine educational messages on beliefs about nicotine, NRT, E-cigarettes, and RNC cigarettes in a convenience sample of adults.

## METHODS

### Study Sample

The authors conducted an online trial in 521 U.S. adults (aged  $\geq 18$  years) who completed a 15-minute survey on Amazon Mechanical Turk in 2018 on Communicating About Cancer Risk Behaviors. After completing items on sociodemographics, literacy,<sup>20</sup> and cancer risk behaviors (e.g., physical activity), participants were randomized in a 2:1:1 ratio to 1 of 3 conditions: nicotine messaging ( $n=263$ ), sun safety messaging (attention control,  $n=128$ ), or no message control ( $n=130$ ). Participants in the no message control condition immediately completed outcome measures. Participants in the 2 messaging intervention conditions completed these items after exposure to the educational messages. This study was deemed exempt by the IRB at the University of Vermont.

### Intervention

Six images were presented to participants in the nicotine messaging condition using a black slide template with smoke and content adapted for a lay audience from several evidence-based sources.<sup>1–3,5,21,22</sup> The 6 tested messages were: (1) *nicotine is the addictive substance in tobacco products*, (2) *nicotine makes it easier for people to start smoking regularly*, (3) *nicotine makes it harder for people to quit smoking*, (4) *nicotine does not cause cancer*, (5) *chemicals in cigarette smoke, not nicotine, largely cause cancer, heart disease, and other health problems related to smoking*, and (6) *nicotine can be used safely long-term in quit smoking products like nicotine patches, gum, or lozenges*. Participants in the sun safety condition also received 6 messages of similar length to the nicotine messages using an orange slide template with the image of a sun; the messages included *indoor tanning and ultraviolet radiation from the sun cause skin cancer and premature aging and wearing sunscreen alone does not prevent skin cancer*.

### Measures

Primary outcomes ascertained beliefs about nicotine, NRT, E-cigarettes, and RNC cigarettes. Secondary outcomes assessed norms about nicotine use, behavioral control regarding cigarette/tobacco use, and intention to use cigarettes, NRT, E-cigarettes, and RNC cigarettes in the next 12 months. These measures are detailed, with their response options, in Tables 1 and 2. Items on the relative harm of E-cigarettes or nicotine products compared with cigarettes were initially asked on a 5-point scale (*much less harmful* to *much more harmful*) but collapsed to a 3-point scale. Nine items on RNC cigarette beliefs were adapted from previous studies<sup>23,24</sup> and assessed on a 5-point scale from *definitely not true* to *definitely true*. Items were summed to create subscales, with higher scale values indicating a greater number of false beliefs. Norms

items on the social acceptability of specific tobacco products and other substances were assessed on a 5-point scale (*not at all to extremely*) and items on people's opinions of using nicotine on a 5-point scale (*very positive* to *very negative*). Acceptability of uses of nicotine was assessed by ranking 3 options from *most acceptable* (1) to *least acceptable* (3). Intention to use specific products in the next 12 months was assessed in past 30-day tobacco users and nonusers, with those reporting *definitely yes*, *probably yes*, and *probably not* coded as susceptible to future use and those reporting *definitely not* coded as not susceptible, in line with other studies.<sup>25,26</sup>

### Statistical Analysis

Bivariate analyses examined, in 2019, using chi-squared tests and *t*-tests, differences in sociodemographic characteristics (age, gender, race/ethnicity, education, and subjective financial situation), past 30-day tobacco use, and response to nicotine, NRT, E-cigarette, and RNC cigarette beliefs, norms, behavioral control, and intention to use by study condition. Because there were no significant differences in the primary outcomes between the 2 control conditions, comparisons focused on the nicotine messaging versus combined control conditions. Multiple linear regression analyses examined the relationship between study conditions and the 4 false beliefs scales, controlling for past 30-day tobacco use status.

## RESULTS

Approximately half of participants were male (52%), 46% were aged 25–34 years, 80% were white, 11% were of Hispanic ethnicity, 87% had at least some college education, and 40% reported past 30-day tobacco or E-cigarette use (Table 3). The study groups did not differ on pre-exposure measures of sociodemographic characteristics, literacy, or past 30-day tobacco use.

Table 1 shows a strong effect of nicotine messaging on reducing false beliefs about nicotine, NRT, E-cigarettes, and RNC cigarettes compared with the combined control conditions. Importantly, the nicotine messaging condition doubled the probability of a correct response (false, 78.3% vs 36.8%) to *nicotine is a cause of cancer* and dramatically reduced the probability of responding *don't know* to this item (5.3% vs 26.0%). It also increased correct responses regarding the contribution of nicotine to health risks and cancer caused by cigarette smoking ( $p < 0.001$ ). Of particular interest, the impact of the educational intervention was specific to nicotine; there was no impact on beliefs about other substances within a cigarette ( $p > 0.40$  for all). In multivariable models, exposure to nicotine messaging remained associated with a lower level of nicotine ( $b = -1.82$ ,  $p < 0.001$ ), NRT ( $b = -1.16$ ,  $p < 0.001$ ), and E-cigarette ( $b = -0.39$ ,  $p = 0.043$ ) false beliefs, after controlling for past 30-day tobacco use; the relationship between study condition and RNC cigarette false beliefs in this model was marginally significant ( $b = -1.13$ ,  $p = 0.054$ ).

**Table 1.** Differences in Nicotine and Cigarette Beliefs, by Study Condition

Beliefs	Study condition		p-value
	Nicotine messaging (n=263)	Combined controls (n=258)	
Thinking about the harm that individual substances within a cigarette may cause, how much harm comes from <sup>a</sup>			
Substances produced when raw tobacco burns? (missing=25)	3.30 (1.31)	3.21 (1.26)	0.44
The nicotine in a cigarette? (missing=23)	2.34 (1.36)	3.13 (1.29)	<0.001
Naturally occurring substances in tobacco? (missing=23)	2.66 (1.23)	2.70 (1.21)	0.69
Things that are added to cigarettes during the manufacturing process? (missing=23)	3.94 (1.11)	3.91 (1.13)	0.79
Nicotine false beliefs			
Nicotine is a cause of cancer <sup>b</sup>			<0.001
False	78.3	36.8	
Don't know	5.3	26.0	
True	16.4	37.2	
In your opinion, how large a part of the health risks of cigarette smoking comes from the nicotine itself? <sup>b</sup>			<0.001
None/small part	76.4	55.8	
Large/very large part	23.6	44.2	
In your opinion, how large a part of the cancer caused by cigarette smoking comes from the nicotine itself? <sup>b</sup>			<0.001
None/small part	84.0	62.8	
Large/very large part	16.0	37.2	
Nicotine false beliefs scale ( $\alpha=0.86$ ) <sup>a,c</sup>	4.90 (2.06)	6.71 (2.48)	<0.001
NRT false beliefs			
It is easy to get addicted to nicotine gum <sup>b</sup>			0.321
False	13.3	11.6	
Don't know	26.2	32.2	
True	60.5	56.2	
Long-term use of nicotine from patches or gums is almost as harmful to health as cigarette smoking <sup>b</sup>			<0.001
False	59.3	33.7	
Don't know	21.3	24.4	
True	19.4	41.9	
Are nicotine products (like gum, patches, lozenges) more likely, about the same, or less likely to cause someone to become addicted as regular cigarettes? <sup>b</sup>			0.016
Less likely	40.7	29.1	
About the same	47.9	55.0	
More likely	11.4	15.9	
Are nicotine products (like gum, patches, lozenges) more likely, about the same, or less likely to cause someone to have a heart attack as regular cigarettes? <sup>b</sup> (missing=1)			<0.001
Less likely	62.7	44.7	
About the same	27.8	44.4	
More likely	9.5	10.9	
Are nicotine products (like gum, patches, lozenges) more likely, about the same, or less likely to cause cancer as regular cigarettes? <sup>b</sup>			0.001
Less likely	72.2	56.2	
About the same	22.1	32.9	
More likely	5.7	10.9	
Relative harm of nicotine products (like gum, patches, lozenges) compared with cigarettes <sup>b</sup>			0.010

(continued on next page)

**Table 1.** Differences in Nicotine and Cigarette Beliefs, by Study Condition (*continued*)

Beliefs	Study condition		p-value
	Nicotine messaging (n=263)	Combined controls (n=258)	
Less harmful	76.4	64.3	
About the same	16.0	24.0	
More harmful	7.6	11.6	
NRT false beliefs scale ( $\alpha=0.74$ ) <sup>a,d</sup>	9.89 (2.63)	11.07 (2.84)	<b>&lt;0.001</b>
E-cigarette false beliefs			
Long-term use of E-cigarettes is almost as harmful to health as cigarette smoking <sup>b</sup>			<b>0.022</b>
False	35.0	30.6	
Don't know	27.8	20.5	
True	37.3	48.8	
Are E-cigarettes more likely, about the same, or less likely to cause someone to have a heart attack as regular cigarettes? <sup>b</sup> (missing=1)			0.059
Less likely	52.9	42.8	
About the same	38.4	48.2	
More likely	8.7	8.9	
Are E-cigarettes more likely, about the same, or less likely to cause cancer as regular cigarettes? <sup>b</sup> (missing=1)			<b>0.038</b>
Less likely	57.0	47.1	
About the same	34.2	45.1	
More likely	8.7	7.8	
Relative harm of E-cigarettes (like JUUL, Vuse, MarkTen, blu, or Joyetech) compared with cigarettes <sup>b</sup> (missing=2)			0.463
Less harmful	61.3	56.6	
About the same	28.4	33.3	
More harmful	10.3	10.1	
E-cigarette false beliefs scale ( $\alpha=0.79$ ) <sup>a,e</sup>	6.58 (2.21)	6.97 (2.24)	<b>0.043</b>
RNC cigarette false beliefs			
Cigarettes that are lower in nicotine are less likely to cause cancer than regular cigarettes <sup>a</sup>	2.05 (1.07)	2.29 (1.01)	<b>0.010</b>
Cigarettes that are lower in nicotine are safer than regular cigarettes <sup>a</sup>	2.16 (1.11)	2.32 (1.09)	0.110
Cigarettes that are lower in nicotine are healthier than regular cigarettes <sup>a</sup>	2.10 (1.09)	2.24 (1.14)	0.137
Cigarettes that are lower in nicotine have fewer chemicals than regular cigarettes <sup>a</sup>	2.14 (1.11)	2.24 (1.12)	0.289
Smoking cigarettes that are lower in nicotine make it easier to quit smoking completely compared with regular cigarettes <sup>a,f</sup>	2.81 (1.13)	3.03 (1.10)	<b>0.024</b>
Cigarettes that are lower in nicotine also have less tar than regular cigarettes <sup>a</sup>	2.40 (1.03)	2.38 (1.03)	0.810
High nicotine content cigarettes are worse for your health than low nicotine cigarettes, even if you smoke the same number of each <sup>a</sup>	2.63 (1.16)	2.82 (1.13)	0.053
A low nicotine cigarette is safer to smoke than a high nicotine cigarette, even if you don't quit <sup>a</sup>	2.35 (1.14)	2.41 (1.08)	0.545
Low nicotine cigarettes are healthier for you than high nicotine cigarettes even before you quit <sup>a</sup>	2.53 (1.16)	2.44 (1.09)	0.372
RNC cigarette false beliefs scale ( $\alpha=0.91$ ) <sup>a,g</sup>	20.99 (6.80)	22.16 (6.53)	<b>0.047</b>

Note: If missing number not provided, there are no missing data on that item. Boldface indicates statistical significance ( $p<0.05$ ).

<sup>a</sup>M (SD).

<sup>b</sup>Column percent.

<sup>c</sup>Nicotine false beliefs scale comprised 3 items (listed above; range 3–11).

<sup>d</sup>NRT false beliefs scale comprised 6 items (listed above; range 6–18).

<sup>e</sup>E-cigarette false beliefs scale comprised 4 items (listed above; range 3–12).

<sup>f</sup>This item was reverse-coded.

<sup>g</sup>RNC cigarette false beliefs scale comprised 9 items (listed above; range 9–39).

NRT, nicotine-replacement therapy; RNC, reduced nicotine content.

**Table 2.** Norms, Behavioral Control, and Intention to Use Nicotine and Tobacco, by Study Condition

Variable	Nicotine messaging (n=263)	Combined controls (n=258)	p-value
Norms, full sample	(n=263)	(n=258)	
How socially acceptable among your peers do you think each of the following products are? <sup>a</sup>			
Nicotine	2.45 (1.10)	2.60 (1.14)	0.132
Caffeine	4.53 (0.79)	4.51 (0.86)	0.776
Alcohol	3.94 (0.97)	4.02 (1.04)	0.410
Marijuana	3.04 (1.23)	3.25 (1.19)	<b>0.049</b>
Cigarettes	2.48 (1.16)	2.47 (1.22)	0.924
E-cigarettes	2.80 (1.17)	3.00 (1.26)	0.059
Nicotine products (i.e., gum, patches, lozenges)	2.89 (1.23)	2.83 (1.30)	0.635
Hookah	2.63 (1.23)	2.77 (1.26)	0.187
Low nicotine cigarettes	2.33 (1.14)	2.42 (1.14)	0.400
Rank the following 3 uses of nicotine in terms of their acceptability to you and people like you <sup>a</sup> (range 1–3)			
Nicotine delivered via the patch for cessation of tobacco use (missing=78)	1.76 (0.86)	1.90 (0.86)	0.086
Nicotine delivered via the E-cigarettes for either cessation or harm reduction (missing=78)	1.89 (0.67)	1.85 (0.69)	0.533
Nicotine delivered via E-cigarettes for purposes other than cessation or harm reduction (i.e., recreational use of E-cigarettes) (missing=78)	2.35 (0.80)	2.25 (0.84)	0.196
Opinion of using nicotine <sup>a</sup>			
Most people	3.78 (0.90)	3.79 (0.86)	0.845
People who are important to you	3.74 (1.01)	3.80 (0.96)	0.483
Behavioral control among past 30–day tobacco users	(n=97)	(n=109)	
How confident are you that you could resist smoking a cigarette in situations where others are smoking? <sup>b</sup>			0.722
Not at all confident	23.7	23.9	
Somewhat confident	33.0	36.7	
Moderately confident	21.7	15.6	
Very confident	21.6	23.9	
How confident are you that you can quit smoking cigarettes/using tobacco products totally and for good if and when you wanted to? <sup>b</sup>			0.789
Not at all confident	19.6	19.3	
Somewhat confident	42.3	41.3	
Moderately confident	17.5	13.8	
Very confident	20.6	25.7	
If a tobacco product made a claim that it was less harmful to health than other tobacco products, how likely would you be to use that product? <sup>b</sup>			0.439
Very likely	16.5	10.1	
Somewhat likely	32.0	34.9	
Somewhat unlikely	24.7	25.7	
Very unlikely	16.5	22.9	
Don't know	10.3	6.4	
Intention to use among past 30–day tobacco users	(n=97)	(n=109)	
Cigarettes <sup>b</sup>			0.768
No	12.4	13.8	
Yes	87.6	86.2	
E-cigarettes <sup>b</sup>			0.539
No	9.3	11.9	
Yes	90.7	88.1	

(continued on next page)

**Table 2.** Norms, Behavioral Control, and Intention to Use Nicotine and Tobacco, by Study Condition (*continued*)

Variable	Nicotine messaging	Combined controls	p-value
Low nicotine cigarettes <sup>b</sup>			0.597
No	21.7	24.8	
Yes	78.3	75.2	
NRT <sup>b</sup>			0.441
No	23.7	28.4	
Yes	76.3	71.6	
Intention to use among non-past 30–day tobacco users	(n=166)	(n=149)	
Cigarettes <sup>b</sup>			0.378
No	85.5	81.9	
Yes	14.5	18.1	
E-cigarettes <sup>b</sup>			0.143
No	80.1	73.2	
Yes	19.9	26.8	
Low nicotine cigarettes <sup>b</sup>			0.382
No	88.6	85.2	
Yes	11.4	14.8	
NRT <sup>b</sup>			0.912
No	91.0	90.6	
Yes	9.0	9.4	

Note: If missing number not provided, there are no missing data on that item. Boldface indicates statistical significance ( $p < 0.05$ ).

<sup>a</sup>M (SD).

<sup>b</sup>Column percent.

NRT, nicotine-replacement therapy.

**Table 3.** Participant Characteristics and Baseline Smoking Beliefs, by Study Condition

Characteristics	Study condition		Total % (n=521)	p-value
	Nicotine messaging % (n=263)	Combined controls % (n=258)		
Sex				0.516
Female	46.8	49.6	48.2	
Male	53.2	50.4	51.8	
Age, years				0.605
18–24	9.5	11.6	10.6	
25–34	44.9	46.9	45.9	
35–44	27.4	20.9	24.2	
45–54	8.4	10.5	9.4	
55–64	6.8	7.4	7.1	
≥65	3.0	2.7	2.9	
Hispanic ethnicity				0.524
No	88.6	90.3	89.4	
Yes	11.4	9.7	10.6	
Race				0.342
White	82.1	79.1	80.6	
Black or African American	8.4	6.6	7.5	
American Indian or Alaska Natives	1.1	0.8	1.0	
Asian	5.3	8.5	6.9	
More than 1 race	2.7	3.1	2.9	
Other	0.4	1.9	1.2	

(continued on next page)

**Table 3.** Participant Characteristics and Baseline Smoking Beliefs, by Study Condition (*continued*)

Characteristics	Study condition			p-value
	Nicotine messaging % (n=263)	Combined controls % (n=258)	Total % (n=521)	
Highest level of education completed				0.282
Less than high school	0.4	1.6	1.0	
High school/GED	14.1	10.9	12.5	
Some college/Associate's degree	35.7	32.9	34.4	
Bachelor's/Advanced degree	49.8	54.7	52.2	
Subjective financial status				0.949
Live comfortably	22.1	23.3	22.6	
Meet needs with a little left	43.7	44.2	44.0	
Just meet basic expenses	30.4	28.3	29.4	
Don't meet basic expenses	3.8	4.3	4.0	
Single Item Literacy Screener				0.468
Adequate reading ability	87.5	85.3	86.4	
Limited reading ability	12.5	14.7	13.6	
Use of tobacco products, past 30 days				0.434
None	63.1	57.8	60.5	
Other tobacco products only	5.7	6.2	6.0	
E-cigarettes	6.1	10.1	8.1	
Cigarettes	18.6	17.8	18.2	
Cigarettes and E-cigarettes	6.5	8.1	7.3	
Baseline smoking beliefs <sup>a</sup>				
Nicotine is the main substance in tobacco that makes people want to smoke	1.95 (1.03)	2.08 (1.13)	2.01 (1.08)	0.179
Smoking behavior is something basic about a person that they can't change very much	4.89 (1.26)	4.91 (1.32)	4.90 (1.29)	0.851

Note: Missing data: none. Column percentages unless otherwise noted. Boldface indicates statistical significance ( $p < 0.05$ ).

<sup>a</sup>M (SD).

There were no differences in nicotine-related norms, behavioral control, or intention to use tobacco or nicotine products by study condition (Table 2). The only marginally significant difference between groups was for social acceptability of marijuana ( $p=0.049$ ).

## DISCUSSION

Findings from the current study support that a brief nicotine messaging intervention—similar to the messages likely to be seen on warning labels or in media campaigns—can correct misperceptions of nicotine, NRT, E-cigarette, and RNC cigarette use in a general population sample of adults. Brief exposure to nicotine messages in this pilot study, however, did not impact norms about nicotine, behavioral control, or intention to use tobacco or nicotine products.

### Limitations

This study used an online convenience sample and a single, brief exposure to sample nicotine education

messages. Although it provides encouraging preliminary evidence of the potential for messaging to correct misperceptions of nicotine, studies with repeated exposures in a population sample are needed to determine whether public education on nicotine would produce similar results in U.S. adults.

## CONCLUSIONS

Public education is an essential complement to the Food and Drug Administration's efforts to move smokers away from combusted tobacco products and prevent nonusers from trying nicotine and tobacco products. Communication via mass media, warnings, and effective labeling are central components of such educational efforts and must convey correct information in a way that the public understands. Studies with more intensive exposure to such messages are needed to determine the durability of these effects and extension to behavioral outcomes, as are studies to examine their effects in subgroups of interest (e.g., tobacco users).

## ACKNOWLEDGMENTS

The authors wish to thank Richard O'Connor for his contributions to study measures. The authors were supported by NIH under awards R03CA212694 and P20GM103644 (ACV and JCW), U54DA036114 (ACV), U54DA031659 (ECD and AAS), and U54CA229973 (DM, ECD, JNC, and AAS). The content is solely the responsibility of the authors and does not necessarily represent the official views of NIH.

No financial disclosures were reported by the authors of this paper.

## REFERENCES

- IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. *Smokeless Tobacco and Some Tobacco-Specific N-Nitrosamines*. WHO, International Agency for Research on Cancer, 2007.
- IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. *Tobacco Smoking*. WHO, International Agency for Research on Cancer, 2012.
- IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. *Smokeless Tobacco*. WHO, International Agency for Research on Cancer, 2012.
- HHS. *How Tobacco Smoke Causes Disease: the Biology and Behavioral Basis for Smoking-Attributable Disease: A Report of the Surgeon General*. Atlanta, GA: U.S.: HHS, CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2010.
- HHS. *The Health Consequences of Smoking—50 Years of Progress. A Report of the Surgeon General*. Atlanta, GA: HHS, CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2014.
- Benowitz NL, Burbank AD. Cardiovascular toxicity of nicotine: implications for electronic cigarette use. *Trends Cardiovasc Med*. 2016;26(6):515–523. <https://doi.org/10.1016/j.tcm.2016.03.001>.
- O'Brien EK, Nguyen AB, Persoskie A, Hoffman AC. U.S. adults' addiction and harm beliefs about nicotine and low nicotine cigarettes. *Prev Med*. 2017;96:94–100. <https://doi.org/10.1016/j.ypmed.2016.12.048>.
- Byron MJ, Jeong M, Abrams DB, Brewer NT. Public misperception that very low nicotine cigarettes are less carcinogenic. *Tob Control*. 2018;27(6):712–714. <https://doi.org/10.1136/tobaccocontrol-2017-054124>.
- Villanti AC, Naud S, West JC, et al. Prevalence and correlates of nicotine and nicotine product perceptions in U.S. young adults, 2016. *Addict Behav*. 2019;98:106020. <https://doi.org/10.1016/j.addbeh.2019.06.009>.
- Ferguson SG, Gitchell JG, Shiffman S, et al. Providing accurate safety information may increase a smoker's willingness to use nicotine replacement therapy as part of a quit attempt. *Addict Behav*. 2011;36(7):713–716. <https://doi.org/10.1016/j.addbeh.2011.02.002>.
- Etter JF, Perneger TV. Attitudes toward nicotine replacement therapy in smokers and ex-smokers in the general public. *Clin Pharmacol Ther*. 2001;69(3):175–183. <https://doi.org/10.1067/mcp.2001.113722>.
- Borrelli B, Novak SP. Nurses' knowledge about the risk of light cigarettes and other tobacco "harm reduction" strategies. *Nicotine Tob Res*. 2007;9(6):653–661. <https://doi.org/10.1080/14622200701365202>.
- Bansal MA, Cummings KM, Hyland A, Giovino GA. Stop-smoking medications: who uses them, who misuses them, and who is misinformed about them? *Nicotine Tob Res*. 2004;6(suppl 3):S303–S310. <https://doi.org/10.1080/14622200412331320707>.
- Cummings KM, Hyland A, Giovino GA, et al. Are smokers adequately informed about the health risks of smoking and medicinal nicotine? *Nicotine Tob Res*. 2004;6(suppl 3):S333–S340. <https://doi.org/10.1080/14622200412331320734>.
- Wikmans T, Ramstrom L. Harm perception among Swedish daily smokers regarding nicotine, NRT-products and Swedish Snus. *Tob Induc Dis*. 2010;8:9. <https://doi.org/10.1186/1617-9625-8-9>.
- Shiffman S, Ferguson SG, Rohay J, Gitchell JG. Perceived safety and efficacy of nicotine replacement therapies among U.S. smokers and ex-smokers: relationship with use and compliance. *Addiction*. 2008;103(8):1371–1378. <https://doi.org/10.1111/j.1360-0443.2008.02268.x>.
- Donny EC, Denlinger RL, Tidey JW, et al. Randomized trial of reduced-nicotine standards for cigarettes. *N Engl J Med*. 2015;373(14):1340–1349. <https://doi.org/10.1056/NEJMsa1502403>.
- Higgins ST, Heil SH, Sigmon SC, et al. Addiction potential of cigarettes with reduced nicotine content in populations with psychiatric disorders and other vulnerabilities to tobacco addiction. *JAMA Psychiatry*. 2017;74(10):1056–1064. <https://doi.org/10.1001/jamapsychiatry.2017.2355>.
- H.R. 1256—111th Congress: Family Smoking Prevention and Tobacco Control Act. GovTrack.us (database of federal legislation); 2009.
- Morris NS, MacLean CD, Chew LD, Littenberg B. The Single Item Literacy Screener: evaluation of a brief instrument to identify limited reading ability. *BMC Fam Pract*. 2006;7:21. <https://doi.org/10.1186/1471-2296-7-21>.
- U.S. Food and Drug Administration. FDA's plan for tobacco and nicotine regulation. [www.fda.gov/TobaccoProducts/NewsEvents/ucm568425.htm](http://www.fda.gov/TobaccoProducts/NewsEvents/ucm568425.htm). Published 2017. Accessed February 7, 2018.
- U.S. Food and Drug Administration. Modifications to labeling of nicotine replacement therapy products for over-the-counter human use. [www.federalregister.gov/documents/2013/04/02/2013-07528/modifications-to-labeling-of-nicotine-replacement-therapy-products-for-over-the-counter-human-use](http://www.federalregister.gov/documents/2013/04/02/2013-07528/modifications-to-labeling-of-nicotine-replacement-therapy-products-for-over-the-counter-human-use). Published 2013. Accessed July 3, 2018.
- Mercincavage M, Saddleson ML, Gup E, et al. Reduced nicotine content cigarette advertising: how false beliefs and subjective ratings affect smoking behavior. *Drug Alcohol Depend*. 2017;173:99–106. <https://doi.org/10.1016/j.drugalcdep.2016.12.022>.
- Strasser AA, Tang KZ, Tuller MD, Cappella JN. PREP advertisement features affect smokers' beliefs regarding potential harm. *Tob Control*. 2008;17(suppl 1):i32–i38. <https://doi.org/10.1136/tc.2007.022426>.
- Pierce JP, Choi WS, Gilpin EA, Farkas AJ, Merritt RK. Validation of susceptibility as a predictor of which adolescents take up smoking in the United States. *Health Psychol*. 1996;15(5):355–361. <https://doi.org/10.1037//0278-6133.15.5.355>.
- Pierce JP, Farkas AJ, Evans N, Gilpin E. An improved surveillance measure for adolescent smoking? *Tob Control*. 1995;4(suppl 1):S47–S56. <https://doi.org/10.1136/tc.4.suppl1.s47>.