

Immunological Aspects

Impact of individual-level factors on *Ex vivo* mycobacterial growth inhibition: Associations of immune cell phenotype, cytomegalovirus-specific response and sex with immunity following BCG vaccination in humans

Satria A. Prabowo^{a,b,*}, Steven G. Smith^{a,b}, Karin Seifert^{a,1}, Helen A. Fletcher^{a,b}

^a Department of Immunology and Infection, Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, UK

^b Tuberculosis Centre, London School of Hygiene and Tropical Medicine, London, UK



ARTICLE INFO

Keywords:

Tuberculosis vaccine
BCG
Growth inhibition assay
Cytomegalovirus
Sex
NK cell

ABSTRACT

Understanding factors associated with varying efficacy of Bacillus Calmette-Guérin (BCG) would aid the development of improved vaccines against tuberculosis (TB). In addition, investigation of individual-level factors affecting mycobacterial-specific immune responses could provide insight into confounders of vaccine efficacy in clinical trials. Mycobacterial growth inhibition assays (MGIA) have been developed to assess vaccine immunogenicity *ex vivo* and provide a measure of immune function against live mycobacteria. In this study, we assessed the impact of immune cell phenotype, cytomegalovirus (CMV)-specific response and sex on *ex vivo* growth inhibition following historical BCG vaccination in a cohort of healthy individuals (n = 100). A higher frequency of cytokine-producing NK cells in peripheral blood was associated with enhanced *ex vivo* mycobacterial growth inhibition following historical BCG vaccination. A CMV-specific response was associated with T-cell activation, a risk factor for TB disease and we also observed an association between T-cell activation and *ex vivo* mycobacterial growth. Interestingly, BCG-vaccinated females in our cohort controlled mycobacterial growth better than males. In summary, our present study has shown that individual-level factors influence capacity to control mycobacterial growth following BCG vaccination and the MGIA could be used as a tool to assess how vaccine candidates may perform in different populations.

1. Introduction

Tuberculosis (TB) is the number one cause of death from an infectious disease worldwide and it is currently estimated that a quarter of the world population is infected with *Mycobacterium tuberculosis* (*Mtb*) [1,2]. The introduction of Bacillus Calmette-Guérin (BCG) vaccination and chemotherapy in the past century provided optimism to fight the disease. Despite this, drug-resistant TB is now a major risk to global health security, and BCG as the only licensed vaccine for TB is known to have a variable efficacy against contagious adult pulmonary TB [3,4]. BCG remains as the most widely used vaccine worldwide, primarily because it provides good protection against TB in children [3]. Understanding factors associated with varying BCG protection could aid the development of improved vaccination practice as well as novel vaccines

against TB.

It has been proposed that the observed variation in BCG efficacy is attributed to individual-level factors which influence host mycobacteria-specific immune responses [5–7]. In a recent systematic review, protection following BCG vaccination was shown to vary according to the geographical latitudes in which the vaccine was given. In the UK, a country where exposure to environmental mycobacteria and/or *Mtb* is regarded to be lower (latitude > 40°), BCG is known to provide efficacy of up to 80% against pulmonary TB [8] and vaccination of school-aged children could provide protection for more than 20 years [9]. Another factor that may influence the mycobacteria-specific immune response is sex. Globally, TB case rates are much higher in men than in women, as reflected by a global male to female ratio (M:F) of 1.7 for case notifications in 2016 [10]. Males contribute to 65% of TB cases worldwide

* Corresponding author. Department of Immunology and Infection, Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, UK.

E-mail address: satria.prabowo@lshtm.ac.uk (S.A. Prabowo).

¹ Current address: Federal Institute for Drugs and Medical Devices, Bonn, Germany.

<https://doi.org/10.1016/j.tube.2019.101876>

Received 30 December 2018; Received in revised form 2 August 2019; Accepted 29 September 2019

Available online 30 September 2019

1472-9792/Crown Copyright © 2019 Published by Elsevier Ltd. All rights reserved.

and although it is thought that socioeconomic and cultural factors are contributing to the observed sex bias, differences in the immune responses between the sexes also play a role [11,12]. It is generally acknowledged that females exhibit more robust immune responses towards infection and vaccination compared to males [13]. In the context of susceptibility to TB, differences in immune cells frequencies and functions have been thought to contribute to higher TB rates in males [14]. With regard to BCG vaccination, there is currently limited evidence concerning the impact of sex on its protective effect against pulmonary TB in adults. Interestingly, BCG is thought to provide a non-specific protective effect against unrelated pathogens, thus contributing in reduction of overall cause of mortality, and this effect is more pronounced in females rather than males [15–17].

Recently, Fletcher et al. found that T-cell activation is an immune correlate of risk of TB disease in BCG-vaccinated infants in a study enrolling a large cohort of infants [18]. Chronic exposure to antigen from persistent viral or bacterial infection is known to drive continuous T-cell activation which could lead to dysfunction of antigen specific T-cells [19]. Further to the findings of the infant study, it was identified that cytomegalovirus (CMV)-specific IFN- γ responses were associated with T-cell activation and could have contributed to increased risk of developing TB disease [20].

The mycobacterial growth inhibition assay (MGIA) has been developed as a measure of vaccine immunogenicity *ex vivo*. Following optimisation works in the past few years [21–23], the assay has gained attention for its potential ability to detect vaccine-mediated inhibition of growth following BCG vaccination in adults and infants [24–26]. The assay described in the present study involves direct co-culture of peripheral blood mononuclear cells (PBMCs) with mycobacteria, and subsequent measurement of mycobacterial growth inhibition as a functional assessment of vaccine response. Several studies have demonstrated the ability of the MGIA to detect changes in the innate and adaptive compartment following vaccination [25,27–30]. Recently, Joosten and colleagues (2018) found that the capacity to control mycobacterial growth following recent *Mtb* exposure or BCG vaccination is associated with nonclassical monocytes, and this observation is reflective of the trained innate immune mechanism [26]. In a study by Jensen et al., IFN- γ was associated with reduction of mycobacterial growth *ex vivo* following immunisation with a TB vaccine candidate in mice [31]. However, in that study the source IFN- γ was not found among the investigated vaccine-specific T-cells, suggesting potential contribution from other cell types, such as NK cells.

In this study, we demonstrated the impact of immune cell phenotype, CMV-specific response and sex on vaccine-specific mycobacterial growth inhibition following historical BCG vaccination in adult healthy volunteers.

2. Materials and methods

2.1. Study participants and ethics statement

We recruited 100 healthy adult participants with (i) no history of BCG vaccination or (ii) a history of BCG vaccination more than 6 months before study enrolment. Verbal interviews were conducted to determine eligibility based on the absence of any major chronic illness, current medication administration or symptoms of infection. Participants were aged 18–80 years with no evidence of exposure or infection with TB. Participants were excluded if they were suffering from any persistent medical condition or infection. Sample size was calculated based on the assumption of effect size 0.70, with power 0.8 and significance level 0.05. Written informed consent was obtained from all participants prior to enrolment in the study. Individuals were recruited under protocols approved by the LSHTM Observational Research Ethics Committee (ref 8762 and 10485). All procedures were conducted in accordance with the Declaration of Helsinki, as agreed by the World Medical Association General Assembly (Washington, 2002) and ICH Good Clinical Practice

(GCP).

2.2. PBMCs isolation and IFN- γ enzyme-linked immunospot (ELISpot) assay

Peripheral blood (50 ml) was collected and processed within 6 h. PBMCs isolation and IFN- γ ELISpot assay were performed as previously described [32]. PBMCs were cryopreserved in FBS (Labtech International Ltd, Uckfield, UK) containing 10% DMSO (Sigma-Aldrich) and stored in -80°C freezer using CoolCell containers (VWR International, Lutterworth, UK). PBMCs were thawed and an *ex vivo* IFN- γ ELISpot assay was performed to assess antigen-specific response. PBMCs were incubated overnight for 18 h with 20 $\mu\text{g}/\text{ml}$ purified protein derivative (PPD) (Oxford Biosystem, Oxfordshire, UK). Positive control Phytohemagglutinin (PHA) (10 $\mu\text{g}/\text{ml}$, Sigma-Aldrich) and negative control (medium-only) wells were included for each participant samples. Results are reported as spot forming cells (SFC) per million PBMCs, calculated by subtracting the mean of the unstimulated wells from the mean of antigen wells and correcting for the numbers of PBMC in the wells. Spots were quantified using an automated plate reader with ELISpot 5.0 software as well as checked visually.

2.3. *Ex vivo* mycobacterial growth inhibition assay

The growth inhibition assay was performed using cryopreserved PBMCs of the study participants, as previously described [32]. In brief, a 2-ml screw-cap tubes containing 3×10^6 PBMCs in 600 ml of medium were rotated at 37°C with ~ 100 Colony Forming Units (CFU) of BCG Pasteur Aeras strain (Rockville, MD, USA) for 4 days. The PBMCs were then lysed with sterile water, and the lysate transferred to a Bactec MGIT supplemented with PANTA antibiotics and OADC enrichment broth (all from Becton Dickinson, Oxford, UK). The tube was placed in a Bactec MGIT 960 and incubated until growth was detected (measured as time to positivity [TTP]). Use of a standard curve enables conversion of the TTP of a sample tube into bacterial numbers (log CFU) (Supplementary Fig. S1). All work with cells pre-BCG infection and involving BCG infected samples was done in Biosafety Level (BSL) 2 laboratory.

2.4. Enzyme-linked immunosorbent assay (ELISA)

MGIA supernatants were analysed to assess cytokine concentrations by ELISA. The levels of following cytokines were measured: IFN- γ , interleukin (IL)-12p40, IL-6 [BD OptiEIA kits, Becton Dickinson, UK], tumor necrosis factor alpha (TNF- α), granulocyte-macrophage colony-stimulating factor (GM-CSF), interferon-gamma-induced protein 10 (IP-10), granzyme B, IL-32, IL-22 [R&D Systems, Abingdon, UK], IL-10, IL-17 [BioLegend, London, UK] and perforin [Abcam, Cambridge, UK]. Assays were performed according to the manufacturers' instruction.

2.5. Flow cytometric immune phenotyping

PBMC were washed and stained with 1 $\mu\text{l}/\text{ml}$ Live Dead Blue Stain (Invitrogen), followed by staining with the following titrated antibody for the Lymphocyte panel: 2.5 μl CD3-AF700 (clone UCHT1, Ebioscience, Loughborough, UK), 1.25 μl CD4-APC/Cy7 (clone RPA-T4, BioLegend), 1.25 μl CD8-Superbright645 (clone RPA-T8, Ebioscience), 2.5 μl CD19-FITC (clone HIB19, BioLegend), 2.5 μl CD56-APC (clone HCD56, BioLegend), 2.5 μl CD16-BV510 (clone 3G8, BioLegend), 5 μl HLA-DR-PE (clone L243, BioLegend), 5 μl LAG3-PE/Cy7 (clone 11C3C65, BioLegend) and 1.25 μl PD1-BV421 (clone EH12.2H7, BioLegend). For the Monocyte panel, the cells were stained with the following titrated antibodies: 2.5 μl CD3-AF700 (clone UCHT1, Ebioscience), 2.5 μl CD19-FITC (clone HIB19, BioLegend), 2.5 μl CD14-BV421 (clone HCD14, BioLegend), 2.5 μl CD16-BV510 (clone 3G8, BioLegend), 1.25 μl CD86-APC/Cy7 (clone IT2.2, BioLegend), 5 μl HLA-DR-PE (clone L243, BioLegend), 5 μl CD206-APC (clone 15-2,

Table 1
Characteristics of study participants.

Characteristic	Total Participants: 100	
	Naïve (n = 37)	BCG Vaccinated (n = 63)
Female [no. (%)]	28 (75.7%)	42 (66.7%)
Median age [yr (range)]	31 (23–70)	39 (24–80)
Average time since BCG vaccination [yr (range)]	–	29.4 (10–58)
Country of Origin	8 (21.6%)	44 (69.8%)
UK [no. (%)]		

BioLegend), 5 µl CD163-BV605 (clone GHI/61, BioLegend), 2.5 µl CD64-APC/Cy7 (clone 10.1, BioLegend) and 5 µl CD123-BV650 (clone 6H6, BioLegend). Fluorescence minus one (FMO) controls were set using cells for each antibody and used to guide gating. Cells were acquired on a BD LSR II flow cytometer. Data was analysed with FlowJo software version 10.4 (Treestar Inc., USA). Results are presented as percentages of cells after gating out of dead cells and doublets. Gating strategies for the lymphocyte and monocyte panels are described in [Supplementary Figs. S2A and S2B](#).

2.6. Intracellular cytokine staining (ICS) flow cytometry

The ICS flow cytometry was performed as previously described [32]. In brief, PBMCs were then incubated alone (medium only) as a negative control, with 5 µg/ml *Staphylococcus enterotoxin B* (SEB; Sigma, UK) as a positive control, with ~100 CFU BCG (as per the MGIA protocol) and with 10 µg/ml CMV peptide pool (5 peptides, 2 µg/ml/peptides, ANASPEC, Fremont, CA, USA). The CMV peptide pool used is the same as the Fletcher et al. study [20]. The incubation with BCG was performed for 4 days and the addition of SEB and CMV was performed on Day 3. Two hours after the addition of SEB and CMV to the respective tubes, brefeldin A (Sigma, UK) was added to all tubes which were then incubated for 18 h at 37 °C until Day 4. Data was acquired using an LSRII flow cytometer (BD Biosciences) and FACSDiva acquisition software (BD Biosciences). ICS flow cytometry data was analysed using FlowJo software version 10.4 (TreeStar Inc., Ashland, OR, USA). Samples were gated sequentially on singlet, live, CD14⁻CD19⁻, CD3⁺ (lymphoid), CD4⁺, CD8⁺ cells and negative control stimulation tubes were used to set cytokine gates (see [Supplementary Fig. S3](#), ICS gating).

Table 2
Summary of mean cytokine responses measured with ELISA assays, assessed from MGIA supernatant samples after 4 days of co-culture. Comparisons were made between naïve and BCG-vaccinated groups (unpaired *t*-test), the values indicate mean of concentration in pg/ml [95% CI]. Correlations were assessed with *ex vivo* mycobacterial growth among responders (Spearman's correlation). A *p* value < 0.05 was considered statistically significant (in bold), and after a multiple testing correction only values with *p* < 0.0063 were considered significant (underlined). n = 37 BCG-naïve and n = 63 BCG-vaccinated participants. ****p* < 0.001.

Cytokine (pg/ml)	Comparison			Correlation with <i>ex vivo</i> mycobacterial growth	
	Naïve	BCG-vaccinated	<i>p</i> -value	<i>r</i>	<i>p</i> -value
IFN-γ	12.47 [8.245–16.69]	23.37 [10.8–35.94]	0.1962	–0.027	0.8432
IP-10	111.7 [55.42–168]	204.5 [112.1–297]	0.1505	0.19	0.1158
TNF-α	37.97 [2.547–73.4]	97.98 [38.61–157.4]	0.1471	–0.35	0.0558
IL-12	27.6 [3.033–52.17]	63.61 [20.83–106.4]	0.2299	–0.23	0.3158
IL-10	52.55 [31.07–74.03]	59.99 [36.56–83.41]	0.6688	<u>0.37***</u>	0.0003
GM-CSF	7.729 [–1.688–17.15]	88.54 [26.19–150.9]	0.0512	–0.37	0.1552
IL-6	356.7 [246.1–467.3]	315 [236.5–393.4]	0.5293	0.071	0.5449
IL-17	0.00 [0.00–0.00]	0.1596 [0.00–0.4083]	0.3291	–0.13	0.2141

2.7. Statistical analysis

To identify statistical significance of *ex vivo* growth inhibition (log CFU values) and ELISA responses, students *t*-test were used. Mann-Whitney *U* Test was performed to identify significant differences of the ELISpot, cell surface flow cytometry and ICS responses between groups. Spearman's correlation coefficient was used to test for correlations between growth inhibition and immune responses. A multiple comparison correction was included (Bonferroni), as indicated in each figure legend. Statistical analyses were performed in GraphPad Prism 7 (GraphPad, La Jolla, CA, USA).




Fig. 1. Growth inhibition and immune responses following historical BCG vaccination. Assessment was performed from 37 BCG-naïve and 63 BCG-vaccinated participants. (A) Growth inhibition was compared using BCG input ~ 100 Colony Forming Unit (CFU) as immune target (unpaired *t*-test). Data is presented as total number of log CFUs per sample, which was determined by use of a standard curve. (B) IFN-γ production from PBMC following stimulation with PPD was compared (Mann-Whitney test). Numbers above each group represent median (range). SFC, spot forming cells. (C) The correlation between *ex vivo* growth inhibition and PPD-specific IFN-γ response was assessed (Spearman's correlation). A *p* value < 0.05 was considered statistically significant. Dots and squares represent individual data points, and the central lines indicate the median response with inter-quartile range (IQR). *****p* < 0.0001.

Table 3

Correlation of immune cell frequencies in peripheral blood and *ex vivo* mycobacterial growth inhibition. Assessment was performed from 16 BCG-naïve and 34 BCG-vaccinated participants. Correlations were performed from a total of 50 participants, as well as from each naïve and BCG-vaccinated groups respectively (Spearman's correlation). A p value < 0.05 was considered statistically significant (in bold), and after a multiple testing correction only values with p < 0.0031 were considered significant (underlined). Note: The ML ratio was obtained by dividing the percentage of monocytes by the sum of the percentages of T- and B-cells. The NK cell ratio was obtained by dividing the percentage of cytokine-producing by cytotoxic NK cells. *p < 0.05, **p < 0.01.

Leukocyte subsets	Correlation with <i>ex vivo</i> mycobacterial growth					
	All participants		Naïve		BCG-vaccinated	
	r	p-value	r	p-value	r	p-value
T-cells	-0.068	0.6367	0.29	0.2708	-0.30	0.0866
CD4 T-cells	-0.041	0.7764	0.17	0.5172	-0.091	0.6080
CD8 T-cells	0.24	0.0938	0.36	0.1714	0.093	0.6011
CD4/CD8 ratio	-0.16	0.2718	-0.20	0.4579	-0.058	0.7448
NK cells	-0.27	0.0593	-0.71**	0.0028	-0.19	0.2833
Cytokine NK cell	-0.26	0.0702	-0.47	0.0679	-0.41*	0.0147
Cytotoxic NK cell	-0.25	0.0814	-0.64**	0.0093	-0.19	0.2699
NK cell ratio	-0.2	0.1602	-0.35	0.1866	-0.087	0.6241
Monocytes	0.12	0.4244	-0.0088	0.9758	0.13	0.4638
ML ratio	0.064	0.6609	-0.044	0.8714	0.083	0.6390
M1 monocytes	-0.076	0.5993	-0.28	0.2867	-0.031	0.8610
M2 monocytes	-0.16	0.2784	-0.16	0.5458	-0.12	0.4978
M1/M2 ratio	0.059	0.6831	-0.17	0.5283	0.15	0.3939
CD64 ⁺ monocytes	-0.063	0.6659	-0.29	0.2664	0.028	0.8759
CD123 ⁺ monocytes	-0.072	0.6169	-0.27	0.3025	0.015	0.9313
Suppressor monocytes	0.21	0.1414	0.31	0.2381	0.089	0.6149

3. Results

3.1. Demographics of enrolled participants

One hundred participants were enrolled in the study; 37 vaccine-naïve volunteers with no history of BCG vaccination and 63 volunteers previously-vaccinated with BCG (average time since vaccination 29.4 years prior to enrolment). Table 1 summarises the characteristics of the study participants. Almost 70% of the BCG-vaccinated participants were from the UK.

3.2. Assessment of *ex vivo* growth inhibition and mycobacterial antigen-specific cytokine responses

The growth inhibition assay was performed to assess impact of historical BCG vaccination on *ex vivo* mycobacterial growth control. Using cryopreserved PBMCs, enhanced growth inhibition in PBMCs from BCG-vaccinated individuals was observed compared to vaccine-naïve individuals (median log CFU 1.680 and 2.027, p < 0.0001, Fig. 1A). The IFN- γ ELISpot assay was performed to measure the magnitude of the mycobacteria-specific response. The secretion of IFN- γ in response to PPD was elevated in samples from vaccinated individuals in comparison to unvaccinated individuals (median SFC 109.5 and 48, p < 0.0001, Fig. 1B). There was a significant inverse correlation between higher IFN- γ ELISpot response and lower mycobacterial growth (p = 0.022, Spearman r = -0.23, Fig. 1C).

Trends for higher production of Th1-type cytokines (IFN- γ , IP-10, TNF- α , IL-12) as well as GM-CSF were observed in the BCG-vaccinated group compared to the vaccine-naïve group (Table 2). There was a statistically significant correlation between higher IL-10 production and higher mycobacterial growth (Spearman r = 0.37, p = 0.0003, Table 2). Meanwhile, historical BCG-vaccination was associated with significantly increased frequency of mycobacterial antigen specific IL-2⁺ CD4 T-cells in the BCG-vaccinated group upon 4 days of stimulation with BCG (p = 0.008, Supplementary Fig. S4). Similar trends were observed with the frequencies of IFN- γ ⁺ as well as TNF- α ⁺ CD4 T-cells (Supplementary Fig. S4). There were no significant correlations between the frequencies of BCG-specific CD4 and CD8 T-cells and mycobacterial growth, although the observed trends suggest that these cells may contribute to control of growth (Supplementary Table S1).

3.3. Associations between historical BCG vaccination and the frequency of circulating leukocyte subsets

Historical BCG vaccination was not associated with frequencies of circulating leukocytes in T-cell, NK cell and monocyte compartments (Supplementary Table S2). However, significant correlations were observed between the frequencies of NK cells and enhanced control of mycobacterial growth *ex vivo* in the naïve and BCG-vaccinated groups (p < 0.05, Spearman's correlations, Table 3). In the BCG-vaccinated group, higher frequency of cytokine-producing NK cells was associated with reduced mycobacterial growth (Spearman r = -0.41, p = 0.015, Fig. 2A). A higher production of perforin was observed from the cells of BCG-vaccinated participants compared to naïve (p = 0.018, Fig. 2B). The production of perforin significantly correlated with enhanced growth inhibition (Spearman r = -0.44, p = 0.013, Fig. 2C and Supplementary Table S3), and the association was still significant when the correlation was performed in the BCG-vaccinated group only (Spearman r = -0.36, p = 0.037, data not shown). Correlations with other




Fig. 2. NK cells correlations. A higher frequency of cytokine-producing NK cells (CD56^{bright} CD16^{+/−}) correlated with enhanced *ex vivo* mycobacterial growth inhibition (Spearman's correlation) (A). A perforin ELISA was performed from MGIA supernatants and the response was compared between vaccination groups (unpaired *t*-test) (B). The production of perforin was associated with enhanced *ex vivo* growth inhibition (Spearman's) (C). A p value < 0.05 was considered statistically significant. *p < 0.05, **p < 0.01.

Table 4

Correlation of CMV-specific T-cell responses and T-cell activation. Associations were investigated from 3 different subsets of CMV-specific cytokine⁺ T-cells producing IFN- γ ⁺, IL-2⁺ or TNF- α ⁺, respectively. Three markers were used for T-cell activation: HLA-DR, LAG3 and PD1. A p value < 0.05 was considered statistically significant (in bold), and after a multiple testing correction only values with p < 0.0083 were considered significant (underlined) (Spearman's correlation). n = 50 participants, consisted of 16 BCG-naïve and n = 34 BCG-vaccinated participants. *p < 0.05, **p < 0.01.

CMV-specific cytokine ⁺ T-cells	Correlation with activated T-cells					
	HLA-DR ⁺ CD4 T-cells		LAG3 ⁺ CD4 T-cells		PD1 ⁺ CD4 T-cells	
	r	p-value	r	p-value	r	p-value
IFN- γ ⁺ CD4 T-cells	0.026	0.8748	-0.004	0.9805	0.20	0.2112
IL-2 ⁺ CD4 T-cells	-0.045	0.7823	-0.056	0.7310	-0.0082	0.9601
TNF- α ⁺ CD4 T-cells	0.054	0.7401	0.058	0.7239	0.091	0.5757
	HLA-DR ⁺ CD8 T-cells		LAG3 ⁺ CD8 T-cells		PD1 ⁺ CD8 T-cells	
	r	p-value	r	p-value	r	p-value
	IFN- γ ⁺ CD8 T-cells	0.31	0.0552	0.39*	0.0140	<u>0.44**</u>
IL-2 ⁺ CD8 T-cells	-0.087	0.5917	0.0024	0.9885	-0.15	0.3609
TNF- α ⁺ CD8 T-cells	0.28	0.0799	0.35*	0.0281	0.33*	0.0375

measured NK cell associated markers (granzyme, IL-32, IL-22) did not reach significance (Supplementary Table S3).

3.4. Impacts of CMV-specific T-cell response and T-cell activation on ex vivo mycobacterial growth inhibition

CMV-specific T-cells producing IFN- γ ⁺ and TNF- α ⁺, notably in the CD8 compartment, were significantly associated with the frequency of T-cells expressing LAG3 and PD1 markers (p < 0.05, Spearman's correlations, Table 4 and Fig. 3A–D). Historical BCG-vaccination was not associated with differences in CMV-specific response nor T-cell activation (Supplementary Tables S2 and S4). However, T-cell activation was shown to correlate with higher growth of mycobacteria ex vivo, particularly in the naïve group (Fig. 3 and Supplementary Table S5). LAG3⁺ CD4 T-cells were significantly associated with growth of mycobacteria (p = 0.047), with a similar trend for LAG3⁺ CD8 T-cells (p = 0.072) (Fig. 3 F and I).

3.5. Impact of sex on ex vivo mycobacterial growth inhibition and cytokine responses, and its association with immune cell phenotype

In this study, we demonstrated that sex was associated with differences in immune response following historical BCG vaccination. First, BCG-vaccinated females exhibited a superior capacity to control mycobacterial growth when compared to males (p = 0.029, Fig. 4B). In contrast, males showed a trend towards higher IFN- γ response from PPD-stimulated PBMCs as well as higher IP-10 production in the MGIA supernatant, both in naïve and BCG-vaccinated groups (Fig. 4C–F).




Fig. 3. CMV-specific responses were associated with higher CD8 T-cell activation, expressing markers LAG3 (A–B) and PD1 (C–D) respectively. Activated CD4 and CD8 T-cells (E–J) were correlated with higher growth of mycobacteria, notably in the naïve groups (F, I). A p value < 0.05 was considered statistically significant (Spearman's correlation). n = 50 participants, consisted of 16 BCG-naïve and 34 BCG-vaccinated participants. *p < 0.05, **p < 0.01.




Fig. 4. Sex impact on growth inhibition and immune responses following historical BCG vaccination. Assessment was performed from 37 BCG-naïve (A,C,E) and 63 BCG-vaccinated participants (B,D,F). (A–B) Growth inhibition was compared between sex and data was presented as total number of log CFUs per sample (unpaired *t*-test). (C–D) IFN- γ production from PBMC following stimulation with PPD was compared (Mann-Whitney test). Numbers above each group represent median (range). SFC, spot forming cells. (E–F) IP-10 was measured from MGIA supernatants using ELISA assay (mean, unpaired *t*-test). Dots and squares represent individual data points, and the central lines indicate the median response with IQR. * $p < 0.05$.

Supplementary Table S6 summarises the sex comparisons of all measured cytokines from the MGIA supernatants.

In the BCG-vaccinated group, females had a higher frequency of cytokine-producing NK cells ($p = 0.018$, Fig. 5A). There was also a higher CD4/CD8 ratio in females compared to males in the naïve group ($p = 0.028$, Fig. 5B). Interestingly, there was a higher frequency of monocytes in males in the BCG-vaccinated group ($p = 0.049$, Fig. 5C), with a trend of higher monocyte-to-lymphocyte (ML) ratio in BCG-vaccinated males compared to females ($p = 0.08$, Supplementary Table S7). In terms of T-cell activation, BCG-vaccinated females exhibited a lower frequency of LAG3⁺ CD8 T-cells ($p = 0.0297$, Fig. 5D). While in the naïve group, females also had lower frequencies of activated CD8 T-cells expressing HLA-DR, LAG3 and PD1 ($p < 0.05$, Supplementary Table S7). The lower frequencies of activated T-cells in females may be a consequence of lower CMV-specific CD8 T-cells response (Fig. 5E and Supplementary Table S7).

4. Discussion

The present study reports that mycobacterial growth inhibition *ex vivo* is enhanced following historical BCG vaccination in adult healthy volunteers. In this study, the average time since BCG vaccination was 29.4 years prior to enrolment. Our results are in line with previous studies such as Fletcher et al. [24] and Prabowo et al. [32] which detected the impact of historical BCG vaccination after more than 20 years using the same PBMC-based MGIA. Most vaccinated individuals enrolled in our study are UK participants in which BCG vaccination is known to be effective [9]. A higher IFN- γ response was also observed in the BCG-vaccinated group compared to the naïve group using the ELISpot assay, reflecting the presence of mycobacterial-specific memory cells. Moreover, there was a significant correlation between IFN- γ response and lower mycobacterial growth. Several published MGIA studies reported increased IFN- γ production following BCG vaccination [24,33,34], and BCG-specific IFN- γ response measured by ELISpot assay is known to be associated with reduced TB disease risk following BCG vaccination in infants [18]. The ELISpot assay measures all cells that




Fig. 5. Comparison by sex of immune cells phenotype (A–C), T-cell activation (D) and CMV-specific T-cell response (E). Assessment was performed from 16 BCG-naïve and 34 BCG-vaccinated participants. The box plots show the minimum and maximum values (ends of the whiskers), the median (band near the middle of the box) and interquartile ranges. Blue and red colour represent males and females, respectively. A p value < 0.05 was considered statistically significant (Mann-Whitney). *p < 0.05, **p < 0.01, ***p < 0.0001. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

secrete IFN- γ in response to antigen stimulation, including NK cells and $\gamma\delta$ T-cells in addition to conventional T-cells. Focusing on the conventional T-cells response, in our study, we did not observe a significant association between Th1-type cytokine-expressing T-cells and *ex vivo* mycobacterial growth inhibition. This finding was in contrast with the study of Smith et al. [25] which showed an association between MGIA control capacity and the frequency of polyfunctional CD4 T-cells using studying a small cohort of BCG-vaccinated infants. However, our results were consistent with the finding of Joosten et al. [26] using the same PBMC-based MGIA, as well as with a study by Kagina et al. [35] which showed no association between polyfunctional T-cells and the risk to develop TB disease following BCG vaccination.

We also observed trends of higher Th1-type cytokines in the MGIA supernatants from BCG-vaccinated participants compared to the naïve. Interestingly, there was a strong significant correlation between IL-10 production and reduced control of mycobacterial growth. This observation replicates earlier findings, in which IL-10 was associated with reduced *ex vivo* growth inhibition, and was significantly predictive of mycobacterial growth through inhibition of other pro-inflammatory cytokines [36,37]. IL-10 is known to have immunosuppressive activity by inhibiting T-cell proliferation and IFN- γ production, leading to reduced macrophage activation [38]. The capacity of individuals to produce IL-10 may need to be considered when assessing TB vaccine effects in clinical trials.

In this study, the frequency of NK cells – in particular cytokine-producing NK cells – is associated with enhanced *ex vivo* mycobacterial growth inhibition following historical BCG vaccination. This may account for our correlation between IFN- γ ELISpot response and control of mycobacterial growth as IFN- γ secreting NK cells will be measured in addition to CD4 and CD8 positive T-cells. Our results again support a recent finding, in which a greater frequency of putative cytokine-producing CD16⁺ NK cells was associated with reduced mycobacterial growth in the multiple regression analysis of MVA85A correlate of risk

study [36,39]. Cytokine-producing NK cells are the main source of NK-cell derived cytokines such as IFN- γ , TNF- α and GM-CSF [40], which were modestly increased in the MGIA supernatants of the BCG-vaccinated group in our study. Initially, cytotoxicity and cytokine-producing functions of NK cells were regarded as two distinct functions with little synergy between them [40,41]. However, it was recently shown that IFN- γ and TNF- α could synergistically enhance NK cell cytotoxicity [42]. In our study, cells obtained from BCG-vaccinated participants produced a higher level of perforin and the secretion of this lytic granules was associated with enhanced growth inhibition.

Although considered a component of innate immune system, an emerging body of evidence has revealed that NK cells can also behave in a memory-like manner following infection or vaccination [reviewed in Refs. [43,44]]. NK cells isolated from pleural fluid express the memory marker CD45RO and produce higher amounts of IFN- γ and IL-22 in response to stimulation with IL-12, IL-15 and BCG when compared with CD45RO⁺ cells [45,46]. Even though NK cells do not have antigen receptors generated by genetic rearrangement, they possess receptors which allow direct antigenic contact, resulting in subsequent cellular activation [44]. This process will generate antigen-specific NK cells, which lead to enhanced response following re-exposure with the same stimulus [47,48]. In addition, work by Kleinnijenhuis et al. reveals that BCG vaccination promotes augmented secondary responses towards the same and unrelated stimulus through trained innate immunity mechanism [49]. The growth inhibition assay has recently been shown to be able to detect contribution from the trained innate immune compartment, following *Mtb* exposure or BCG vaccination, by the role of nonclassical monocytes [26]. Our present study has shown the additional contribution of NK cells to *ex vivo* mycobacterial growth control, and in line with this, recent clinical trials also reported that immune cells associated with protection from TB disease and after BCG vaccination were not T-cells, but IFN- γ -producing NK cells [50,51].

Furthermore, we have demonstrated that a CMV-specific response

may be associated with T-cell activation, in particular in the CD8 compartment, and this activation is correlated with mycobacterial growth *ex vivo*. In HIV, T-cell activation has been established as a risk factor for acquisition of infection as well as progression from infection to disease [52–54]. In TB, evidence has emerged denoting the role of CMV and T-cell activation on TB disease risk [18,20], and our study is the first to show such association with *ex vivo* mycobacterial growth. In this study, we chose to measure CMV-specific T-cell cytokine response with ICS flow cytometry rather than with serology, as evidence in the literature showed that CMV-antibody levels do not correlate with the size of the T-cell response against CMV and the ICS method is more sensitive for detection of CMV-specific cytokine-producing T-cells [55,56]. Unfortunately in this study due to a technical limitation, we were unable to perform CMV serology in our cohort samples. CMV infection is recognised to drive the expansion of NKG2C⁺ NK cells [57], which do not respond well to cytokine stimulation discussed above [58,59]. Further studies are required to better understand the interplay between CMV-specific response, T-cell activation and NK cells in the context of BCG vaccination.

Differences in TB disease notification rates between the sexes are well documented and thought to be a result of biological factors, in addition to social factors [11,14,60]. Therefore, it is of interest that our study demonstrated a higher capacity of BCG-vaccinated females to control mycobacterial growth *ex vivo* compared to males. In conjunction to this data, we found that females had a higher frequency of cytokine-producing NK cells, and lower frequency of activated T-cells as well as CMV-specific response. In addition, females also had a lower monocyte frequency, with a trend of a lower ML ratio compared to males. Altogether, these individual-level factors appear to contribute to the enhanced growth inhibition in females following BCG vaccination. Such sex specific effect has also been observed with measles and smallpox vaccines, where females are more protected than males following vaccination [61,62]. The epidemiological observation that the sex bias in TB does not arise until puberty has suggested the important role of sex hormones [11]. In general, testosterone is considered to downregulate the Th1 response, whereas estrogen is believed to enhance it [14]. Moreover, genetic or epigenetic differences between sex may also play a role as well in the observed sex-differential protective effect [13].

In summary, we have demonstrated the impact of individual-level factors on *ex vivo* mycobacterial growth inhibition in a cohort of healthy, adult volunteers. Our results indicate that immune cell phenotype, cytomegalovirus-specific response and sex have impacts on immunity following BCG vaccination. These *ex vivo* observations are reflective of epidemiological data and published human studies, and such impacts may need to be considered when testing TB vaccine candidates in trial populations. Importantly, researchers should consider the impact of sex in clinical vaccine studies, as the impact of sex in infectious diseases is common but often neglected [63]. The MGIA assay offers an *ex vivo* testing platform for assessment of a wide range of candidate TB vaccines, either using BCG or virulent *Mtb* as the immune target, with the ability to reflect inter-individual variation which may be important for vaccine effectiveness. The *ex vivo* MGIA is therefore an important additional tool for the TB vaccine community and should continue to be assessed for its ability to act as a correlate of vaccine-induced protection.

Declaration of competing interest

All authors declare no competing interests.

Acknowledgements

We would like to thank Carolynne Stanley for participant recruitment and blood sample collection, and Ayad Eddaoudi and Stephanie Canning for assistance with flow cytometry. HF has received support for

this project from EC HORIZON2020 TBVAC2020 (grant no. 643381). Satria Arief Prabowo (SP) received a PhD scholarship from the Indonesian Endowment Fund for Education (LPDP). KS was supported by the UK Medical Research Council (MRC) and the UK Department for International Development (DFID) under the MRC/DFID Concordat agreement (grant reference MR/J008702/1).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.tube.2019.101876>.

Author contributions

SP conceived and planned the experiments, supervised by HF. SP performed laboratory work and analysed the results supervised by HF, KS and SS. SP wrote the first draft of the manuscript. All authors reviewed and approved the final version of the manuscript.

References

- [1] Houben RMGJ, Dodd PJ. The global burden of latent tuberculosis infection: a re-estimation using mathematical modelling. *PLoS Med* 2016;13:e1002152.
- [2] Floyd K, Glaziou P, Zumla A, Raviglione M. The global tuberculosis epidemic and progress in care, prevention, and research: an overview in year 3 of the End TB era. *Lancet Resp Med* 2018;6:299–314.
- [3] Mangtani P, Abubakar I, Ariti C, Beynon R, Pimpin L, Fine PE, et al. Protection by BCG vaccine against tuberculosis: a systematic review of randomized controlled trials. *Clin Infect Dis: Off Publ Infect Dis Soc Am* 2014;58:470–80.
- [4] Pai M, Behr MA, Dowdy D, Dheda K, Divangahi M, Boehme CC, et al. Tuberculosis. *Nat Rev Dis Prim* 2016;2:16076.
- [5] Fine PE. Variation in protection by BCG: implications of and for heterologous immunity. *Lancet* 1995;346:1339–45.
- [6] Black GF, Weir RE, Floyd S, Bliss L, Warddorff DK, Crampin AC, et al. BCG-induced increase in interferon-gamma response to mycobacterial antigens and efficacy of BCG vaccination in Malawi and the UK: two randomised controlled studies. *Lancet* 2002;359:1393–401.
- [7] Rhodes SJ, Knight GM, Fielding K, Scriba TJ, Pathan AA, McShane H, et al. Individual-level factors associated with variation in mycobacterial-specific immune response: gender and previous BCG vaccination status. *Tuberculosis* 2016;96:37–43.
- [8] B.C.G. AND vole bacillus vaccines in the prevention of tuberculosis in adolescents; first (progress) report to the Medical Research Council by their Tuberculosis Vaccines Clinical Trials Committee. *Br Med J* 1956;1:413–27.
- [9] Mangtani P, Nguipdop-Djompo P, Keogh RH, Sterne JAC, Abubakar I, Smith PG, et al. The duration of protection of school-aged BCG vaccination in England: a population-based case-control study. *Int J Epidemiol* 2017;1–9.
- [10] World Health Organization. Global tuberculosis report. Geneva: WHO; 2017.
- [11] Neyrolles O, Quintana-Murci L. Sexual inequality in tuberculosis. *PLoS Med* 2009;6:e1000199.
- [12] Horton KC, MacPherson P, Houben RM, White RG, Corbett EL. Sex differences in tuberculosis burden and notifications in low- and middle-income countries: a systematic review and meta-analysis. *PLoS Med* 2016;13:e1002119.
- [13] Fish EN. The X-files in immunity: sex-based differences predispose immune responses. *Nat Rev Immunol* 2008;8:737–44.
- [14] Nhamoyebonde S, Leslie A. Biological differences between the sexes and susceptibility to tuberculosis. *J Infect Dis* 2014;209:S100–6.
- [15] Stensballe LG, Nante E, Jensen IP, Kofoed PE, Poulsen A, Jensen H, et al. Acute lower respiratory tract infections and respiratory syncytial virus in infants in Guinea-Bissau: a beneficial effect of BCG vaccination for girls community based case-control study. *Vaccine* 2005;23:1251–7.
- [16] Roth A, Sodemann M, Jensen H, Poulsen A, Gustafson P, Weise C, et al. Tuberculin reaction, BCG scar, and lower female mortality. *Epidemiology* 2006;17:562–8.
- [17] Stensballe LG, Sorup S, Aaby P, Benn CS, Greisen G, Jeppesen DL, et al. BCG vaccination at birth and early childhood hospitalisation: a randomised clinical multicentre trial. *Arch Dis Child* 2017;102:224–31.
- [18] Fletcher HA, Snowden MA, Landry B, Rida W, Satti I, Harris SA, et al. T-cell activation is an immune correlate of risk in BCG vaccinated infants. *Nat Commun* 2016;7:11290.
- [19] Klenerman P, Hill A. T cells and viral persistence: lessons from diverse infections. *Nat Immunol* 2005;6:873–9.
- [20] Muller J, Matsumiya M, Snowden MA, Landry B, Satti I, Harris SA, et al. Cytomegalovirus infection is a risk factor for TB disease in Infants. *bioRxiv*; 2017. <https://doi.org/10.1101/222646>. Accessed online, <https://www.biorxiv.org/content/10.1101/222646v3>.
- [21] Tanner R, O’Shea MK, Fletcher HA, McShane H. *In vitro* mycobacterial growth inhibition assays: a tool for the assessment of protective immunity and evaluation of tuberculosis vaccine efficacy. *Vaccine* 2016;0264–410X.

- [22] Brennan MJ, Tanner R, Morris S, Scriba TJ, Achkar JM, Zelmer A, et al. The cross-species mycobacterial growth inhibition assay (MGIA) project, vol 24. 2010-2014. Clinical and vaccine immunology: CVI; 2017.
- [23] Tanner R, O'Shea MK, White AD, Muller J, Harrington-Kandt R, Matsumiya M, et al. The influence of haemoglobin and iron on in vitro mycobacterial growth inhibition assays. *Sci Rep* 2017;7:43478.
- [24] Fletcher HA, Tanner R, Wallis RS, Meyer J, Manjaly ZR, Harris S, et al. Inhibition of mycobacterial growth in vitro following primary but not secondary vaccination with *Mycobacterium bovis* BCG. *Clin Vaccine Immunol : CVI* 2013;20:1683-9.
- [25] Smith SG, Zelmer A, Blitz R, Fletcher HA, Dockrell HM. Polyfunctional CD4 T-cells correlate with in vitro mycobacterial growth inhibition following *Mycobacterium bovis* BCG-vaccination of infants. *Vaccine* 2016;34:5298-305.
- [26] Joosten SA, van Meijngaarden KE, Arend SM, Prins C, Oftung F, Korsvold GE, et al. Mycobacterial growth inhibition is associated with trained innate immunity. *J Clin Invest* 2018;128:1837-51.
- [27] Naranbhai V, Hill AV, Abdool Karim SS, Naidoo K, Abdool Karim Q, Warimwe GM, et al. Ratio of monocytes to lymphocytes in peripheral blood identifies adults at risk of incident tuberculosis among HIV-infected adults initiating antiretroviral therapy. *J Infect Dis* 2014;209:500-9.
- [28] Naranbhai V, Kim S, Fletcher H, Cotton MF, Violari A, Mitchell C, et al. The association between the ratio of monocytes:lymphocytes at age 3 months and risk of tuberculosis (TB) in the first two years of life. *BMC Med* 2014;12:120.
- [29] Naranbhai V, Moodley D, Chipato T, Stranix-Chibanda L, Nakabaiito C, Kamateeka M, et al. The association between the ratio of monocytes: lymphocytes and risk of tuberculosis among HIV-infected postpartum women. *J Acquir Immune Defic Syndr* 2014;67:573-5.
- [30] Naranbhai V, Fletcher HA, Tanner R, O'Shea MK, McShane H, Fairfax BP, et al. Distinct transcriptional and anti-mycobacterial profiles of peripheral blood monocytes dependent on the ratio of monocytes: lymphocytes. *EBioMedicine* 2015; 2:1619-26.
- [31] Jensen C, Lindebo Holm L, Svensson E, Aagaard C, Ruhwald M. Optimisation of a murine splenocyte mycobacterial growth inhibition assay using virulent *Mycobacterium tuberculosis*. *Sci Rep* 2017;7:2830.
- [32] Prabowo SA, Zelmer A, Stockdale L, Ojha U, Smith SG, Seifert K, et al. Historical BCG vaccination combined with drug treatment enhances inhibition of mycobacterial growth ex vivo in human peripheral blood cells. *Sci Rep* 2019;9.
- [33] Hoft DF, Worku S, Kampmann B, Whalen CC, Ellner JJ, Hirsch CS, et al. Investigation of the relationships between immune-mediated inhibition of mycobacterial growth and other potential surrogate markers of protective *Mycobacterium tuberculosis* immunity. *J Infect Dis* 2002;186:1448-57.
- [34] Kampmann B, Tena GN, Mzazi S, Eley B, Young DB, Levin M. Novel human in vitro system for evaluating antimycobacterial vaccines. *Infect Immun* 2004;72:6401-7.
- [35] Kagina BM, Abel B, Scriba TJ, Hughes EJ, Keyser A, Soares A, et al. Specific T cell frequency and cytokine expression profile do not correlate with protection against tuberculosis after bacillus Calmette-Guerin vaccination of newborns. *Am J Respir Crit Care Med* 2010;182:1073-9.
- [36] Tanner R. Development of mycobacterial growth inhibition assays for the early evaluation and gating of novel TB vaccine candidates. Doctoral Thesis. University of Oxford; 2015.
- [37] Anwar S. Impact of helminth infection on antimycobacterial immune responses in UK migrants. Doctoral Thesis. London School of Hygiene & Tropical Medicine; 2017.
- [38] Redford PS, Murray PJ, O'Garra A. The role of IL-10 in immune regulation during *M. tuberculosis* infection. *Mucosal Immunol* 2011;4:261-70.
- [39] Tameris MD, Hatherill M, Landry BS, Scriba TJ, Snowden MA, Lockhart S, et al. Safety and efficacy of MVA85A, a new tuberculosis vaccine, in infants previously vaccinated with BCG: a randomised, placebo-controlled phase 2b trial. *Lancet* 2013;381:1021-8.
- [40] Cooper MA, Fehniger TA, Caligiuri MA. The biology of human natural killer-cell subsets. *Trends Immunol* 2001;22:633-40.
- [41] Barcelos W, Sathler-Avelar R, Martins-Filho OA, Carvalho BN, Guimaraes TM, Miranda SS, et al. Natural killer cell subpopulations in putative resistant individuals and patients with active *Mycobacterium tuberculosis* infection. *Scand J Immunol* 2008;68:92-102.
- [42] Wang R, Jaw JJ, Stutzman NC, Zou Z, Sun PD. Natural killer cell-produced IFN-gamma and TNF-alpha induce target cell cytolysis through up-regulation of ICAM-1. *J Leukoc Biol* 2012;91:299-309.
- [43] Paust S, von Andrian UH. Natural killer cell memory. *Nat Immunol* 2011;131: 500-8.
- [44] Choreno Parra JA, Martinez Zuniga N, Jimenez Zamudio LA, Jimenez Alvarez LA, Salinas Lara C, Zuniga J. Memory of natural killer cells: a new chance against *Mycobacterium tuberculosis*? *Front Immunol* 2017;8:967.
- [45] Fu X, Liu Y, Li L, Li Q, Qiao D, Wang H, et al. Human natural killer cells expressing the memory-associated marker CD45RO from tuberculous pleurisy respond more strongly and rapidly than CD45RO- natural killer cells following stimulation with interleukin-12. *Immunology* 2011;134:41-9.
- [46] Fu X, Yu S, Yang B, Lao S, Li B, Wu C. Memory-like antigen-specific human NK cells from TB pleural fluids produced IL-22 in response to IL-15 or *Mycobacterium tuberculosis* antigens. *PLoS One* 2016;11:e0151721.
- [47] Sun JC, Beilke JN, Lanier LL. Adaptive immune features of natural killer cells. *Nature* 2009;457:557-61.
- [48] Sun JC, Madera S, Bezman NA, Beilke JN, Kaplan MH, Lanier LL. Proinflammatory cytokine signaling required for the generation of natural killer cell memory. *J Exp Med* 2012;209:947-54.
- [49] Kleinnijenhuis J, Quintin J, Preijers F, Joosten LA, Jacobs C, Xavier RJ, et al. BCG-induced trained immunity in NK cells: role for non-specific protection to infection. *Clin Immunol* 2014;155:213-9.
- [50] Suliman S, Geldenhuys H, Johnson JL, Hughes JE, Smit E, Murphy M, et al. *Bacillus calmette-guerin* (BCG) revaccination of adults with latent *Mycobacterium tuberculosis* infection induces long-lived BCG-reactive NK cell responses. *J Immunol* 2016;197:1100-10.
- [51] Roy Chowdhury R, Vallania F, Yang Q, Lopez Angel CJ, Darboe F, Penn-Nicholson A, et al. A multi-cohort study of the immune factors associated with *M. tuberculosis* infection outcomes. *Nature* 2018 Aug;560(7720):644-8.
- [52] Appay V, Sauce D. Immune activation and inflammation in HIV-1 infection: causes and consequences. *J Pathol* 2008;214:231-41.
- [53] Wittkop L, Bitard J, Lazaro E, Neau D, Bonnet F, Mercie P, et al. Effect of cytomegalovirus-induced immune response, self antigen-induced immune response, and microbial translocation on chronic immune activation in successfully treated HIV type 1-infected patients: the ANRS CO3 Aquitaine Cohort. *J Infect Dis* 2013;207:622-7.
- [54] Gronborg HL, Jespersen S, Honge BL, Jensen-Fangel S, Wejse C. Review of cytomegalovirus coinfection in HIV-infected individuals in Africa. *Rev Med Virol* 2017;27.
- [55] Clari MA, Munoz-Cobo B, Solano C, Benet I, Costa E, Remigia MJ, et al. Performance of the QuantiFERON-cytomegalovirus (CMV) assay for detection and estimation of the magnitude and functionality of the CMV-specific gamma interferon-producing CD8(+) T-cell response in allogeneic stem cell transplant recipients. *Clin Vaccine Immunol: CVI* 2012;19:791-6.
- [56] Terrazzini N, Bajwa M, Vita S, Thomas D, Smith H, Vescovini R, et al. Cytomegalovirus infection modulates the phenotype and functional profile of the T-cell immune response to mycobacterial antigens in older life. *Exp Gerontol* 2014; 54:94-100.
- [57] Guma M, Angulo A, Vilches C, Gomez-Lozano N, Malats N, Lopez-Botet M. Imprint of human cytomegalovirus infection on the NK cell receptor repertoire. *Blood* 2004;104:3664-71.
- [58] Beziat V, Dalgard O, Asselah T, Halfon P, Bedossa P, Boudifa A, et al. CMV drives clonal expansion of NKG2C+ NK cells expressing self-specific KIRs in chronic hepatitis patients. *Eur J Immunol* 2012;42:447-57.
- [59] Wu Z, Sinzger C, Frascaroli G, Reichel J, Bayer C, Wang L, et al. Human cytomegalovirus-induced NKG2C(hi) CD57(hi) natural killer cells are effectors dependent on humoral antiviral immunity. *J Virol* 2013;87:7717-25.
- [60] Rhines AS. The role of sex differences in the prevalence and transmission of tuberculosis. *Tuberculosis* 2013;93:104-7.
- [61] Haralambieva IH, Ovsyannikova IG, Kennedy RB, Larrabee BR, Shane Pankratz V, Poland GA. Race and sex-based differences in cytokine immune responses to smallpox vaccine in healthy individuals. *Hum Immunol* 2013;74:1263-6.
- [62] de Bree LCJ, Koeken V, Joosten LAB, Aaby P, Benn CS, van Crevel R, et al. Non-specific effects of vaccines: current evidence and potential implications. *Semin Immunol* 2018;06. 1044-5323.
- [63] van Lunzen J, Altfeld M. Sex differences in infectious diseases-common but neglected. *J Infect Dis* 2014;209:S79-80.