



Impulsivity in the self-harm and suicidal behavior of young people: A systematic review and meta-analysis



Catherine M. McHugh^{a,b,*}, Rico Sze Chun Lee^{a,c}, Daniel F. Hermens^{a,d}, Amy Corderoy^e, Matthew Large^f, Ian B. Hickie^{a,c}

^a CRE Optimize, Brain and Mind Centre, University of Sydney, 94 Mallett street, Camperdown, NSW, 2050, Australia

^b Centre for Research Excellence in Suicide Prevention, University of Sydney, Australia

^c Brain and Mental Health Laboratory, Monash University, Melbourne, Australia

^d Sunshine Coast Mind and Neuroscience Thompson Institute, University of the Sunshine Coast, Birtinya, Australia

^e University of Notre Dame Australia, School of Medicine, Sydney, Australia

^f University of New South Wales, School of Psychiatry, Sydney, Australia

ARTICLE INFO

Keywords:

Suicidal behavior
Deliberate self-harm
Non-suicidal self-injury
Young people
Adolescent
Impulsivity
Delay discounting
Cognitive inhibition
Response inhibition

ABSTRACT

Background: Impulsivity is considered a possible phenotype underlying the expression of self-harm and suicidal behaviors. Yet impulsivity is not a unitary construct and there is evidence that different facets of impulsivity follow different neurodevelopmental trajectories and that some facets may be more strongly associated with such behaviors than others. Moreover, it is unclear whether impulsivity is a useful predictor of self-harm or suicidal behavior in young people, a population already considered to display heightened impulsive behavior.

Methods: A systematic review and meta-analysis of studies published in Medline, PubMed, PsychInfo or Embase between 1970 and 2017 that used a neurocognitive measure to assess the independent variable of impulsivity and the dependent variable of self-harm and/or suicidal behavior among young people (mean age < 30 years old).

Results: 6183 titles were identified, 141 full texts were reviewed, and 18 studies were included, with 902 young people with a self-harm or suicidal behavior and 1591 controls without a history of these behaviors. Deficits in inhibitory control (13 studies, SMD 0.21, p-value = 0.002, 95% confidence interval (CI) (0.08–0.34), prediction interval (PI) = 0.06–0.35) and impulsive decision-making (14 studies, SMD 0.17, p-value = 0.008, 95% CI (0.045–0.3), PI = 0.03–0.31) were associated with self-harm or suicidal behavior. There were no significant differences between measures of different facets of impulsivity (i.e. delay discounting, risky decision-making, cognitive or response inhibition) and self-harm or suicidal behavior.

Conclusion: Multiple facets of impulsivity are associated with suicidal behavior in young people. Future suicide research should be designed to capture impulsive states and investigate the impact on different subtypes of impulsivity.

1. Introduction

Impulsivity has emerged as a promising phenotype underlying suicidal and self-harm behavior (Mann et al., 2009). Though impulsivity has long been considered an important contributor to the expression of these behaviors, research into the association has been limited by the use of heterogeneous definitions of impulsivity (Anestis et al., 2014; Gvion 2011). Impulsivity is now widely considered ‘a predisposition toward rapid, unplanned reactions to internal or external stimuli with diminished regard to the negative consequences of these reactions to the impulsive individual or to the other’ (Moeller et al., 2001).

Historically, much of the literature examining impulsivity and self-harm or suicidal behavior has used self-report measures of impulsivity exclusively, such as the Barratt Impulsivity scale (Anestis et al., 2014; Hamza et al., 2015), though recent reviews have made distinctions between self-report and neurocognitive measures (Liu et al., 2017). Neurocognitive testing for impulsivity has been growing in favor, in part due to the ability of these tests to capture best performance through experiential testing and a growing interest in proximal, or state-based, factors associated with self-harm and suicidal behavior (Christensen et al., 2016). In a recent meta-analysis, Liu et al. (2017) found the relationship between impulsivity measured proximally

* Corresponding author. 94 Mallett street, Camperdown, NSW, 2050, Australia.
E-mail address: catherine.mchugh@sydney.edu.au (C.M. McHugh).

	Definition
Impulsivity	A predisposition toward rapid, unplanned reactions to internal or external stimuli with diminished regard to the negative consequences of these reactions to the impulsive individual or to others (Moeller, Barratt et al. 2001)
Response inhibition	The process of the overriding of a planned or already initiated action (Bari and Robbins 2013)
Cognitive inhibition	The stopping or overriding of a mental process, in whole or in part, with or without intention (MacLeod 2007)
Delay discounting	The tendency to choose small rewards that are given immediately over larger rewards that are given after a relative delay (Fineberg, Chamberlain et al. 2014) Also referred to as temporal discounting, or time-preference.
Probabilistic discounting	The tendency to choose larger but less likely option, over the smaller, more likely option. (Robbins and Dalley 2016)
Suicidal behavior	Any intentional action that may cause a person to die.
Suicide attempt	A potentially self-injurious behavior associated with at least some intent to die.
Non-suicidal self-injury	Self-injurious behavior with no intent to die.
Deliberate self-harm	Any type of self-injurious behavior, including suicide attempts and non-suicidal self-injury.
Suicidal ideation	Active: Thoughts about taking action to end one's life, including identifying a method, having a plan, or having intent to act. Passive: Thoughts about death or wanting to be dead without any plan or intent

Fig. 1. Definitions of impulsivity and suicidal behavior.

(within the last month) to suicide attempts was much stronger than attempts occurring at a more distal time point. Self-report measures, which have generally been designed to capture trait-based impulsivity, may be less objective than neurocognitive tasks in capturing impulsive states. Neurocognitive models of impulsivity have delineated several separate, but related, constructs including impulsive decision-making deficits (delay discounting and risky decision-making) and inhibitory control (response and cognitive inhibition) (Dalley and Robbins, 2017). Though definitions of impulsivity are closer to consensus than in the past, there is still no agreed set of measures used by researchers and clinicians (Fineberg et al., 2014). This review considers varying definitions of impulsivity based on the instruments used within selected studies (Fig. 1) and uses the term impulsivity to refer to impulsive decision-making deficits and deficits of inhibitory control. Importantly previous reviews have not assessed whether the association between impulsivity and self-harm and suicidal behavior is moderated by the facet of impulsivity measured (Liu et al., 2017, Hamza et al., 2015; Lockwood et al., 2017).

This review considers self-harm and suicidal behavior as a broad construct that includes a number of more specific constructs such as suicide attempts (SA), non-suicidal self-injury (NSSI) and deliberate self-harm (DSH) and uses the terms as they have been used in the primary studies. Despite evidence these behaviors are distinct from each other, in terms of intent, severity, frequency and method, previous findings have suggested these behaviors may represent differing degrees of severity within the same phenotype or differing phenotypes with significant overlap. (Muehlenkamp and Gutierrez, 2007; Hawton et al., 2012) The high degree of co-morbidity of these behaviors in young people would support such an idea (Muehlenkamp and

Gutierrez, 2007; Nock et al., 2006). Yet it is not clear whether a particular behavior, whether suicide attempt or non-suicidal self-injury, may be more strongly associated with impulsivity. Liu et al. (2017) found impulsivity more strongly associated with suicide attempts than non-suicidal self-injury. Hamza et al. (2015) found neurocognitive measures were not associated with NSSI, though both reviews included relatively few studies of NSSI.

1.1. Neurocognitive development and suicidal behavior

Deficits in executive function are associated with expression of suicidal behavior, including decision-making and problem-solving deficits, memory biases (overgeneralization, preferential recall of negatively valenced stimuli) and attentional biases (Iorfino et al., 2016; Carballo et al., 2008). Impulsive decision-making and deficits of inhibitory control of cognition and behavior have been linked with self-harm and suicidal behaviors in older age groups and with other associated psychopathology, such as substance abuse (Bickel et al., 2007; Dombrovski et al., 2011; Richard-Devantoy et al., 2012). Less is known about the role of these deficits in the suicidal behavior of young people. Cognitive and motor inhibitory control do not reach full maturity until early adulthood (Hooper et al., 2004). Decision making in developmentally normative young people is considered more impulsive than adults. This impulsivity has been characterized as being a result of higher reward dependency, greater sensation seeking and reduced tolerance for ambiguity of outcome (Steinberg, 2008; Blakemore and Robbins, 2012; Tymula et al., 2012). Thus, the question should be asked: how does the suicide and self-harm literature on impulsivity apply to young people, and in an already impulsive cohort is it useful in

identifying those at risk? Further characterizing impulsivity in young people engaging in self-harm or suicidal behaviors may be an important step in understanding the contribution of these development-related vulnerabilities to the expression of the behavior.

1.2. The stress-diathesis model: the importance of the impulsive state

While ethical and practical issues have constrained our ability to do research in populations of individuals who are acutely suicidal and distressed, there is evidence that in individuals who have very recently engaged in near fatal deliberate self-harm, intensity of suicidal ideation and intent can fluctuate rapidly (Douglas et al., 2004). Suicidal ideation and behavior have only a weak predictive ability for later completed suicide (Ribeiro et al., 2016; McHugh et al., 2019). Similarly there is evidence that suicide attempters with high trait impulsivity are as likely to plan an attempt as suicide attempters with low trait-impulsivity (May and Klonsky, 2016; Witte et al., 2008). These findings have raised important questions about the relevance of measuring suicidal intent or planning, which has been an essential component of suicide prevention in clinical settings. The limitations of this approach have driven the investigation of cognitive phenotypes, which recognize the importance of the gene-environment interaction in the development of psychopathology (Mann et al., 2009). Mann's stress-diathesis model (1999) of suicidal behavior proposes that individuals may have a trait-like disposition (diathesis) to expressing suicidal behavior when exposed to stressors, or state variables, such as symptoms of psychiatric illness, psychosocial stress or substance use.

Until recently, most research into associations between impulsivity and self-harm and suicidal behaviors have approached impulsivity as a 'state-independent' trait. It has been considered a personality trait, yet personality is now understood to be more dynamic with maladaptive features becoming accentuated in times of stress (Newton-Howes et al., 2015; McHugh and Balaratnasingam, 2018). What is not clear is how an impulsive trait may interact with impulsive cognitive states to produce suicidal behaviors.

1.2. Measures of impulsivity

1.2.1. Impulsive decision-making

Impulsive decision making can be considered to include several facets, including delay discounting, probabilistic discounting and reflection impulsivity. Delay discounting in economics refers to the individual's tendency to reduce value attached to an outcome based on time until outcome (Bickel et al., 2007). That is, the choice to accept smaller sooner rewards, rather than larger, later rewards. This is also referred to as temporal discounting or time preference. Tests of delay discounting may be self-report or experiential, the latter of which involves real monetary or food rewards. Probabilistic discounting refers to the choice between a smaller reward with higher probability of return, or larger reward, with lower probability of return. Decision-making tasks such as the Iowa Gambling Task (IGT), or the Cambridge Gambling Task (CGT) can be considered to measure probabilistic discounting, but also tap into other neurocognitive constructs such as reinforcement learning (IGT) or delay aversion (CGT) (Brevers et al., 2013; Cambridge Cognition, 2018). As these measures are not purely measures of probabilistic discounting they will be referred to as measures of risky decision-making. In this review, the term impulsive decision-making will be used to refer measures of delay discounting and these risky decision-making tasks. Explanations of neurocognitive tests of discounting discussed in this review are included in Appendix A1.

1.2.2. Inhibitory control

Inhibitory control consists of cognitive and motor inhibition and is an essential executive function. In decision making terms, having sufficient inhibitory control may allow an individual to consider value and response selection, and enable them to choose the option with the

highest objective value, or utility. Motor inhibition, hereafter referred to as response inhibition, is the failure to inhibit pre-potent responses to stimuli. Behavioral tests of response inhibition include the Go/No Go impulsivity paradigm (Dougherty et al., 2005) and the Stop signal test (SST) (Cambridge Cognition, 2018).

Cognitive inhibition, 'the stopping or overriding of a mental process, in whole or in part, with or without intention' (MacLeod, 2007), requires mediating control of attentional processes and working memory. Measures of selective and sustained attention, such as the continuous performance task (CPT), are also often used to measure this type of impulsivity (Dougherty et al., 2002). Commission errors are believed to reflect response initiation aspects of impulsivity because they result from incomplete processing of stimuli leading to rapid but incorrect responses. The Stroop test is another commonly used measure of cognitive inhibition. Explanations of neurocognitive tests of inhibitory control discussed in this review are included in Appendix A1.

The current review aims to establish whether these neurocognitive measures of impulsivity can be used to differentiate young people engaging in self-harm or suicidal behaviors from those who do not, and which, if any, sub-type of impulsivity is more strongly associated with these behaviors. We aim to establish whether age or gender moderates the relationship between impulsivity and such behaviors. A further aim of this review was to understand whether use of neurocognitive measures of impulsivity has identified evidence of impulsivity being more strongly associated with recent self-harm or suicidal behavior, rather than a lifetime history of these behaviors.

2. Methods

Methods of review including eligibility criteria, data collection and synthesis were specified in advance in the form of a review protocol and registered with PROSPERO (CRD42017081260). PRISMA guidelines for systematic review were followed (Liberati et al., 2009).

2.1. Search strategy

Medline, PubMed, PsychInfo, Embase databases were searched for relevant key words and MeSH terms which included 'suicid*' OR 'suicide, attempted' OR 'self-injurious behavior' OR 'auto-mutilation' AND 'impulsiv*' OR 'decision making' OR 'delay discounting' OR 'inhibitory control' OR 'response inhibition' OR 'executive function' OR 'cognitive control'. Studies identified from references of relevant articles were also included. Search criteria selected only studies that were reported in English and published between January 1980 and December 2017.

2.2. Eligibility criteria

Using a proforma the first author (CM) screened titles and abstracts for the following inclusion criteria: i) a mean age between 12 and 30 years, ii) compared a group with a self-harm or suicidal behavior to a group without such behaviors iii) used a neurocognitive measure of impulsivity. Self-harm or suicidal behavior was defined as any non-fatal self-injurious behavior, with intent to harm self and/or intent to end one's life, and included suicide attempts (SA), non-suicidal self-injury (NSSI) and deliberate self-harm (DSH). Terms were used as they were in the original studies. A neurocognitive test was considered any task designed to measure impulsivity. Studies with mean age between 12 and 30 years were included to capture changing impulsivity with development.

Studies were excluded if they measured suicidal ideation without behavior, had a control group with self-harm or suicidal behavior, did not report on the association between impulsivity and these behaviors or used a self-report measure of impulsivity exclusively. Articles published in languages other than English were also excluded.




Fig. 2. PRISMA flowchart.

2.3. Identification of studies

Fig. 2 outlines the process undertaken to identify studies for this review. 6183 titles were screened to determine appropriateness for screening of abstract. 333 titles and abstracts were screened by two authors (CM and AC). At that stage 192 studies were excluded based on eligibility criteria. The eligibility stage led to 141 full texts being assessed and subsequently a further 123 studies being excluded. The remaining 18 studies were included in this review.

2.4. Synthesis of results

A proforma was used to extract data from the included studies. This included information on study design (cross-sectional, longitudinal and period of follow up), sample characteristics (sample size, age, gender, population), outcome measures (NSSI, SA, or DSH), recency of self-harm or suicidal behavior (lifetime, or within 1 year), measure of impulsivity used, key findings and limitations. Key findings were evidence of an association between the impulsivity measure and outcome measure.

Neurocognitive measures were grouped together a priori based on

the facet (or subtype) of impulsivity the measure was considered to capture—delay discounting, risky decision-making, response inhibition or cognitive inhibition. Key metrics were extracted and examined to ensure appropriateness of entering into meta-analysis. Where studies reported several metrics from the same measure efforts were made to extract the same metrics from each study (ie. commission errors on the CPT, overall proportional bets on IGT) Where studies reported multiple outcome groups (ie. multiple attempts vs. single attempts separately) the two groups were pooled, due to the small number of studies reporting these as separate groups.

2.5. Assessment of study quality

Study quality was assessed using the Newcastle-Ottawa scale for case-control studies and cohort studies (Wells et al, 2018). Studies scoring 5 or less were considered to be of low quality and at greater risk of bias.

2.6. Meta-analysis

The pooled effect size across all facets of impulsivity was estimated

with a random effects meta-analysis chosen a priori for all estimates because of the diversity of study populations and differences in the methods used in the primary research. Between-study heterogeneity in effect sizes was examined using I-square, Q-value statistics and prediction intervals. The possibility of publication bias was assessed using Egger's regression (Egger et al., 1997) and was quantified using Duval and Tweedie's trim and fill method (Egger et al., 1997; Duval and Tweedie, 2000). Subgroups according to the facets of impulsivity, strength of reporting and age and sex distribution of the samples were compared using a mixed effects model. Comprehensive Meta-Analysis (CMA; Version 3, Biostat, Englewood NJ) was used for the main analysis.

3. Results

3.1. Study characteristics

Eighteen studies were included in the meta-analysis. Fourteen studies with a total of 632 subjects measured impulsive decision-making, four measured delay discounting, 3 via delay discounting (DD) tasks and 1 via Two Choice Impulsivity Paradigm (TCIP). Nine studies measured risky decision-making, 4 via IGT, 4 via CGT, 1 via the Information Sampling Task (IST).

Thirteen studies with a total of 1195 subjects measured inhibitory control. Eight studies measured response inhibition (6 via SST, 3 via Go No-go or goStop Impulsivity paradigm), six studies measured cognitive inhibition (5 via CPT, 1 via the Stroop test). Three studies used multiple measures, allowing for comparison between different neurocognitive measures of impulsivity. Table 1 summarizes the characteristics of included studies.

3.1.1. Assessment of study quality

The median NOS score of the 18 included studies was 4 (range 3–7). Additional detail is included in the supplementary material (Appendix A: 2).

3.2. Meta-analysis

Young people engaging in self-harm or suicidal behavior were significantly more impulsive than those without such behavior (standardized difference in mean 0.19, 95% CI = 0.10–0.28, p-value < 0.000, PI = 0.0934–0.29), see Fig. 3. The degree of variance was less than expected due to sampling error alone ($Q = 23.05$, $df = 26$, $I\text{-squared} = 0$). The funnel plot showed no evidence of publication bias (Appendix A: 3). Egger's regression was not significant (intercept = 0.03, t-value = 0.0034, $df = 25$, p-value = 0.49). Duval and Tweedie's trim and fill did not identify any missing hypothetical studies.

Both impulsive decision-making and deficits of inhibitory control were significantly associated with self-harm and suicidal behavior. Deficits in inhibitory control were associated with a greater effect size (SMD 0.21, p-value = 0.002, 95% CI (0.08–0.34), PI = 0.06–0.35), than impulsive decision-making (SMD 0.17, p-value = 0.01, 95% CI (0.05–0.3), PI = 0.03–0.31), though the difference between the two groups was non-significant ($Q\text{-value} = 0.13$, $df = 1$, p-value = 0.72).

Measures of risky decision-making (SMD 0.26, p-value = 0.001, 95% CI 0.11–0.42) and response inhibition (0.22, p-value = 0.005, 95% CI (0.06–0.38), differentiated young people with such behaviors from those without. Measures of cognitive inhibition (ie. CPT) (SMD 0.14, p-value = 0.24, CI -0.09-0.37) and measures of delay discounting (SMD -0.02, p-value = 0.9, CI -0.3-0.3) did not differentiate the suicidal young people from controls. Despite response inhibition and risky decision-making being significantly associated with self-harm or suicidal behavior, but not cognitive inhibition or delay discounting, subtype of impulsivity measure used did not significantly moderate the relationship between impulsivity and these behaviors ($Q\text{-value} = 3.09$, $df = 3$, p-value = 0.38).

3.2.1. Moderator analysis

None of the variables selected a priori significantly moderated the effect between impulsivity and self-harm and suicidal behavior. In some cases, subgroups reached statistical significance while others did not. See Table 2. Young people who had engaged in self-harm or suicidal behavior were significantly more impulsive than healthy controls (20 studies, SMD = 0.19, 95% CI = 0.09–0.30, p-value = 0.0002), but not psychiatric controls ($n = 7$, SMD = 0.18, 95% CI = -0.14-0.49, p-value = 0.27), though this was a non-significant difference ($Q\text{-value} = 0.01$, $df = 1$, p-value = 0.92). Study quality, whether low quality (24 studies, SMD = 0.206, 95% CI = 0.111–0.302, p-value = 0.000) or high quality (3 studies, SMD = 0.016, 95% CI = -0.315-0.348, p-value = 0.923), did not moderate the relationship between impulsivity and outcome measure.

Type of self-harm or suicidal behavior measure, whether suicide attempt ($n = 15$, SMD = 0.18, 95% CI = 0.02–0.34, p-value = 0.003) or NSSI ($n = 11$, SMD = 0.20, 95% CI = 0.07–0.33, p-value = 0.003), did not moderate the relationship between impulsivity and such behaviors ($Q\text{-value} = 0.04$, $df = 1$, p-value = 0.85). Timing of self-harm or suicidal behavior, whether lifetime history ($n = 13$, SMD = 0.17, 95% CI = 0.04–0.31, p-value = 0.02) or recent ($n = 14$, SMD = 0.21, 95% CI = 0.07–0.34, p-value = 0.002), was associated with similar effect sizes ($Q\text{-value} = 0.12$, $df = 1$, p-value = 0.73).

Age of sample was not a moderator of the relationship between impulsivity and self-harm or suicidal behavior. The relationship between impulsivity and these behaviors was similar in samples of young adults ($n = 15$, SMD = 0.19, 95% CI = 0.07–0.30, p-value = 0.002) and adolescents (age < 18 years old) ($n = 12$, SMD = 0.19, 95% CI 0.01–0.37, p-value = 0.04, $Q\text{-value} = 0.003$, $df = 1$, p-value = 0.96). Gender distribution of the sample also did not moderate the relationship. Though studies with sample sizes of more than 70% females ($n = 13$, SMD = 0.15, 95% CI = 0.004–0.29, p-value = 0.04) had a smaller effect size than those with more even gender distribution ($n = 14$, SMD = 0.23, 95% CI = 0.10–0.36, p-value < 0.001, $Q\text{-value} = 0.72$, $df = 1$, p-value = 0.40), the difference was non-significant.

4. Discussion


This review confirms that there are associations between multiple facets of impulsivity and self-harm or suicidal behaviors in young people. Thus, we have provided an important addition to previous meta-analyses which have found self-report measures of impulsivity and such behaviors are weakly associated (Anestis et al., 2014; Hamza et al., 2015) whereas neurocognitive measures of impulsivity and self-harm and suicidal behaviors are associated with a medium to large effect size (Liu et al., 2017).

While in several included studies the same sample showed differential deficits of decision-making, cognitive and response inhibition, our overall findings that subtype of impulsivity did not moderate the relationship is an important one. As evidence from the substance use disorder literature has previously suggested that delay discounting and response inhibition are separate domains (De Wit, 2009) and there is evidence that these facets are related to distinct neurocircuits (Fineberg et al., 2014) we had hypothesized that in suicidal populations there may be independence of these executive function deficits. The current review has not found any evidence of this. Although identifying discrete neurocognitive mechanisms and their associated neurocircuits may lead to further characterization of phenotypes of suicidal behavior and ultimately personalized interventions, at present the broader construct of impulsivity may be just as useful as a therapeutic target. Similarly dividing the broader construct of self-harm and suicidal behavior into more specific behaviors, suicide attempts and non-suicidal self-injury, did not moderate the relationship with impulsivity, which may indicate these behaviors are part of the same neurocognitive phenotype. Our finding that impulsivity differentiates between young people with self-

Table 1
Neurocognitive measures of impulsivity.

Study	N total (subjects, controls)	Age range in years (mean)	Population	Design	Outcome measure	Cognitive test Self-report	Findings
Ackerman et al., (2015) (Ackerman, McBee-Strayer et al., 2015)	14 SA 14 psychiatric controls	15-19 (16.9)	Community, ED F 64%	Case-control	SA Recent (1yr)	CGT	CGT scores higher in SA
Allen and Hooley (2015) (Allen and Hooley, 2015)	33 NSSI 31 healthy controls	18-53 (23.7)	Community F 75%	Case-control	NSSI Lifetime	SST	No differences between NSSI and HCs on any dimension of the SST
Bridge et al., 2012 (Bridge, McBee-Strayer et al., 2012)	40 SA 40 psychiatric controls	13-18 (15.5)	Community F 75%	Case-control	SA Recent (1yr)	IGT BIS	SA performed worse on IGT than controls, difference persisted once affective disorder, psychotropic use, self-report impulsivity and hostility adjusted for
Bridge et al., 2015 (Bridge et al., 2015)	40 SA 40 psychiatric controls	13-18 (15.6)	Community F 75%	Case-control	SA Recent (1yr)	DD	SA were more aggressive, those on medication were less aggressive, no differences on DDQ
Chamberlain et al., (2013) (Chamberlain et al., 2013)	16 SA 288 healthy controls	18-29 (21.8)	Community F 30%	Case-control	SA Lifetime	CGT	CGT performance was associated with a significant incremental benefit in predicting suicidality, over clinical and demographic variables
Fikke et al., (2011) (Fikke et al., 2011)	33 high-NSSI 29 low- NSSI 35 healthy controls	14-15 (14.8)	Schools F 71%	Case-control	NSSI Recent (1yr)	SST	Low-severity NSSI more impulsive than controls and high-severity NSSI
Gianetta et al. 2012 (Gianetta et al., 2012)	23 DSH 23 healthy controls	10-14 (11.6)	Schools F 30%	Case-control	DSH Recent (1 year)	Stroop, YSR	SI and DSH assoc. with impulsivity by self-report Cognitive measures not associated
Glenn et al. 2010 (Glenn and Klonsky, 2010)	82 NSSI 82 healthy controls	(19)	Community F 73%	Case-control	NSSI Lifetime	SST	NSSI vs. HC: no difference in Stop signal results
Grant et al. 2014 (Grant et al., 2014)	32 any suicidality 142 no suicidality	18-29 (21.7)	Community F 31%	Case-control	NSSI Lifetime	UPPS CGT	Urgency differences greater than lack of premed. or sensation seeking Suicidality associated with relative impairments in decision making and cognitive flexibility
Horesh 2001 (Horesh, 2001)	37 SA 23 psychiatric controls	(15)	Inpatient F 34%	Case-control	SA Current, recent	BIS, EIS CPT: TOVA ICS	IMP higher in suicide attempters Correlation between self-report and cog. measure low, but significant
Janis et al. 2009a (Janis and Nock, 2009)	64 NSSI 30 healthy controls	12-19 (17)	Outpatient F 80%	Case-control	NSSI Lifetime	IGT, CPT	No difference between self-harmers and controls
Janis et al. 2009b (Janis and Nock, 2009)	20 NSSI 20 healthy controls	18-30 (23-24)	Outpatient F100%	Case-control	NSSI Recent	IGT, CPT, DD K-SCADS-PL	No difference between self-harmers and controls
Mathias et al. 2011 (Mathias et al., 2011)	22 multiple SA 15 SA 22 psychiatric controls	13-17	Inpatients F 100%	Case-control	SA Lifetime	TCIP, goStop	Multiple SA > single SA > no attempts on TCIP No differences on goStop
Milner (2015) (Milner, 2015)	30 SA 34 healthy controls	18-55 (26.8)	Inpatient F 60%	Case-control	SA Recent	SST, DD, IST, CPT, Go-no go BIS, UPPS-P	No significant differences on any of the tasks
Oldershaw et al. 2009 (Oldershaw et al., 2009)	30 current DSH 24 past DSH 22 depressed controls	12-18 (15-16.5)	Outpatient Schools F 80%	Case-control	DSH Current, Lifetime	IGT	IMP higher in current DSH > past DSH > depressed controls > healthy controls Past DSH able to learn low risk strategy
Pan et al., (2011) (Pan et al., 2011)	57 healthy controls 15 SA 15 depressed controls	13-17 (16.2)	Outpatient F 57%	Case-control	SA Lifetime	Go-no go	No significant differences in any of the go-no go dimensions
Schatten et al., 2015 (Schatten et al., 2015)	14 healthy controls 48 NSSI	18-29 (21.9)	Population F 64%	Case-control	NSSI Lifetime	IGT	NSSI status not associated with IGT performance either before or after social exclusion task
Stewart et al., (2015) (Stewart et al., 2015)	59 SA 101 depressed controls	13-18 (15.6)	Inpatient F 77%	Case-control	SA Recent (1 yr)	Grad CPT	Disinhibition moderates relationship between CSA and recent suicide attempt

Meta Analysis



Meta Analysis

Fig. 3. Forest plot of impulsivity.

Table 2
Moderators of effect.

Group	Number Studies	Point estimate	Standard error	Variance	Lower limit	Upper limit	Z-value	P-value	Q-value	df	p-value
Cognitive inhibition	6.000	0.139	0.119	0.014	-0.094	0.372	1.168	0.243			
Delay Discounting	4.000	-0.019	0.149	0.022	-0.310	0.273	-0.125	0.900			
Risky decision-making	9.000	0.262	0.079	0.006	0.107	0.416	3.324	0.001			
Response inhibition	8.000	0.219	0.080	0.006	0.062	0.375	2.739	0.006			
Total between									3.093	3.00	0.377
Lifetime	13.000	0.173	0.071	0.005	0.034	0.311	2.440	0.015			
Recent	14.000	0.207	0.068	0.005	0.073	0.340	3.032	0.002			
Total between									0.119	1.00	0.730
Healthy	20.000	0.193	0.053	0.003	0.090	0.296	3.663	0.000			
Psychiatric	7.000	0.177	0.160	0.026	-0.136	0.490	1.108	0.268			
Total between									0.009	1.00	0.926
Adolescent	12.000	0.192	0.092	0.008	0.011	0.372	2.082	0.037			
Adult	15.000	0.186	0.060	0.004	0.069	0.303	3.109	0.002			
Total between									0.003	1.00	0.956
Less	14.000	0.230	0.066	0.004	0.102	0.358	3.511	0.000			
Over 70% female	13.000	0.147	0.073	0.005	0.004	0.290	2.011	0.044			
Total between									0.721	1.000	0.396
Impulsive decision-making	14.000	0.173	0.065	0.004	0.045	0.300	2.657	0.008			
Inhibitory control	13.000	0.206	0.066	0.004	0.076	0.336	3.110	0.002			
Total between									0.128	1.00	0.721
NSSI	11.000	0.197	0.066	0.004	0.068	0.327	2.992	0.003			
SA	15.000	0.177	0.081	0.007	0.019	0.336	2.191	0.028			
Total between									0.036	1.00	0.849
High quality	3	0.016	0.169	0.029	-0.315	0.348	0.097	0.923			
Low quality	24	0.206	0.049	0.002	0.111	0.302	4.234	0.000			
Total within									1.159	1.00	0.282

harm and suicidal behavior and healthy controls, but not psychiatric controls must be interpreted with caution. Few studies used psychiatric control groups, such that the number of psychiatric controls in the meta-analytic sample was far fewer than healthy controls (161 vs. 1430 respectively). Thus our meta-analysis was likely underpowered to detect an effect between suicidal and self-harming young people and psychiatric controls.

4.1. Risky decision-making predicts self-harm and suicidal behaviors

Risky decision-making tasks such as the IGT and CGT, but not delay discounting tasks, differentiated young people with self-harm or suicidal behavior from controls. The majority of studies of risky decision-making used the Iowa Gambling task (IGT), which is often considered to measure multiple executive functions, such as reinforcement learning, in addition to probabilistic discounting, which may confound interpretation of our findings. Nonetheless these results are consistent with findings in adult populations (Gorlyn et al., 2013) and add to the

existing literature as other meta-analysis have not included these decision-making tasks (Liu et al., 2017).

Contrary to our hypothesis deficits of delay discounting did not differentiate young people with self-harm or suicidal behavior from controls. However, only a small number of studies (4) measured delay discounting, which suggests the analysis may have been underpowered in testing our hypothesis. Discounting is sensitive to a number of different conditions including order of magnitude (i.e. the larger the amount, the more steeply people discount), sign effect (positive outcomes discounted more than negative outcomes), duration (distant outcomes discounted more than proximal) and affective states (in 'hot' affective states persistence of current preferences overestimated) (Tucker et al., 2010). One potential model of decision-making in self-harm and suicidal behavior may involve the individual under stress valuing the smaller sooner rewards (ie. cessation of emotional pain) over the larger, later rewards that may come with continuing living. The current value of life may be suddenly decreased due to loss of social role or connection. Put alternatively the costs associated with these behaviors in the present, such as physical pain and fear, may be less than perceived costs of continued living with emotional pain. This is consistent with evidence from healthy populations that individuals in 'hot' affective states, such as in emotional crisis, are likely to overestimate the likelihood that current conditions will persist (Madden and Johnson, 2010). Under this model of self-harm and suicidal decision-making the individual may believe the immediate choice of the self-harm or suicidal act is the option with the highest utility. Other hallmarks of the suicidal crisis, including intoxication and sleeplessness have all been shown to cause steeper discounting of future gain (Tucker et al., 2010).

4.2. Deficits in inhibitory control are associated with self-harm and suicidal behavior in young people

Young people with self-harm or suicidal behavior were significantly slower in responding appropriately to stimulus in response inhibition tasks relative to controls. This finding is in keeping with current psychological models of suicidal behavior. A deficit in response inhibition, which is the motor, or behavioral component of inhibitory control may explain why some individuals engaging in self-harm and suicidal behavior describe having engaged in the act before they have had a chance to stop themselves (Skegg, 2005).

Measures of cognitive inhibition, however, did not differentiate young people with self-harm and suicidal behavior from controls, contrary to our hypothesis. Only six studies measured cognitive inhibition, which likely makes our analysis underpowered. We had hypothesized that deficits in cognitive inhibition, or the ability to override mental processes, may have particular effects on the suicidal mind, resulting in increased ideation, a predictor of behavior (Nock et al., 2010). Once an individual is in crisis they may be unable to shift attention from negative thoughts or thoughts of self-harm or suicide, which is likely to lead to increased frequency and perceived lack of control over ideas of self-harm or suicide. Suicidal ideation and behavior scales that measure frequency and control over suicidal thoughts have previously been shown to be more robust measures of suicidal ideation, relative to dichotomous descriptors (Batterham et al., 2015). Deficits of cognitive inhibition may be related to rumination, which has also been associated with these behaviors (Rogers and Joiner, 2017).

4.3. The role of state-based impulsivity is poorly understood

Only two studies used individuals who had self-harmed or made a suicide attempt within the last week to month (Horeh, 2001; Oldershaw et al., 2009) and less than half the studies recorded suicide attempts or self-harm within the previous year (Janis and Nock, 2009; Stewart et al., 2015; Ackerman et al., 2015; Bridge et al.; Bridge et al., 2015; Fikke et al., 2011; Giannetta et al., 2012). The only study that

compared current and former self-harmers, found adolescents who were current self-harmers made more high-risk choices on IGT than those with a history of self-harm, or healthy controls (Oldershaw et al., 2009). Current self-harmers were the only group who did not improve over the course of the task. This deficit of studies examining impulsivity and cognition proximally to self-harm and suicidal behavior in young people is particularly important, given the findings from Liu et al's (2017) meta-analysis that in adults the odds of impulsivity in recent (< 1 month) suicide attempters was greatly increased.

It is possible that decision-making processes are more state based than measures of inhibitory control, which would mean the methodological limitations of the literature may mask the true effect of these deficits on self-harm and suicidal behavior. Indeed previous reviews have found inhibitory control may be less impacted by stress conditions than other executive functions (Shields et al., 2016). Discounting of reward has been shown to be strongly impacted by other state variables, including both acute intoxication and chronic substance use (Bickel et al., 1999). Certainly a growing body of evidence suggests conditions of stress may significantly impact decision-making and inhibitory control in healthy individuals. It will be vital for research to get closer in time to self-harm and suicidal behaviors and measure associated state-based variables. For example, in this review few studies controlled for current or past substance use. Traditionally, measuring cognitive states has been difficult, in both clinical and research settings. Advances in technology, however, have increased both the accessibility and sophistication of neurocognitive testing, such as CanTab or Cogstat (Fray et al., 1996; Maruff et al., 2009). With these advances it may be possible to measure suicidal cognition in real-time, outside the domains of the clinical or research setting (Nock et al., 2009). Using measures that quantify state differences are likely to be more useful in understanding etiology, recognizing risk clinically and in prevention.

4.4. Impulsivity and self-harm or suicidal behavior in development

Impulsivity is associated with suicidal behavior in young people. Despite impulsive choice and behavior being a feature of normative development, deficits of inhibitory control and discounting still meaningfully differentiated groups of young people with self-harm or suicidal behavior from those without. While previous reviews have examined facets of impulsivity in development of healthy adolescents and populations of substance using young people, less is known about changing impulsivity in young people engaging in self-harm or suicidal behaviors. This analysis found the average age of the sample, either over 18 years or below, did not moderate the association between impulsivity and these behaviors. It is possible that dividing age dichotomously introduces error into the analysis. Examining age associations in meta-analysis can be problematic as each study generally reports the mean age of the sample, rather than for cases and controls. No individual studies in the current review had a study design sufficient to comment on changing impulsivity with age, which will be an important question for future research.

Half of the included studies had a sample of greater than 70% female, which suggests there may be sampling bias in the literature on impulsivity and self-harm and suicidal behavior. If sex is associated with either delayed development of inhibitory control or decision-making, the current literature may have underestimated the relationship between these behaviors and impulsivity due to sampling bias. Self-harm, with or without suicidal intent, is considered to be more prevalent in females (Skegg, 2005). Cultural difference in expression of emotion, and biological factors (ie. testosterone and aggression), have both been hypothesized to contribute to such an effect (Bresin and Schoenleber, 2015). Yet studies using broader definitions of self-harm, have shown less of a gender difference in prevalence, which suggests methodological issues in suicide research may at least in part contribute to this finding (Muehlenkamp and Gutierrez, 2004). Rippon et al. (2014) have outlined the way in which methodological error (ie.

inappropriate consideration of age and development) can confound relationships between sex and various outcomes in the neurosciences. Gender bias in sampling is particularly important to consider in the study of self-harm and suicidal behavior.

4.5. Limitations

The current review had a limited ability to look at important moderators, including other executive function deficits, and potential confounders such as substance use. This reflects the relatively small number of studies examining neurocognitive measures of impulsivity in young people with self-harm or suicidal behavior. We were only able to broadly examine the role of individual characteristics, such as age or gender, at a study level, which likely introduces bias. There were relatively few studies reporting facets of impulsivity, which likely resulted in the meta-analysis being underpowered to detect an effect. It is still not clear how to measure clinically meaningful differences in impulsive decision-making and inhibitory control. This makes it difficult to know whether the findings of this review are clinically meaningful.

5. Conclusion

The current review has demonstrated that impulsivity differentiates young people engaging in self-harm or suicidal behavior from healthy controls. This relationship exists despite the age or gender of the sample. There is limited evidence of a differential association between particular facets of impulsivity and the expression of such behaviors in young people. Deficits of response inhibition and riskier decision-making were associated with these behaviors, though deficits of cognitive inhibition and steeper delay discounting were not significantly associated. Recency of self-harm or suicidal behavior or type of behavior measured did not moderate the relationship. Future research into self-harm and suicidal behaviors and decision making should focus on state impulsivity by studying decision making processes in populations with recent self-harm or suicidal behavior. Greater understanding of trajectories of impulsive decision-making and behavior across adolescence and adulthood would increase our understanding of the contribution of this important executive function in expression of self-harm and suicidal behavior.

Funding

This work was supported by the Royal Australian and New Zealand College of Psychiatry New Investigator Grant.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jpsychires.2019.05.012>.

References

Ackerman, J.P., McBee-Strayer, S.M., Mendoza, K., et al., 2015. Risk-sensitive decision-making deficit in adolescent suicide attempters. *J. Child Adolesc. Psychopharmacol.* 25, 109–113.

Allen, K.J., Hooley, J.M., 2015. Inhibitory control in people who self-injure: evidence for impairment and enhancement. *Psychiatr. Res.* 225 (3), 631–637.

Anestis, M.D., Soberay, K.A., Gutierrez, P.M., et al., 2014. Reconsidering the link between impulsivity and suicidal behavior. *Pers. Soc. Psychol. Rev.* 18, 366–386.

Batterham, P.J., Ftanou, M., Pirkis, J., et al., 2015. A systematic review and evaluation of measures for suicidal ideation and behaviors in population-based research. *Psychol. Assess.* 27, 501.

Bickel, W.K., Miller, M.L., Yi, R., et al., 2007. Behavioral and neuroeconomics of drug addiction: competing neural systems and temporal discounting processes. *Drug Alcohol Depend.* 90 (Suppl. 1), S85–S91.

Bickel, W.K., Odum, A.L., Madden, G.J., 1999. Impulsivity and cigarette smoking: delay discounting in current, never, and ex-smokers. *Psychopharmacology* 146, 447–454.

Blakemore, S.J., Robbins, T.W., 2012. Decision-making in the adolescent brain. *Nat. Neurosci.* 15, 1184–1191.

Bresin, K., Schoenleber, M., 2015. Gender differences in the prevalence of nonsuicidal

self-injury: a meta-analysis. *Clin. Psychol. Rev.* 38, 55–64.

Brevers, D., Bechara, A., Cleeremans, A., Noël, X., 2013. Iowa Gambling Task (IGT): twenty years after – gambling disorder and IGT. *Front. Psychol.* 4, 665.

Bridge, J.A., McBee-Strayer, S.M., Cannon, E.A., et al., 2012. Impaired decision making in adolescent suicide attempters. *J. Am. Acad. Child Adolesc. Psychiatry* 51 (4), 394–403.

Bridge, J.A., Reynolds, B., McBee-Strayer, S.M., et al., 2015. Impulsive aggression, delay discounting, and adolescent suicide attempts: effects of current psychotropic medication use and family history of suicidal behavior. *J. Child Adolesc. Psychopharmacol.* 25, 114–123.

Carballo, J.J., Akamnonu, C.P., Oquendo, M.A., 2008. Neurobiology of suicidal behavior. An integration of biological and clinical findings. *Arch. Suicide Res.* 12, 93–110.

Chamberlain, S.R., Odlaug, B.L., Schreiber, L.R.N., Grant, J.E., 2013. Clinical and neurocognitive markers of suicidality in young adults. *J. Psychiatr. Res.* 47 (5), 586–591.

Christensen, H., Cuijpers, P., Reynolds, C.F., et al., 2016. Changing the direction of suicide prevention research: a necessity for true population impact. *JAMA Psychiatr.* 73, 435–436.

Cambridge Cognition, 2018. CANTAB® [Cognitive Assessment Software]. All rights reserved. www.cantab.com.

Dalley, J.W., Robbins, T.W., 2017. Fractionating impulsivity: neuropsychiatric implications. *Nat. Rev. Neurosci.* 18, 158–171.

De Wit, H., 2009. Impulsivity as a determinant and consequence of drug use: a review of underlying processes. *Addict. Biol.* 14, 22–31.

Dombrowski, A.Y., Szanto, K., Siegle, G.J., et al., 2011. Lethal forethought: delayed reward discounting differentiates high-and low-lethality suicide attempts in old age. *Biol. Psychiatry* 70, 138–144.

Dougherty, D.M., Marsh, D.M., Mathias, C.W., 2002. Immediate and delayed memory tasks: a computerized behavioral measure of memory, attention, and impulsivity. *Behav. Res. Methods Instrum. Comput.* 34, 391–398.

Dougherty, D.M., Mathias, C.W., Marsh, D.M., et al., 2005. Laboratory behavioral measures of impulsivity. *Behav. Res. Methods* 37, 82–90.

Douglas, J., Cooper, J., Amos, T., et al., 2004. "Near-fatal" deliberate self-harm: characteristics, prevention and implications for the prevention of suicide. *J. Affect. Disord.* 79, 263–268.

Duval, S., Tweedie, R., 2000. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* 56, 455–463.

Egger, M., Smith, G.D., Schneider, M., et al., 1997. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 315, 629–634.

Fikke, L.T., Melinder, A., Landro, N.I., 2011. Executive functions are impaired in adolescents engaging in non-suicidal self-injury. *Psychol. Med.* 41, 601–610.

Fineberg, N.A., Chamberlain, S.R., Goudriaan, A.E., Stein, D.J., Vanderschuren, L.J.M.J., Gillan, C.M., Shekar, S., Gorwood, P.A.P.M., Voon, V., Morein-Zamir, S., Denys, D., Sahakian, B.J., Moeller, F.G., Robbins, T.W., Potenza, M.N., 2014. New developments in human neurocognition: clinical, genetic and brain imaging correlates of impulsivity and compulsivity. *CNS Spectr.* 19 (1), 69–89.

Fray, P.J., Robbins, T.W., Sahakian, B.J., 1996. Neuropsychiatric applications of CANTAB. *Int. J. Geriatr. Psychiatry* 11, 329–336.

Giannetta, M.M., Betancourt, L.M., Brodsky, N.L., et al., 2012. Suicidal ideation and self-harm behavior in a community sample of preadolescent youth: a case-control study. *J. Adolesc. Health* 50, 524–526.

Glenn, C.R., Klonsky, E., 2010. A multimethod analysis of impulsivity in nonsuicidal self-injury. *Personal. Disord.: Theor. Res. Treat.* 1 (1), 67–75.

Gorlyn, M., Keilp, J.G., Oquendo, M.A., et al., 2013. Iowa Gambling Task performance in currently depressed suicide attempters. *Psychiatr. Res.* 207, 150–157.

Grant, J.E., Derbyshire, K., Leppink, E., Chamberlain, S.R., 2014. Suicidality in non-treatment seeking young adults with subsyndromal gambling disorder. *Psychiatr. Q.* 85 (4), 513–522.

Gvion, Y., Apter, A., 2011. Aggression, impulsivity, and suicide behavior: a review of the literature. *Archives of Suicide Research* 15 (2), 93–112.

Hamza, C.A., Willoughby, T., Heffer, T., 2015. Impulsivity and nonsuicidal self-injury: a review and meta-analysis. *Clin. Psychol. Rev.* 38, 13–24.

Hawton, K., Saunders, K.E., O'Connor, R.C., 2012. Self-harm and suicide in adolescents. *The Lancet* 379 (9834), 2373–2382.

Hooper, C.J., Luciana, M., Conklin, H.M., et al., 2004. Adolescents' performance on the Iowa Gambling Task: implications for the development of decision making and ventromedial prefrontal cortex. *Dev. Psychol.* 40, 1148.

Hoshish, N., 2001. Self-report vs. computerized measures of impulsivity as a correlate of suicidal behavior. *Crisis: J. Crisis Interv. Suicide* 22, 27–31.

Iorfino, F., Hickie, I.B., Lee, R.S.C., et al., 2016. The underlying neurobiology of key functional domains in young people with mood and anxiety disorders: a systematic review. *BMC Psychiatry* 16.

Janis, I.B., Nock, M.K., 2009. Are self-injurers impulsive?: results from two behavioral laboratory studies. *Psychiatr. Res.* 169, 261–267.

Liberati, A., Altman, D.G., Tetzlaff, J., et al., 2009. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med.* 6, e1000100.

Liu, R.T., Trout, Z.M., Hernandez, E.M., Cheek, S.M., Gerlus, N., 2017. A behavioral and cognitive neuroscience perspective on impulsivity, suicide, and non-suicidal self-injury: meta-analysis and recommendations for future research. *Neurosci. Biobehav. Rev.* 83, 440–450.

Lockwood, J., Daley, D., Townsend, E., Sayal, K., 2017. Impulsivity and self-harm in adolescence: a systematic review. *Eur. Child Adolesc. Psychiatry* 26 (4), 387–402.

MacLeod, C.M., 2007. The concept of inhibition in cognition. *Inhib. Cognit.* 3–23.

Madden, G., Johnson, 2010. A delay-discounting primer. In: Madden, G., Bickel, W. (Eds.), *Impulsivity: the Behavioural and Neurological Science of Discounting*. American Psychological Association, Washington.

- Mann, J.J., Arango, V.A., Avenevoli, S., et al., 2009. Candidate endophenotypes for genetic studies of suicidal behavior. *Biol. Psychiatry* 65, 556–563.
- Maruff, P., Thomas, E., Cysique, L., et al., 2009. Validity of the CogState brief battery: relationship to standardized tests and sensitivity to cognitive impairment in mild traumatic brain injury, schizophrenia, and AIDS dementia complex. *Arch. Clin. Neuropsychol.* 24, 165–178.
- Mathias, C.W., Dougherty, D.M., James, L.M., Richard, D.M., Dawes, M.A., Acheson, A., Hill-Kapturczak, N., 2011. Intolerance to delayed reward in girls with multiple suicide attempts. *Suicide Life-Threatening Behav.* 41 (3), 277–286.
- May, A.M., Klonsky, E.D., 2016. "Impulsive" suicide attempts: what do we really mean? *Personal Disord.* 7, 293–302.
- McHugh, C., Balaratnasingam, S., 2018. Impulsivity in personality disorders: current views and future directions. *Curr. Opin. Psychiatr.* 31, 63–68.
- McHugh, C., Corderoy, A., Ryan, C., Hickie, I., Large, M., 2019. Association between suicidal ideation and suicide: meta-analyses of odds ratios, sensitivity, specificity and positive predictive value. *BJPsych Open* 5 (2), E18. <https://doi.org/10.1192/bjo.2018.88>.
- Milner, A.J., 2015. Clarifying the Pathway to Suicide: an Examination of Subtypes of Suicidal Behaviour and Their Association with Impulsiveness. Doctor of Philosophy. Harvard University.
- Moeller, F.G., Barratt, E.S., Dougherty, D.M., et al., 2001. Psychiatric aspects of impulsivity. *Am. J. Psychiatry* 158, 1783–1793.
- Muehlenkamp, J.J., Gutierrez, P.M., 2004. An investigation of differences between self-injurious behavior and suicide attempts in a sample of adolescents. *Suicide Life-Threatening Behav.* 34, 12–23.
- Muehlenkamp, J.J., Gutierrez, P.M., 2007. Risk for suicide attempts among adolescents who engage in non-suicidal self-injury. *Arch. Suicide Res.* 11 (1), 69–82.
- Newton-Howes, G., Clark, L.A., Chanen, A., 2015. Personality disorder across the life course. *The Lancet* 385, 727–734.
- Nock, M.K., Joiner Jr., T.E., Gordon, K.H., Lloyd-Richardson, E., Prinstein, M.J., 2006. Non-suicidal self-injury among adolescents: diagnostic correlates and relation to suicide attempts. *Psychiatr. Res.* 144 (1), 65–72.
- Nock, M.K., Park, J.M., Finn, C.T., et al., 2010. Measuring the suicidal mind. *Psychol. Sci.* 21, 511–517.
- Nock, M.K., Prinstein, M.J., Sterba, S.K., 2009. Revealing the form and function of self-injurious thoughts and behaviors: a real-time ecological assessment study among adolescents and young adults. *J. Abnorm. Psychol.* 118, 816–827.
- Oldershaw, A., Grima, E., Jollant, F., et al., 2009. Decision making and problem solving in adolescents who deliberately self-harm. *Psychol. Med.* 39, 95–104.
- Pan, L.A., Batezati-Alves, S.C., Almeida, J.R., Segreti, A., Akkal, D., Hassel, S., Lakdawala, S., Brent, D.A., Phillips, M.L., 2011. Dissociable patterns of neural activity during response inhibition in depressed adolescents with and without suicidal behavior. *J. Am. Acad. Child Adolesc. Psychiatry* 50 (6), 602–611 e603.
- Ribeiro, J., Franklin, J., Fox, K.R., et al., 2016. Self-injurious thoughts and behaviors as risk factors for future suicide ideation, attempts, and death: a meta-analysis of longitudinal studies. *Psychol. Med.* 46, 225–236.
- Richard-Devantoy, S., Gorwood, P., Annweiler, C., et al., 2012. Suicidal behaviours in affective disorders: a deficit of cognitive inhibition? *Can. J. Psychiatr. - Rev. Can. Psychiatr.* 57, 254–262.
- Rippon, G., Jordan-Young, R., Kaiser, A., et al., 2014. Recommendations for sex/gender neuroimaging research: key principles and implications for research design, analysis, and interpretation. *Front. Hum. Neurosci.* 8, 650.
- Rogers, M.L., Joiner, T.E., 2017. Ruminative, Suicidal Ideation, and Suicide Attempts: A Meta-Analytic Review. Educational Publishing Foundation.
- Schatten, H.T., Andover, M.S., Armev, M.F., 2015. The roles of social stress and decision-making in non-suicidal self-injury. *Psychiatr. Res.* 229 (3), 983–991.
- Shields, G.S., Sazma, M.A., Yonelinas, A.P., 2016. The effects of acute stress on core executive functions: a meta-analysis and comparison with cortisol. *Neurosci. Biobehav. Rev.* 68, 651–668.
- Skegg, K., 2005. Self-harm. *Lancet* 366, 1471–1483.
- Steinberg, L., 2008. A social neuroscience perspective on adolescent risk-taking. *Dev. Rev.* : DR (Dev. Rev.) 28, 78–106.
- Stewart, J.G., Kim, J.C., Esposito, E.C., et al., 2015. Predicting suicide attempts in depressed adolescents: clarifying the role of disinhibition and childhood sexual abuse. *J. Affect. Disord.* 187, 27–34.
- Tucker, J., Simpson, C., Khodneva, J., 2010. Role of time and delay in health decision making. In: Madden, G., Bickel, W. (Eds.), *Impulsivity: the Behavioural and Neurological Science of Discounting*. American Psychological Association, Washington.
- Tymula, A., Rosenberg Belmaker, L.A., Roy, A.K., et al., 2012. Adolescents' risk-taking behavior is driven by tolerance to ambiguity. *Proc. Natl. Acad. Sci. U. S. A.* 109, 17135–17140.
- Wells, G., Shea, B., O'Connell, D., Peterson, J., Welch, V., Losos, M., 2018. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomized studies in meta-analysis. University of Ottawa, Ottawa, pp. 2008.
- Witte, T.K., Merrill, K.A., Stellrecht, N.E., et al., 2008. "Impulsive" youth suicide attempters are not necessarily all that impulsive. *J. Affect. Disord.* 107, 107–116.

Further reading

- Bari, A., Robbins, T.W., 2013. Inhibition and impulsivity: behavioral and neural basis of response control. *Prog. Neurobiol.* 108, 44–79.
- Cognition, C., 2016. CANTAB Stop Signal Task. Available at: <http://www.cambridgecognition.com/tests/stop-signal-task-sst>.