



Individuals with 22q11.2 deletion syndrome show intact prediction but reduced adaptation in responses to repeated sounds: Evidence from Bayesian mapping

Kit Melissa Larsen^{a,b,c,d,e,*}, Morten Mørup^b, Michelle Rosgaard Birknow^{c,d,f}, Elvira Fischer^a, Line Olsen^{c,d}, Michael Didriksen^f, William Frans Christiaan Baaré^a, Thomas Mears Werge^{c,d,j}, Marta Isabel Garrido^{e,g,h,i,1}, Hartwig Roman Siebner^{a,j,k,1}

^a Danish Research Centre for Magnetic Resonance, Centre for Functional and Diagnostic Imaging and Research, Copenhagen University Hospital Hvidovre, Hvidovre, Denmark

^b DTU Compute, Cognitive Systems, Technical University of Denmark, Lyngby, Denmark

^c Institute of Biological Psychiatry, Mental Health Centre Sct. Hans, Copenhagen University Hospital, Boserupvej 2, DK-4000 Roskilde, Denmark

^d iPSYCH, The Lundbeck Foundation Initiative for Integrative Psychiatric Research, Aarhus and Copenhagen, Denmark

^e Queensland Brain Institute, The University of Queensland, St Lucia, 4072 Brisbane, Australia

^f Synaptic Transmission, H. Lundbeck A/S, Ottiliavej 9, DK-2500, Valby, Denmark

^g Centre for Advanced Imaging, The University of Queensland, St Lucia, 4072 Brisbane, Australia

^h Australian Research Council Centre of Excellence for Integrative Brain Function Centre of Excellence for Integrative Brain Function, The University of Queensland, St Lucia, 4072 Brisbane, Australia

ⁱ School of Mathematics and Physics, The University of Queensland, St Lucia, 4072 Brisbane, Australia

^j Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

^k Department of Neurology, Copenhagen University Hospital Bispebjerg, Copenhagen, Denmark

ARTICLE INFO

Keywords:

22q11 deletion syndrome
Dynamic causal modelling
Posterior probability maps
EEG
Mismatch negativity
Repetition suppression

ABSTRACT

One of the most common copy number variants, the 22q11.2 microdeletion, confers an increased risk for schizophrenia. Since schizophrenia has been associated with an aberrant neural response to repeated stimuli through both reduced adaptation and prediction, we here hypothesized that this may also be the case in non-psychotic individuals with a 22q11.2 deletion.

We recorded high-density EEG from 19 individuals with 22q11.2 deletion syndrome (12–25 years), as well as 27 healthy volunteers with comparable age and sex distribution, while they listened to a sequence of sounds arranged in a roving oddball paradigm. Using posterior probability maps and dynamic causal modelling we tested three different models accounting for repetition dependent changes in cortical responses as well as in effective connectivity; namely an adaptation model, a prediction model, and a model including both adaptation and prediction.

Repetition-dependent changes were parametrically modulated by a combination of adaptation and prediction and were apparent in both cortical responses and in the underlying effective connectivity. This effect was reduced in individuals with a 22q11.2 deletion and was negatively correlated with negative symptom severity. Follow-up analysis showed that the reduced effect of the combined adaptation and prediction model seen in individuals with 22q11.2 deletion was driven by reduced adaptation rather than prediction failure. Our findings suggest that adaptation is reduced in individuals with a 22q11.2 deletion, which can be interpreted in light of the framework of predictive coding as a failure to suppress prediction errors.

Abbreviations: DCM, Dynamic Causal Modelling; MMN, Mismatch Negativity; 22q11.2DS, 22q11.2 Deletion Syndrome

* Corresponding author at: Queensland Brain Institute, The University of Queensland, St Lucia, 4072 Brisbane, Australia.

E-mail address: m.larsen@uq.edu.au (K.M. Larsen).

¹ These authors contributed equally to the study.

<https://doi.org/10.1016/j.nicl.2019.101721>

Received 24 October 2018; Received in revised form 23 January 2019; Accepted 12 February 2019

Available online 13 February 2019

2213-1582/ © 2019 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

22q11.2 deletion syndrome (22q11.2DS) is caused by one of the most common copy number variants in humans with a prevalence of 1:2000 to 1:4000 (Goodship et al., 1998; Olsen et al., 2018; Oskarsdóttir et al., 2004; Shprintzen, 2005). The 22q11.2DS is clinically presented with a highly variable phenotype, including a range of somatic disorders, learning problems, and cognitive deficits (Karayiorgou et al., 2010; Robin and Shprintzen, 2005). 22q11.2DS is associated with a high frequency of several neurodevelopmental disorders, including autism spectrum disorder, attention deficit hyperactivity disorder and schizophrenia (Bassett et al., 2008; Hoeft et al., 2017; Karayiorgou et al., 2010; Purcell et al., 2009; Schneider et al., 2014; Stefansson et al., 2008).

The ability to adapt to the ever changing environment and react to deviations within it, is something the healthy brain masters on a daily basis. However, people with schizophrenia show reduced ability to adapt to the environment expressed as a state of aberrant salience (Kapur, 2003). A neurophysiological example of this phenomenon is the typically reduced neural response to repeated stimuli, a process called repetition suppression (RS), often depicted as a consequence of neural fatigue (Grill-Spector et al., 2006). Recent theoretical formulations inspired on predictive coding (Aukstulewicz and Friston, 2016; Friston, 2005; Grotheer and Kovács, 2016) propose that altered RS in schizophrenia may be caused by inaccurate sensory predictions (Baldeweg, 2007; Summerfield et al., 2008; Todorovic and de Lange, 2012; Todorovic et al., 2011). According to this perspective, RS is a consequence of prediction error minimization afforded by adaptation to the environment through learning about incoming sensory input. Neurobiological, repetition-dependent changes in responses to repeated stimuli have previously been explained by experience-dependent changes in effective connectivity (Garrido et al., 2009). These changes in effective connectivity were evident both in extrinsic connections, (between brain areas) as well as in intrinsic connections (within brain area). Extrinsic connections are believed to encode prediction while intrinsic connections are believed to encode prediction precision. Most event-related potential (ERP) studies have focused on the first repetition relative to the initial presentation (Andrade et al., 2015; Henson et al., 2004; Mayer et al., 2009; Stefanics et al., 2018). Very recent, Stefanics et al. (2018) showed that RS was best explained by an exponential model indicating that repetition effects are observable for trials beyond the first repetition, highlighting the necessity to investigate brain responses beyond the first repetition in order to understand the underlying processes of RS.

RS in schizophrenia has mostly been studied through sensory gating where suppression of P50 is seen to be reduced (Adler et al., 1982; Rentsch et al., 2015). Haenschel et al. (2005) found a repetition-dependent enhancement of a slow positive wave in responses to repeated sounds in a roving oddball paradigm (healthy volunteers), called repetition positivity. In another study Baldeweg et al. (2004) showed that individuals with schizophrenia do not show this repetition positivity, suggesting that adaptive cortical processes are impaired in schizophrenia. The MMN slope (across repetitions) was directly correlated with short- and long-term memory as well as disease severity. However, RS has also been shown to be intact in schizophrenia (Coffman et al., 2017; Rosburg, 2018) manifest in comparable ERPs to repeated auditory tones (Potter et al., 2005). RS is sparsely studied in 22q11.2DS with again opposing results where sensory gating as indexed by P50 has been shown to be sometimes intact (Rihs et al., 2013; Vorstman et al., 2009) and other times impaired (Larsen et al., 2018a; Zarchi et al., 2013). Given that the underlying mechanism of RS in 22q11.2DS is still poorly understood, we investigated the brain mechanism underpinning RS in 22q11.2DS and how these mechanisms might potentially deviate from what is seen in healthy controls. We formalized three theoretical models to explain RS: the adaptation model, the prediction model and finally the combined model, in the following referred to as adaptation&

prediction. These three models were tested both at the scalp level using Bayesian mapping for M/EEG (Garrido et al., 2018; Harris et al., 2018; Rosa et al., 2010) and at the connectivity level using dynamical causal modelling (DCM) (David et al., 2006). Firstly, we hypothesized that responses to repeated stimuli would show a parametric modulation with an overall decrease in connectivity within the tested network for the first repetitions, followed by an increase reflecting the prediction of new stimuli, in agreement with the combined adaptation&prediction model. Next, we investigated group differences in 22q11.2DS and healthy controls at the scalp level and connectivity level within the model that best described RS. This unique way of modelling RS allowed us to pinpoint the origin of potential deficits in 22q11.2DS, namely the adaptive and predictive processes underpinning RS.

2. Materials and methods

This paper is a follow up study of (Larsen et al., 2018b), where we report on mismatch negativity (MMN) responses in 22q11.2DS. Participants' demographics and stimuli administered is described in Larsen et al. (2018b) and a summary can be found below for clarity.

2.1. Participants

This study is part of a larger Danish nationwide initiative (Schmoeck et al., 2015). A group of 19 non-psychotic individuals with a verified deletion of 3Mb at chromosome 22q11.2 with no current or previous diagnosis were included. 27 healthy individuals without 22q11.2DS was included as a control group with comparable age distribution (controls age range: 12–25 years; mean age: 15.96, standard deviation (SD) = 2.71 years; 22q11.2DS mean age: 15.47, SD 2.41 years, $t_{44} = -0.63$ $p = .53$) and sex ratio (male/female controls: 18/9, cases: 13/6, $\chi^2 = 0.02$, $p = .90$). The following exclusion criteria were applied to controls: presence of a) schizophrenia, schizotypal and delusional disorders (ICD10 DF20–29); b) bipolar disorder (ICD10 DF30–31); c) depression (ICD DF32–33) except for a past episode of mild or moderate depression (ICD10 DF 32.0 or 32.1); d) substance abuse; or e) a first degree relative with a psychotic illness. The regional Ethical Committee of Copenhagen (project id: H-3-2012-136) and the Danish Data protection Agency (project id: 2007-58-0015) approved the study. All participants underwent a verbal and written informed consent process. Participants under the age of 18 provided a verbal assent while their parent's completed written consent.

2.2. Diagnosis and symptoms

The International Classification of Diseases (ICD-10) system was used to evaluate the presence of current psychiatric disorders. Intellectual functioning was assessed using Reynolds Intellectual Screening Test (RIST) (Reynolds and Kamphaus, 2011). We used the Mini International Neuropsychiatric Interview (Sheehan et al., 1998) or the Mini International Neuropsychiatric Interview for Children and Adolescents (Sheehan et al., 2013) to diagnose anxiety, affective disorder and disturbance of activity and attention/attention deficit disorder without hyperactivity. To indicate presence of autism spectrum disorders we used the Social Communication Questionnaire lifetime with a clinical cut-off of 15 (Rutter et al., 2005; Rutter et al., 2003). Screening for current psychosis and rating the severity of schizophrenia-related symptoms was done using the Structured Interview for Prodromal Syndromes (SIPS) (McGlashan et al., 2012; Miller et al., 2003). The schizophrenia-related symptoms were assessed within the four domains: positive (i.e. delusional ideas, persecutory ideas, grandiosity, hallucinations, and disorganized communication), negative (anhedonia or withdrawal, avolition, decreased expression of emotions, decreased experience of emotion or self, impoverished thinking, and deterioration of role functioning), disorganized (odd behavior and appearance, bizarre thinking, trouble with focus and attention, and

personal hygiene), and general symptoms (sleep disturbance, dysphoric mood, motor disturbances, impaired tolerance to normal stress). Two experienced and certified clinicians conducted all clinical interviews.

To test for group differences in SIPS scores we used Wilcoxon rank sum test. A two sample *t*-test was used to test for differences in IQ levels between individuals with 22q11.2DS and controls.

Of the 19 individuals with a 22q11.2 deletion, 1 was diagnosed with affective disorder, 2 with disturbance of activity/attention deficit disorder without hyperactivity, 7 with anxiety or phobia and 1 with both autism spectrum disorder and anxiety or phobia. Only one individual with 22q11.2DS took medication acting on the central nervous system at the time of examination (20 mg retalin). None of the participants had psychosis but the 22q11.2DS group had significantly elevated SIPS scores for all four SIPS symptom domains; negative ($W = 497.5$, $p < .001$), positive ($W = 376$, $p = .004$), disorganized ($W = 416.5$, $p < .001$) and generalized ($W = 324.5$, $p = .037$) symptoms, relative to the control group. The raw sum of negative symptoms in individuals with 22q11.2DS ranged from 1 to 16 (mean = 6.7, $SD = 3.7$), from 0 to 12 for positive symptoms (mean = 2.7, $SD = 3.1$), from 0 to 6 for disorganized symptoms (mean = 1.7, $SD = 1.8$), and from 0 to 7 for generalized symptoms (mean = 0.9, $SD = 1.9$).

The 22q11.2DS group had an IQ (median = 82.0, 90th percentile = 94.4, 10th percentile = 63.8) below the control group (median 108.0, 90th percentile = 127.0, 10th percentile = 95.2, $t_{44} = -7.01$, $p < .001$). See Table 1 for a summary of demographics and clinical scores.

2.3. Stimuli

The roving paradigm was adapted from (Garrido et al., 2008) and comprised of roving sequences of sounds ranging from 1 to 9 drawn from a discrete uniform distribution, Fig. 1A. Each tone in the roving paradigm was a pure sinusoidal tone with frequency 1000 Hz or 1200 Hz and had a duration of 50 ms with a 5 ms rise and fall time. Tones were delivered binaurally via insert-earphones (E-A-RTONE 3A Indianapolis, US), at 85 dB SPL, generated with the Cogent toolbox running in Matlab (http://www.vislab.ucl.ac.uk/cogent_2000.php). With this paradigm it is possible to study the responses to repeated stimulation and thereby the parametric effect of repetition. Participants sat in a comfortable chair and watched a silent movie displaying underwater scenery free of any sudden or salient visual events during the 15 min of recording. Participants were instructed to ignore the sounds.

Table 1

Summary of group data for demographical and clinical data. The content of the table replicates the content of the table in Larsen et al. (2018b).

Measures	Control group	22q11.2 group	Group statistics
Age	Mean 15.96 SD = 2.71	Mean 15.47 SD = 2.41	$t_{44} = -0.63$, $p = .53$
Sex	18 males/9 females	13 males/6 females	$\chi^2 = 0.02$, $p = .90$
IQ	Median = 108.0	Median = 82.0	$t_{44} = -7.01$, $p < .001$
	90th percentile = 127.0	90th percentile = 94.4	
	10th percentile = 95.2	10th percentile = 63.8	
	Mean 109.0 SD = 12.5	Mean 77.7 SD = 16.06	
SIPS – subscales			
Negative	Mean 0.59 SD = 1.04 Range 0–4	Mean 6.68 SD = 3.67 Range 1–16	$W = 477$, $p \leq .001$
Positive	Mean 0.81 SD = 1.49 Range 0–6	Mean 2.74 SD = 3.07 Range 0–12	$W = 305.5$, $p = .008$
Disorganized	Mean 0.11 SD = 0.42 Range 0–2	Mean 1.68 SD = 1.83 Range 0–6	$W = 404$, $p < .001$
Generalized	Mean 0.15 SD = 0.46 Range 0–2	Mean 0.95 SD = 1.90 Range 0–7	$W = 312.5$, $p = .027$
Diagnosis			
MD	$N = 0$	$N = 1$	
ADHD/ADD	$N = 0$	$N = 2$	
Anxiety or phobia	$N = 0$	$N = 8$	
ASD	$N = 0$	$N = 1$	

Audiometric testing was performed prior to the experiment, (20 dB random test Oscilla USB-310 Tablet screening audiometer, Aarhus, Denmark). At 1000 Hz the observed threshold levels were (mean = 20.1, $SD = 0.5$) for controls and (mean = 23.4, $SD = 4.0$) for 22q11.2DS.

2.4. Data acquisition and pre-processing

EEG data were recorded using a 128 channel ActiveTwo Biosemi System (BioSemi, Amsterdam, Netherlands), with a sampling frequency of 4096 Hz. Pre-processing included; band pass filtering between 0.5 Hz - 40 Hz using a second order Butterworth filter, downsampling to 500 Hz and finally epoching with a peristimulus window of -100 ms to 400 ms. The preprocessing was carried out using EEGLAB (Delorme and Makeig, 2004). Baseline correction was applied using the average over the time window -100 ms to -10 ms. Re-referencing to the average reference, artefact removal, scalp analysis, and the DCM analysis were performed using SPM12 (<http://www.fil.ion.ucl.ac.uk/spm/>). Epochs were rejected if their values exceeded $\pm 100 \mu V$. One of the participants (belonging to the 22q11.2DS group) was discarded because the majority of epochs were rejected with this approach (above 80%).

The three models accounting for repetition dependent effects

Since we were interested in the repetition dependent changes in ERPs and effective connectivity, we explored three different models, given below for tone $r = 1, \dots, 9$.

Adaptation model: $x_1(r) = \exp(-r)$

Prediction model: $x_2(r) = \exp(r)$ (1)

Combined adaptation&prediction model: $x_3(r) = x_1(r) + x_2(r)$

We chose the exponential function, given that responses are typically seen heavily reduced in the first repetition, whereas responses seem to become similar thereafter, which is in line with recent findings (Stefanics et al., 2018), see Fig. 1B. The adaptation model postulates that responses decrease with the number of repetitions. Conversely, the prediction model postulates that responses will increase with repetitions, reflecting formation of an expectation that a new event will occur. Finally the adaptation&prediction model is a combination of the adaptation and prediction model in that the initial exponential decay will capture changes due to habituation or adaptation and the growing exponential towards the end will capture formation of an expectation, or prediction.




Fig. 1. Experimental design of the roving paradigm and the three different repetition effects models. A: The tone repetition, R_N , varies randomly between 0 and 8 (maximum of 9 tones). The sequences of tones vary by having a frequency of either 1000 Hz or 1200 Hz. Stimulus onset asynchrony is fixed at 500 ms. B: The three parametric models for repetition-specific effects: the adaptation model, the prediction model and the adaptation&prediction model. C: Model space for DCM models. Each family consisted of the same DCMs, but deviates in the parametric modulation between conditions, that is, the effect that repetitions of tones has on ERPs.

2.5. Posterior probability maps

To compare the adaptation, prediction, and adaptation&prediction models for ERPs, we used posterior probability maps (Garrido et al., 2018; Harris et al., 2018; Rosa et al., 2010). Epoch data were converted into scalp-map images of dimension 64×64 obtained using interpolation. After the conversion to scalp-map images smoothing using a Gaussian kernel specified by a FWHM of 8 mm^2 in the spatial dimension and 10 ms in the temporal dimension was performed.

Individual participant voxel-wise whole brain log-evidences were

calculated using regressors describing the hypothesized relationship amongst the tone repetitions i.e. the equations in (1). The log-evidence for each model were estimated using the variational Bayes first-level model specification (Penny et al., 2005). Group level (pooled across controls and 22q11.2DS) posterior probability maps were calculated using the random effects approach (RFX) for each model. These probability maps can then be used to compare between the three different models for each voxel and time point.




Fig. 2. Grand average difference responses for controls and 22q11.2DS from channel Fz. Left: responses to each tone repetition for controls. Right: Corresponding responses for individuals with 22q11.2DS. First row represent the mean of the responses whereas the second row represents the mean with the shaded area representing one standard deviation from the mean.

2.6. Dynamic causal model specification

To investigate the underlying connectivity network of RS, we used DCM which is a hypothesis driven method for estimating effective connectivity between brain areas (David et al., 2006; Friston et al., 2003). We have previously used the same paradigm to study MMN responses in 22q11.2DS (Larsen et al., 2018b) where we formulated a set of models motivated by previous studies on MMN generators (Doeller et al., 2003; Grau et al., 2007; Opitz et al., 2002; Rinne et al., 2000) as well as previous model comparisons of MMN generation (Boly et al., 2011; Garrido et al., 2008). This network includes bilateral sources in the primary auditory cortex (A1), superior temporal gyrus (STG) and inferior frontal gyrus (IFG), with the IFG usually being most consistent in the right side. The bilateral sources in A1 receive the input. RS in a roving MMN paradigm has been previously studied using DCM (Garrido et al., 2009) with bilateral A1 and STG sources being included. We defined 16 models starting with the right and left A1 and building up the remaining models by adding hierarchical levels until we had a full network comprising the six sources: bilateral A1, STG and IFG, Fig. 1B. Each of the parametric DCM models were estimated for each participant individually with all nine tones in the same model. We defined each of these parametric forms as families that only deviate in the specific condition-specific parametric effect.

2.7. Bayesian model selection

RFX Bayesian model selection was used for the pooled group to test which of the repetition dependent models best described the data overall both at the scalp level (posterior probability maps) and connectivity (DCM). This model comparison is based on the free energy

that accounts for both model evidence and model complexity (Rosa et al., 2010). Posterior and exceedance probabilities were used to compare the models.

2.8. Assessing group differences at the scalp level and connectivity level

2.8.1. Spatiotemporal analysis

In order to test for group differences within the winning model at the scalp level, spatio-temporal analysis was performed over the whole sensor-space (i.e. all electrodes) and time (0 ms to 400 ms) using a full factorial 2×9 design with factors group (controls and 22q11.2DS) and condition (repetitions) and age and sex included as covariates. Weights under the winning model, given by Eq. (1) were entered as contrast allowing to assess group differences in the parametric effect present i.e., how much of the winning model is present in controls compared to 22q11.2DS. All p values reported are thresholded using $p < .05$ FWE corrected at cluster level. To enable the investigation as to whether scalp data activity was correlated with the clinical symptoms in the 22q11.2DS group, we extracted activity from single-participant contrast images associated with the parametric effect. The activity was extracted from a square region with a size of $10 \text{ mm} \times 10 \text{ mm}$ around the peak difference between controls and patients with respect to the winning model at the scalp level. Correlation with negative symptoms were performed using Pearson correlation. We did not pursue correlations with positive, disorganized and generalized symptoms as this in our opinion would not be meaningful. All individuals with a 22q11.2 deletion scored low and below clinically significant thresholds. Furthermore, the variation in symptoms scores was low and extremely right (positive) skewed with five (38%) of the individuals with a 22q11.2 deletion having a score of zero.

2.8.2. Connectivity parameters

Bayesian model averaging (BMA) was carried out within the family with highest exceedance probability, to allow for group comparison of the connectivity parameters (B-parameters) (Penny et al., 2010). A one-way ANCOVA with group as a factor (controls and 22q11.2DS) and age and sex as covariates was performed for each of the parameters. Results are reported both uncorrected as well as corrected for multiple comparison using Bonferroni.

3. Results

Repetition dependent changes can be explained by a combination of adaptation and prediction both in cortical responses and in effective connectivity

We have previously confirmed the presence of repetition dependent responses in this paradigm in the form of a detectable MMN response (Larsen et al., 2018b). Fig. 2 shows difference waves for each repetition of the tones for controls and individuals with 22q11.2DS. Visually it can be seen that controls (left column) show a rapid decay from first to second repetition (in blue) compared to the rest of the repetitions whereas there seems to be no clear pattern between responses to the tone repetitions in the 22q11.2DS group (right column).

At the scalp level the posterior probability maps showed that the combined adaptation&prediction model outperformed the adaptation and prediction models throughout all time points, see Fig. 3A and B, where probabilities are shown summed across space and pooled across the two groups (controls and 22q11.2DS). However, the model's out-performance was most pronounced in the middle of the epoch, from 50 ms to 350 ms. Summing across both time and space, (Fig. 3C), the

model probability for the combined model outperforms the two other models. It is observed that the model probabilities in the baseline period (Fig. 3E) is close to equal meaning that the bias towards a winning model is very small. The exceedance probability of the DCMs for the combined model is 1 (Fig. 3D), meaning that at the connectivity level, this model is a clear winner as well. Together, these results show that repetition-dependent changes in ERPs and in effective connectivity are best explained by adaptation and prediction formation. The spatial distribution of the probabilities in Fig. 3F, thresholded at posterior probability $p = .83$, shows that the spatial distribution of the combined adaptation&prediction model involves electrodes throughout the fronto-central area.

Within the combined adaptation&prediction model the DCM model selection of the models shown in Fig. 1C, did not reveal a clear winning model, which is why we kept the model comparison at the family level.

3.1. 22q11.2DS show reduced adaptation but not prediction

Knowing that repetition dependent changes are explained by a combination of adaptation and prediction processes manifested both in the ERPs and in condition-dependent connectivity changes, we next tested for group differences within this model. Analysis of the scalp maps of the repeated stimuli for the contrast controls greater than 22q11.2DS for the combined adaptation&prediction model revealed an effect peaking at 92 ms, see Fig. 4A. Further, we observed an effect around 172 ms whereas the reversed contrast, 22q11.2DS greater than controls revealed an effect at 74 ms. However, the effect at 74 ms was a small set of points that fell in the interpolated area where no electrodes were positioned. We therefore see this effect as spurious and have not




Fig. 3. Exceedance and posterior probabilities of the three models. A: posterior probability as a function of time, summed across space for the adaptation model (turquoise), the prediction model (yellow) and the combined adaptation&prediction model (blue). B: Same as A, for exceedance probabilities. C: Posterior and exceedance probability for the model comparison at the scalp level, summed across space and time. D: Posterior and exceedance probability for DCMs with connectivity modulations according to the adaptation, prediction, and the adaptation&prediction family. E: Same as C, for the baseline period only. F: Spatial distribution of the combined adaptation&prediction model thresholded at posterior probability $p = .83$. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)




Fig. 4. T-maps show that attenuated adaptation processes drive reduced repetition suppression in individuals with 22q11.2DS. A: Group effect for the winning adaptation&prediction model. There is a significant cluster in the fronto-central area peaking at 92 ms. B: To determine the driver of this effect, we also show the map of the main effect of group for the adaptation model (no effect observed for the prediction model). All results are shown at $p < .05$ FWE corrected at cluster level. C and D: Correlation with activity associated with both the adaptation&prediction model (C) and adaptation model (D) with negative symptoms. The higher degree of negative symptoms in 22q11.2DS were associated with less amount of activation associated with both the adaptation&prediction and adaptation model. p -values shown are corrected for multiple comparison using Bonferroni.

depicted in Fig. 4. To delineate if the group difference at 92 ms was driven by the adaptation or the prediction processes, we looked at the group effects for the adaptation model and prediction model Fig. 4B. Our results indicate that the group difference was driven by the adaptation process, suggesting that the ability to adapt to the tones is impaired in individuals with 22q11.2DS.

We then investigated whether the degree to which repetition dependent changes in cortical responses in 22q11.2DS were associated with symptomatology. To do so, we restricted our search to a square region round the peak difference in the controls greater than 22q11.2 contrast for both the adaptation&prediction model and the adaptation model, where group differences were observed. The strength of the parametric modulator extracted from this region for each individual with 22q11.2DS was negatively correlated with the severity of negative symptoms both for the adaptation&prediction model (Fig. 4C) ($\rho = -0.578$, $p = .012$, unc., $p = .024$ corrected) and the adaptation model (Fig. 4D) ($\rho = -0.662$, $p = .003$, unc., $p = .006$ corrected). Hence, the more severe the negative symptoms, the less activity in the areas associated with the adaptation&prediction and the adaptation model, where the strongest correlation was seen for the adaptation model.

Since 22q11.2DS is associated with hearing loss (Jiramongkolchai et al., 2016) and lower IQ levels (Gothelf et al., 2007; Vorstman et al., 2015) we did a post-hoc analysis to test whether these variables could explain the observed effects in the spatio-temporal analysis. IQ showed

a positive effect at fronto-temporal channels at around 80 ms and 160 ms whereas a positive effect of hearing levels were found at central electrodes around 156 ms and 358 ms as well as at 106 ms over right temporal channels, see Fig. 1 in supplementary material. All effects of IQ and hearing were observed in areas and time points different from the observed group effects. Group effects persisted even after adding hearing levels and IQ as a covariate in the analysis, suggesting that even though these showed effects on the EEG, they do not fully account for the observed group effects.

3.2. Group differences in connectivity strength

Individuals with a 22q11.2 deletion showed a stronger modulation than controls in the extrinsic connection from right IFG to right STG ($F_{1,41} = 6.147$, $p = .017$) in the B parameter associated with an adaptation effect. However, it did not survive correction for multiple comparisons using a conservative Bonferroni correction for 12 test, i.e. connections ($\alpha = 0.05/12 = 0.004$). There was no group difference observed in the B parameters associated with the prediction effect. There was no effect of the covariates sex and age. The group effect persisted when adding hearing levels as covariate, but disappeared when adding IQ as a covariate, suggesting that IQ was driving this effect.

4. Discussion

This study provides evidence that adaptation to repeated sounds is diminished in a group of young non psychotic individuals with 22q11.2DS. Our results suggest that repetition-dependent changes both in ERPs and in effective connectivity are modulated by a combination of adaptation and prediction processes. Furthermore, we found that group differences in the relationship between ERP activity and stimuli repetition was driven by reduced adaptation in individuals with 22q11.2DS in the early ERP component N1, at fronto-central electrodes. Critically, the degree to which this relationship between ERP activity over repetitions was present, correlated negatively with the degree of negative symptoms in 22q11.2DS. Results therefore suggest that repetition dependent changes in cortical responses in 22q11.2DS are associated with negative symptoms.

RS is characterized by a reduction in neural activity, or adaptation, caused by repeated stimuli (Buckner et al., 1998; Grill-Spector et al., 1999; Henson et al., 2000; Kourtzi and Kanwisher, 2001), a phenomenon thought to be mediated by synaptic communication. Predictive coding theories have re-interpreted RS as the neural mechanism underpinning perceptual learning and inference (Aukstulewicz and Friston, 2016; Friston, 2005). Our findings suggest that repetition-dependent changes both in ERPs and in brain connectivity can be explained by a model combining both adaptation and prediction processes. The adaptation component of our combined model predicts an initial exuberant prediction error occurring immediately after a change in sound statistics, which is then followed by decreases in neuronal responses caused by sound repetition. While this accounts for the neurophysiological data evoked by the first half of the sound trains, the prediction component resembles an expectation build-up, as if the participant began to expect an eventual change sometime during the second half of the sound trains. This is in line with the notion that responses to repeated sounds are not only caused by simple mechanisms as neural fatigue, but are likely to be caused by fulfilled expectations (Aukstulewicz and Friston, 2016; Summerfield et al., 2008; Todorovic et al., 2011) including forward message passing of prediction error and backward message passing of predictions or expectations. The spatio-temporal analysis within the combined adaptation&prediction model revealed a group difference driven by the adaptation component, whereas no difference was seen in the prediction component. This is in line with our previous work, where no difference was seen in MMN responses between the two groups (indicating that prediction is preserved) (Larsen et al., 2018b). There is however opposing results in the literature on the change detection mechanism in 22q11.2DS. While MMN evoked by a duration deviant has been shown to be reduced in 22q11.2DS (Baker et al., 2005) using a classical oddball paradigm, no difference across five deviants types; duration, frequency, gap, intensity and location was observed in (Zarchi et al., 2013). While MMN was found to be preserved in (Larsen et al., 2018b), we found a general increased response to tones, evidenced by increased N1 responses in 22q11.2, suggesting either increased sensitivity to tones, or reduced adaptation. Here, we show that the adaptation component is reduced in 22q11.2DS.

We found a correlation with the ERP activity over repetitions and negative symptoms in the 22q11.2DS group. Specifically, reduced adaptation to repetitive sounds was associated with greater negative symptoms in 22q11.2DS, suggesting that adaptation to repeated sounds might play a role in the generation of negative symptoms in 22q11.2DS. Predictive coding ideas have been previously discussed in the context of psychosis and are reminiscent of Kapur's model of psychosis (Kapur, 2003). The relation between the adaptive processes and the negative symptoms in individuals with 22q11.2DS reported here do not speak directly to the emergence of positive symptoms in Kapur's model. However, negative symptoms may be related to positive symptoms (i.e. if beliefs are repeatedly wrong, why would we act on them (Corlett, 2015)). Hence, reduced adaptation to repeated stimuli might be

indirectly related to positive symptoms. However, we were unable to test such a relationship, since positive symptoms were very sparse in this group and none of the individuals met the threshold criteria for clinical relevance. Therefore, this is purely speculative.

RS is very sparsely studied in 22q11.2DS with opposite results on sensory gating as well, with P50 shown to be sometimes intact (Rihs et al., 2013; Vorstman et al., 2009) and other times impaired (Zarchi et al., 2013). However, sensory gating usually entails a paired-click paradigm, excluding the possibility of studying effects beyond the first repetition which have been shown to occur (Stefanics et al., 2018). It is therefore hard to compare results from the present study to previous findings on P50 sensory gating. The small sample size of nonpsychotic individuals with 22q11.2DS is a limitation of the study. The abnormalities in adaptive processes found here warrant replication in larger cohorts. However, it should be noted that all results presented in the paper, have been corrected for multiple comparison, which limits the possibility for false positive results.

Here, we have focused on repetition suppression in 22q11.2DS with a specific emphasis on the susceptibility to psychosis, given the significant higher risk for psychosis associated with the deletion. It is, however, important to note that the microdeletion involves multiple genes and therefore it is not only associated with a high risk of psychosis but with a broader range of psychiatric and neurodevelopmental disorders (Bassett et al., 2008; Olsen et al., 2018; Purcell et al., 2009; Schneider et al., 2014). In conclusion, we show that young non-psychotic individuals with 22q11.2DS are impaired at modulating neural activity to the environmental statistics associated with repeated stimuli.

Financial disclosures

H.R.S. received honoraria as speaker from Lundbeck A/S, Valby, Denmark, Biogen Idec, Denmark A/S, Genzyme, Denmark and MerckSerono, Denmark, honoraria as editor from Elsevier Publishers, Amsterdam, The Netherlands and Springer Publishing, Stuttgart, Germany, travel support from MagVenture, Denmark, and grant support from Biogen Idec, Denmark A/S. M.R.B. is a prior employee at H. Lundbeck A/S, Denmark and received financial support for her PhD from the Innovation Fund Denmark. M.D. is employed with Lundbeck A/S. The authors declare no further biomedical financial interests or potential conflicts of interest.

This article is distributed under the Danish legislation governed by the Privacy Act (act# 429, 31/05/2000), which does not permit data sharing at publicly available repositories or in raw formats. Summary statistics can be obtained through contact with the corresponding author (melissal@drcomr.dk).

Acknowledgements

We thank all participants and families for taking their time to participate in our study and the Danish National 22q11DS Association for their strong support of our work. We highly appreciate the efforts of Anders Vangkilde and Henriette Schmock for their dedicated assistance in the recruitment and clinical assessments. We thank staff involved in the Danish Blood Donor Study, Capital Region Blood bank, Glostrup and <http://www.forsogsperson.dk/> from which our control participants were recruited.

This study was funded by the Lundbeck Foundation, Denmark (R155-2014-1724); Lundbeck Foundation [Grant of Excellence "ContAct" R59 A5399]; Lundbeck Foundation fellowship (R105-9813); The Capital Region's Research Foundation for Mental Health Research; the Australian Research Council Centre of Excellence for Integrative Brain Function (ARC Centre Grant 140100007CE); University of Queensland Fellowship (2016000071) to MIG. H.R.S. holds a professorship in precision medicine at the Faculty of Health Sciences and Medicine, University of Copenhagen, sponsored by the Lundbeckfonden.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.nicl.2019.101721>.

References

- Adler, L.E., Pachtman, E., Franks, R.D., Pecevic, M., Waldo, M.C., Freedman, R., 1982. Neurophysiological evidence for a defect in neuronal mechanisms involved in sensory gating in schizophrenia. *Biol. Psychiatry* 17 (6), 639–654.
- Andrade, G.N., Butler, J.S., Mercier, M.R., Molholm, S., Foxe, J.J., 2015. Spatio-temporal dynamics of adaptation in the human visual system: a high-density electrical mapping study. *Eur. J. Neurosci.* 41 (7), 925–939. <https://doi.org/10.1111/ejn.12849>.
- Auksztulewicz, R., Friston, K., 2016. Repetition suppression and its contextual determinants in predictive coding. *Cortex* 80, 125–140. <https://doi.org/10.1016/j.cortex.2015.11.024>.
- Baker, K., Baldeweg, T., Sivagnanasundaram, S., Scambler, P., Skuse, D., 2005. COMT Val108/158Met modifies mismatch negativity and cognitive function in 22q11 deletion syndrome. *Biol. Psychiatry* 58 (1), 23–31. <https://doi.org/10.1016/j.biopsych.2005.03.020>.
- Baldeweg, T., 2007. ERP repetition effects and mismatch negativity generation: a predictive coding perspective. *J. Psychophysiol.* 21 (3–4), 204–213. <https://doi.org/10.1027/0269-8803.21.34.204>.
- Baldeweg, T., Klugman, A., Gruzelier, J., Hirsch, S.R., 2004. Mismatch negativity potentials and cognitive impairment in schizophrenia. *Schizophr. Res.* 69 (2–3), 203–217. <https://doi.org/10.1016/j.schres.2003.09.009>.
- Bassett, A.S., Marshall, C.R., Lionel, A.C., Chow, E.W.C., Scherer, S.W., 2008. Copy number variations and risk for schizophrenia in 22q11.2 deletion syndrome. *Hum. Mol. Genet.* 17 (24), 4045–4053. <https://doi.org/10.1093/hmg/ddn307>.
- Boly, M., Garrido, M.I., Gosses, O., Bruno, M., Boveroux, P., Schnakers, C., ... Friston, K., 2011. Preserved feedforward but impaired top-down processes in the vegetative state. *Science* 332 (6031), 858–862. <https://doi.org/10.1126/science.1202043>.
- Buckner, R.L., Goodman, J., Burock, M., Rotte, M., Koutstaal, W., Schacter, D., ... Dale, A.M., 1998. Functional-anatomic correlates of object priming in humans revealed by rapid presentation event-related fMRI. *Neuron* 20 (2), 285–296.
- Coffman, B.A., Haigh, S.M., Murphy, T.K., Salisbury, D.F., 2017. Impairment in mismatch negativity but not repetition suppression in schizophrenia. *Brain Topogr.* 30 (4), 521–530. <https://doi.org/10.1007/s10548-017-0571-1>.
- Corlett, P.R., 2015. Answering some phenomenal challenges to the prediction error model of delusions. *World Psychiatry: Off. J. World Psychiatric Assoc. (WPA)* 14 (2), 181–183. <https://doi.org/10.1002/wps.20211>.
- David, O., Kiebel, S.J., Harrison, L.M., Mattout, J., Kilner, J.M., Friston, K.J., 2006. Dynamic causal modeling of evoked responses in EEG and MEG. *NeuroImage* 30 (4), 1255–1272. <https://doi.org/10.1016/j.neuroimage.2005.10.045>.
- Delorme, A., Makeig, S., 2004 Mar 15. EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *J. Neurosci. Methods* 134 (1), 9–21.
- Doeller, C.F.C., Opitz, B., Mecklinger, A., Krick, C., Reith, W., Schröger, E., 2003. Prefrontal cortex involvement in preattentive auditory deviance detection: neuroimaging and electrophysiological evidence. *NeuroImage* 20 (2), 1270–1282. [https://doi.org/10.1016/S1053-8119\(03\)00389-6](https://doi.org/10.1016/S1053-8119(03)00389-6).
- Friston, K., 2005. A theory of cortical responses. *Philos. Trans. R. Soc. Lond. Ser. B Biol. Sci.* 360 (1456), 815–836. <https://doi.org/10.1098/rstb.2005.1622>.
- Friston, K.J., Harrison, L., Penny, W., 2003. Dynamic causal modelling. *NeuroImage* 19 (4), 1273–1302. [https://doi.org/10.1016/S1053-8119\(03\)00202-7](https://doi.org/10.1016/S1053-8119(03)00202-7).
- Garrido, M.I., Friston, K.J., Kiebel, S.J., Stephan, K.E., Baldeweg, T., Kilner, J.M., 2008. The functional anatomy of the MMN: a DCM study of the roving paradigm. *NeuroImage* 42 (2), 936–944. <https://doi.org/10.1016/j.neuroimage.2008.05.018>.
- Garrido, M.I., Kilner, J.M., Kiebel, S.J., Stephan, K.E., Friston, K.J., 2009. Repetition suppression and plasticity in the human brain. *NeuroImage* 48 (1), 269–279. <https://doi.org/10.1016/j.neuroimage.2009.06.034.Repetition>.
- Garrido, M.I., Rowe, E.G., Halász, V., Mattingley, J.B., 2018. Bayesian mapping reveals that attention boosts neural responses to predicted and unpredicted stimuli. *Cereb. Cortex* 28 (5), 1771–1782. <https://doi.org/10.1093/cercor/bhx087>.
- Goodship, J., Cross, I., LiLing, J., Wren, C., 1998. A population study of chromosome 22q11 deletions in infancy. *Arch. Dis. Child.* 79 (4), 348–351.
- Gothelf, D., Feinstein, C., Thompson, T., Gu, E., Penniman, L., Van Stone, E., ... Reiss, A.L., 2007. Risk factors for the emergence of psychotic disorders in adolescents with 22q11.2 deletion syndrome. *Am. J. Psychiatry* 164 (4), 663–669. <https://doi.org/10.1176/appi.ajp.164.4.663>.
- Grau, C., Fuentes-Milla, L., Marco-Pallares, J., 2007. Functional neural dynamics underlying auditory event-related N1 and N1 suppression response. *NeuroImage* 36 (3), 522–531.
- Grill-Spector, K., Kushnir, T., Edelman, S., Avidan, G., Itzhak, Y., Malach, R., 1999. Differential processing of objects under various viewing conditions in the human lateral occipital complex. *Neuron* 24 (1), 187–203. [https://doi.org/10.1016/S0896-6273\(00\)80832-6](https://doi.org/10.1016/S0896-6273(00)80832-6).
- Grill-Spector, K., Henson, R., Martin, A., 2006. Repetition and the brain: neural models of stimulus-specific effects. *Trends Cogn. Sci.* 10 (1), 14–23. <https://doi.org/10.1016/j.tics.2005.11.006>.
- Grother, M., Kovács, G., 2016. Can predictive coding explain repetition suppression? *Cortex* 80, 113–124. <https://doi.org/10.1016/j.cortex.2015.11.027>.
- Haenschel, C., Vernon, D.J., Dwivedi, P., Gruzelier, J.H., Baldeweg, T., 2005. Event-related brain potential correlates of human auditory sensory memory-trace formation. *J. Neurosci.* 25 (45), 10494–10501. <https://doi.org/10.1523/JNEUROSCI.1227-05.2005>.
- Harris, C.D., Rowe, E.G., Randeniya, R., Garrido, M.I., 2018. Bayesian model selection maps for group studies using M/EEG data. *BioRxiv* 365056. <https://doi.org/10.1101/365056>.
- Henson, R., Shallice, T., Dolan, R., 2000. Neuroimaging evidence for dissociable forms of repetition priming. *Science (New York, N.Y.)* 287 (5456), 1269–1272.
- Henson, R., Rylands, A., Ross, E., Vuilleumier, P., Rugg, M., 2004. The effect of repetition lag on electrophysiological and haemodynamic correlates of visual object priming. *NeuroImage* 21 (4), 1674–1689. <https://doi.org/10.1016/j.neuroimage.2003.12.020>.
- Hoeftding, L.K., Trabjerg, B.B., Olsen, L., Mazin, W., Sparsø, T., Vangkilde, A., ... Werge, T., 2017. Risk of psychiatric disorders among individuals with the 22q11.2 deletion or duplication. *JAMA Psychiatry* 74 (3), 282. <https://doi.org/10.1001/jamapsychiatry.2016.3939>.
- Jiramongkolchai, P., Kumar, M.S., Chinnadurai, S., Wootten, C.T., Goudy, S.L., 2016. Prevalence of hearing loss in children with 22q11.2 deletion syndrome. *Int. J. Pediatr. Otorhinolaryngol.* 87, 130–133. <https://doi.org/10.1016/j.ijporl.2016.06.005>.
- Kapur, S., 2003. Psychosis as a state of aberrant salience: a framework linking biology, phenomenology, and pharmacology in schizophrenia. *Am. J. Psychiatr.* 160 (1), 13–23.
- Karayiorgou, M., Simon, T.J., Gogos, J.A., 2010. 22q11.2 microdeletions: linking DNA structural variation to brain dysfunction and schizophrenia. *Nat. Rev. Neurosci.* 11 (6), 402–416. <https://doi.org/10.1038/nrn2841>.
- Kourtzi, Z., Kanwisher, N., 2001. Representation of perceived object shape by the human lateral occipital complex. *Science* 293 (5534), 1506–1509. <https://doi.org/10.1126/science.1061133>.
- Larsen, K.M., Dzaif, I., Siebner, H.R., Garrido, M.I., 2018a. Alteration of functional brain architecture in 22q11.2 deletion syndrome – insights into susceptibility for psychosis. *NeuroImage*. <https://doi.org/10.1016/j.neuroimage.2018.09.001>.
- Larsen, K.M., Mørup, M., Birkenow, M.R., Fischer, E., Hulme, O., Vangkilde, A., ... Garrido, M.I., 2018b. Altered auditory processing and effective connectivity in 22q11.2 deletion syndrome. *Schizophr. Res.* <https://doi.org/10.1016/j.schres.2018.01.026>.
- Mayer, A.R., Hanlon, F.M., Franco, A.R., Teshiba, T.M., Thoma, R.J., Clark, V.P., Canive, J.M., 2009. The neural networks underlying auditory sensory gating. *NeuroImage* 44 (1), 182–189. <https://doi.org/10.1016/j.neuroimage.2008.08.025>.
- McGlashan, T.H., Miller, T.J., Woods, S.W., Rosen, J.L., Hoffmann, R.E., Davidson, L., 2012. Structured Interview for Prodromal Syndromes (SIPS). Norsk version 5.0 Stavanger Universitetssjukehus, Norge.
- Miller, T.J., McGlashan, T.H., Rosen, J.L., Cadenhead, K., Cannon, T., Ventura, J., ... Woods, S.W., 2003. Prodromal assessment with the structured interview for prodromal syndromes and the scale of prodromal symptoms: predictive validity, inter-rater reliability, and training to reliability. *Schizophr. Bull.* 29 (4), 703–715.
- Olsen, L., Sparsø, T., Weinsheimer, S.M., Dos Santos, M.B.Q., Mazin, W., Rosengren, A., ... Werge, T., 2018. Prevalence of rearrangements in the 22q11.2 region and population-based risk of neuropsychiatric and developmental disorders in a Danish population: a case-cohort study. *Lancet Psychiatry* 5 (7), 573–580. [https://doi.org/10.1016/S2215-0366\(18\)30168-8](https://doi.org/10.1016/S2215-0366(18)30168-8).
- Opitz, B., Rinne, T., Mecklinger, A., von Cramon, D.Y., Schröger, E., 2002. Differential contribution of frontal and temporal cortices to auditory change detection: fMRI and ERP results. *NeuroImage* 15 (1), 167–174. <https://doi.org/10.1006/nimg.2001.0970>.
- Oskarsdóttir, S., Vujic, M., Fasth, A., 2004. Incidence and prevalence of the 22q11 deletion syndrome: a population-based study in Western Sweden. *Arch. Dis. Child.* 89 (2), 148–151.
- Penny, W.D., Trujillo-Barreto, N.J., Friston, K.J., 2005. Bayesian fMRI time series analysis with spatial priors. *NeuroImage* 24 (2), 350–362. <https://doi.org/10.1016/j.neuroimage.2004.08.034>.
- Penny, W.D., Stephan, K.E., Daunizeau, J., Rosa, M.J., Friston, K.J., Schofield, T.M., Leff, A.P., 2010. Comparing families of dynamic causal models. *PLoS Comput. Biol.* 6 (3), e1000709. <https://doi.org/10.1371/journal.pcbi.1000709>.
- Potter, D., Summerfelt, A., Gold, J., Buchanan, R.W., 2005. Review of clinical correlates of P50 sensory gating abnormalities in patients with schizophrenia. *Schizophr. Bull.* 32 (4), 692–700. <https://doi.org/10.1093/schbul/sbj050>.
- Purcell, S.M., Wray, N.R., Stone, J.L., Visscher, P.M., O'Donovan, M.C., Sullivan, P.F., Sklar, P., 2009. Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature* 460 (7256), 748–752. <https://doi.org/10.1038/nature08185>.
- Rentsch, J., Shen, C., Jockers-Scherübl, M.C., Gallinat, J., Neuhaus, A.H., 2015. Auditory mismatch negativity and repetition suppression deficits in schizophrenia explained by irregular computation of prediction error. *PLoS One* 10 (5), e0126775. <https://doi.org/10.1371/journal.pone.0126775>.
- Reynolds, C.R., Kamphaus, R.W., 2011. Reynolds Intellectual Assessment Scales (RIAS), Reynolds Intellectual Screening Test (RIST). Dansk version - Vejledning Høgrete Psykologisk Forlag, København, Danmark.
- Rihs, T., A. Tomescu, M.I., Britz, J., Rochas, V., Custo, A., Schneider, M., ... Michel, M., 2013. Altered auditory processing in frontal and left temporal cortex in 22q11.2 deletion syndrome: a group at high genetic risk for schizophrenia. *Psychiatry Res.* 212 (2), 141–149. <https://doi.org/10.1016/j.psychres.2012.09.002>.
- Rinne, T., Alho, K., Ilmoniemi, R., Virtanen, J., Näätänen, R., 2000. Separate time behaviors of the temporal and frontal mismatch negativity sources. *NeuroImage* 12 (1), 14–19.
- Robin, N.H., Shprintzen, R.J., 2005. Defining the clinical spectrum of deletion 22q11.2. *J. Pediatr.* 147 (1), 90–96. <https://doi.org/10.1016/j.jpeds.2005.03.007>.
- Rosa, M.J., Bestmann, S., Harrison, L., Penny, W., 2010. Bayesian model selection maps

- for group studies. *NeuroImage* 49 (1), 217–224. <https://doi.org/10.1016/j.neuroimage.2009.08.051>.
- Rosburg, T., 2018. Auditory N100 gating in patients with schizophrenia: a systematic meta-analysis. *Clin. Neurophysiol.* 129 (10), 2099–2111. <https://doi.org/10.1016/j.clinph.2018.07.012>.
- Rutter, M., Bailey, A., Lord, C., 2003. *The Social Communication Questionnaire (SCQ) Manual. (Form: Lifetime)*. Western Psychological Services, Los Angeles, CA.
- Rutter, M., Bailey, A., Berument, S.K., Lord, C., Pickles, A., 2005. *The Social Communication Questionnaire (SCQ). Dansk version - Vurdering Livsføreb Hogrefe Psykologisk Forlag, København, Danmark*.
- Schmuck, H., Vangkilde, A., Larsen, K.M., Fischer, E., Birkenow, M.R., Jepsen, J.R.M., ... Olsen, L., 2015. The Danish 22q11 research initiative. *BMC Psychiatry* 15 (1), 220. <https://doi.org/10.1186/s12888-015-0594-7>.
- Schneider, M., Debbané, M., Bassett, A.S., Chow, E.W.C., Fung, W.L.A., van den Bree, M., et al., 2014. Psychiatric disorders from childhood to adulthood in 22q11.2 deletion syndrome: results from the international consortium on brain and behavior in 22q11.2 deletion syndrome. *Am. J. Psychiatry* 171 (6), 627–639. <https://doi.org/10.1176/appi.ajp.2013.13070864>.
- Sheehan, D.V., Lecrubier, Y., Sheehan, K.H., Amorim, P., Janavs, J., Weiller, E., ... Dunbar, G.C., 1998. The mini-international neuropsychiatric interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J. Clin. Psychiatry* 22–33. quiz 34–57. Retrieved from. <http://www.ncbi.nlm.nih.gov/pubmed/9881538>.
- Sheehan, D., Shytle, K., Milo, J.J.Y., 2013. *M.I.N.I. International Neuropsychiatric Interview For Children and Adolescents (M.I.N.I.-Kid). Dansk Version 6.0*.
- Shprintzen, R.J., 2005. Velo-cardio-facial syndrome. In: Allanson, A.J., Cassidy, S.B. (Eds.), *Management of Genetic Syndromes. Vol. 2*. Wiley-Liss, New York, pp. 615–632.
- Stefanics, G., Heinze, J., Czigler, I., Valentini, E., Stephan, K.E., 2018. Timing of repetition suppression of event-related potentials to unattended objects. *Eur. J. Neurosci.* <https://doi.org/10.1111/ejn.13972>.
- Stefansson, H., Rujescu, D., Cichon, S., 2008. Large recurrent microdeletions associated with schizophrenia. *Nature* 455, 232–236.
- Summerfield, C., Trittschuh, E.H., Monti, J.M., Mesulam, M.-M., Egner, T., 2008. Neural repetition suppression reflects fulfilled perceptual expectations. *Nat. Neurosci.* 11 (9), 1004–1006. <https://doi.org/10.1038/nn.2163>.
- Todorovic, A., de Lange, F.P., 2012. Repetition suppression and expectation suppression are dissociable in time in early auditory evoked fields. *J. Neurosci.* 32 (39), 13389–13395. <https://doi.org/10.1523/JNEUROSCI.2227-12.2012>.
- Todorovic, A., van Ede, F., Maris, E., de Lange, F.P., 2011. Prior expectation mediates neural adaptation to repeated sounds in the auditory cortex: an MEG study. *J. Neurosci.* 31 (25), 9118–9123. <https://doi.org/10.1523/JNEUROSCI.1425-11.2011>.
- Vorstman, J.A.S., Turetsky, B.I., Sijmens-Morcus, M.E.J., de Sain, M.G., Dorland, B., Sprong, M., ... Kemner, C., 2009. Proline affects brain function in 22q11DS children with the low activity COMT158 allele. *Neuropsychopharmacology* 34 (3), 739–746. <https://doi.org/10.1038/npp.2008.132>.
- Vorstman, J.A.S., Breetvelt, E.J., Duijff, S.N., Eliez, S., Schneider, M., Jalbrzikowski, M., ... International Consortium on Brain and Behavior in 22q11.2 Deletion Syndrome, I. 22q11. B. B. S., 2015. Cognitive decline preceding the onset of psychosis in patients with 22q11.2 deletion syndrome. *JAMA Psychiatry* 72 (4), 377–385. <https://doi.org/10.1001/jamapsychiatry.2014.2671>.
- Zarchi, O., Carmel, M., Avni, C., Attias, J., Frisch, A., Michaelovsky, E., ... Gothelf, D., 2013. Schizophrenia-like neurophysiological abnormalities in 22q11.2 deletion syndrome and their association to COMT and PRODH genotypes. *J. Psychiatr. Res.* 47 (11), 1623–1629. <https://doi.org/10.1016/j.jpsychires.2013.07.004>.