

## Inflammatory adipocyte-derived extracellular vesicles promote leukocyte attachment to vascular endothelial cells



Rebecca M. Wadey<sup>a</sup>, Katherine D. Connolly<sup>a</sup>, Donna Mathew<sup>a</sup>, Gareth Walters<sup>a</sup>, D. Aled Rees<sup>b</sup>, Philip E. James<sup>a,\*</sup>

<sup>a</sup> Cardiff School of Sport and Health Sciences, Cardiff Metropolitan University, 200 Western Avenue, Cardiff, CF5 2YB, UK

<sup>b</sup> Neuroscience and Mental Health Research Institute, Cardiff University, Hadyn Ellis Building, Maindy Road, Cardiff, CF24 4HQ, UK

### HIGHLIGHTS

- Extracellular vesicles (EVs) from inflamed adipocytes increase VCAM expression in vascular endothelial cells.
- EVs from inflamed and hypoxic adipocytes increase leukocyte attachment to vascular endothelial cells.
- Inflammation and hypoxia affect the expression of adipokines in adipocytes and EVs.
- Inflammation and hypoxia affect adipocyte EV yield and size.

### ARTICLE INFO

#### Keywords:

Adipocyte  
Adipokine  
Obesity  
Endothelial dysfunction  
Vesicles

### ABSTRACT

**Background and aims:** Obesity is associated with an increased risk of cardiovascular disease, but the mechanisms involved are not completely understood. In obesity, the adipocyte microenvironment is characterised by both hypoxia and inflammation. Therefore, we sought to determine whether extracellular vesicles (EVs) derived from adipocytes in this setting might be involved in mediating cardiovascular disease, specifically by promoting leukocyte attachment to vascular endothelial cells.

**Methods:** Mature 3T3-L1 adipocytes were incubated for 24 h under control, TNF- $\alpha$  (30 ng/mL), hypoxia (1% O<sub>2</sub>), or TNF- $\alpha$  + hypoxia (30 ng/mL, 1% O<sub>2</sub>) conditions. EVs were isolated by differential ultracentrifugation and analysed by nanoparticle tracking analysis. Primary human umbilical vein endothelial cells (HUVECs) were treated with EVs for 6 h before being lysed for Western blotting to investigate changes in adhesion molecule production, or for use in leukocyte attachment assays.

**Results:** EVs from adipocytes treated with TNF- $\alpha$  and TNF- $\alpha$  + hypoxia increased vascular cell adhesion molecule (VCAM-1) production in HUVECs compared to basal level ( $4.2 \pm 0.6$  and  $3.8 \pm 0.3$ -fold increase, respectively ( $p < 0.05$ )), an effect that was inhibited by an anti-TNF- $\alpha$  neutralising antibody. Production of other adhesion molecules (E-selectin, P-selectin, platelet endothelial cell adhesion molecule and VE-Cadherin) was unchanged. Pre-incubating HUVECs with TNF- $\alpha$  + hypoxia EVs significantly increased leukocyte attachment compared to basal level ( $3.0 \pm 0.4$ -fold increase ( $p < 0.05$ )).

**Conclusions:** Inflammatory adipocyte EVs induce VCAM-1 production in vascular endothelial cells, accompanied by enhanced leukocyte attachment. Preventing adipocyte derived EV-induced VCAM-1 upregulation may offer a novel therapeutic target in the prevention of obesity-driven cardiovascular disease.

### 1. Introduction

Obesity is a risk factor for the development of cardiovascular disease (CVD) [1] but the complex mechanisms linking the two together are still to be fully elucidated. Obesity combined with metabolic syndrome (an umbrella term encompassing insulin resistance,

dyslipidaemia, vascular endothelial dysfunction, hypertension and a hypercoagulable state), further increases the risk of CVD [2,3]. Whilst a complete understanding of the mechanisms that trigger adipose tissue to become metabolically dysregulated in obesity are not fully understood, evidence suggests that hypoxia and inflammation play a role.

Obese adipose tissue contains localised regions of hypoxia [4–7]

\* Corresponding author.

E-mail address: [pjames@cardiffmet.ac.uk](mailto:pjames@cardiffmet.ac.uk) (P.E. James).

<https://doi.org/10.1016/j.atherosclerosis.2019.01.013>

Received 27 April 2018; Received in revised form 21 November 2018; Accepted 10 January 2019

Available online 24 January 2019

0021-9150/© 2019 Elsevier B.V. All rights reserved.

thought to arise due to increased cell number and cell size beyond the distance oxygen can diffuse from capillaries. Free fatty acids and cytokines produced by hypoxic adipocytes attract and activate macrophages and in doing so, induce local tissue inflammation [8]. Under such conditions, physiological adipokine production is dysregulated. For example, release of pro-inflammatory tumour necrosis factor (TNF)- $\alpha$  is increased [9] whilst anti-inflammatory adiponectin is decreased [10]. The elevation of TNF- $\alpha$  and decrease of adiponectin in plasma are individually associated with increased leukocyte attachment to vascular endothelial cells via increased production of vascular endothelial cell adhesion molecules [11,12]. Subsequent phagocytosis of oxidised low-density lipoproteins within vessel walls results in lipid-laden foam cell formation [13] and later, atherosclerotic plaque development.

Like most cells, adipocytes release extracellular vesicles (EVs) [14,15]. EVs are membrane-bound structures that can convey biological information from a cell of origin to a recipient cell to achieve a target effect. Based on diameter and biogenesis they are categorised into three broad groupings: exosomes (~80–100 nm; released from multivesicular bodies upon plasma membrane fusion), microvesicles (~200–1000 nm; shed directly from the plasma membrane) and apoptotic bodies (~1–5  $\mu$ m; released as blebs during apoptosis). *In vitro* data show that adipocytes release EVs containing adipokines [16–19]. In obesity, adipocytes reside in a hypoxic and inflammatory environment, therefore the content and function of adipocyte-derived EVs in this environment may be different to adipocyte-derived EVs from lean adipose tissue. Adipocyte EVs circulating in obesity may contribute to vascular endothelial dysfunction, which later leads to atherosclerotic plaque formation. We sought to test this hypothesis *in vitro* using a leukocyte attachment assay with vascular endothelial cells isolated from umbilical cords and EVs derived from adipocytes cultured in conditions representative of lean and obese (hypoxic and inflammatory) adipose tissue.

## 2. Materials and methods

Ethical approval for this study was granted by the NHS Health Research Authority (ethics committee reference: 14/NW/1459), and Cardiff Metropolitan University's Research Ethics Committee.

### 2.1. Adipocyte culture and EV isolation

3T3-L1 adipocytes were cultured for 14 days as described by Connolly *et al*<sup>16</sup>. For all EV-isolation experiments, serum-containing culture medium was replaced with serum-free medium for 24 h. During this time, adipocytes were exposed to one of four treatments representative of control (95% air/5% CO<sub>2</sub>), inflamed (30 ng/mL TNF- $\alpha$  (13473019, ThermoFisher, USA)), hypoxic (1% O<sub>2</sub>), and inflamed and hypoxic (30 ng/mL TNF- $\alpha$ , 1% O<sub>2</sub>) adipose tissue. Cell media were centrifuged at 4 °C at 1000 g (5 min), 15,000 g (15 min), and finally 100,000 g (1 h) to pellet EVs. EVs were resuspended in 1  $\times$  PBS and quantified by nanoparticle tracking analysis (NTA; a technique that tracks nanoparticles in suspension using a laser, and based on Brownian motion, determines particle size and concentration). NTA was performed using a NanoSight LM10 with a 642 nm laser (Malvern Instruments Ltd, Malvern, UK), using software version 2.3, screen gain 4 and camera level 10. Five, 1-min videos were recorded per sample with analysis screen gain 10 and detection threshold 4. Temperature ranged from 20 to 23.5 °C.

### 2.2. HUVEC isolation and culture

Human umbilical cords were obtained following elective Caesarean sections. Saline was used to flush blood out of the umbilical vein, before one end was clamped. Collagenase type 1A (1 mg/mL; C5894, Sigma) in Medium 199 (M199; 31150, ThermoFisher) at 37 °C, was syringed into the vein until the cord became taut and then the end of the cord was

also clamped. After 15 min, one clamp was released allowing the vascular endothelial cell suspension to be collected into a Falcon tube. Copious growth medium (M199 containing 10% (v/v) foetal bovine serum (10500-064, ThermoFisher), human epidermal growth factor (1 ng/mL; 13453029, ThermoFisher), hydrocortisone (1  $\mu$ g/mL; H0888, Sigma), gentamycin (35  $\mu$ g/mL; G1272, Sigma) and amphotericin (0.5  $\mu$ g/mL; A2942, Sigma) was added to terminate the enzymatic digestion. Cells were centrifuged (300 g, 5 min, 4 °C), resuspended in growth medium, and plated in 96-well plates for leukocyte attachment assays or 6-well plates for protein assays. Plates were pre-coated with 1% (w/v) bovine skin gelatin (G9391, Sigma) in 1  $\times$  PBS. HUVECs were given 2 h to adhere, before the medium was aspirated and replaced. HUVECs reached confluency within 4–7 days, were never passaged, and were used for experiments within 7 days.

### 2.3. Leukocyte isolation

Blood (10 mL) was obtained from healthy volunteers by venepuncture and transferred into a universal container (UC) containing Heparin (100  $\mu$ L, 5000 I.U./mL, Wockhardt, India). Dextran (2.5 mL, 6% (w/v), Sigma) dissolved in 1  $\times$  balanced salt solution (BSS; 0.13 M NaCl, 2.6 mM KCl, 8.0 mM Na<sub>2</sub>HPO<sub>4</sub>, 1.83 mM KH<sub>2</sub>PO<sub>4</sub>, pH 7.4) was added and mixed by a single inversion. Blood was transferred into a fresh UC and allowed to fractionate for 1 h. The buffy coat layer (~1.5 mL) was transferred into a fresh UC. Cells were collected by centrifugation (300 g, 2 min, room temperature) and the pellet resuspended in sterile H<sub>2</sub>O to burst any contaminating erythrocyte membranes. After 10 s the UC was filled with BSS and the leukocytes were pelleted as above. The pellet was resuspended in Krebs-BSA (0.1% (w/v) bovine serum albumin (BSA) in 1  $\times$  Krebs (1.2 M NaCl, 0.48 M KCl, 0.12 M KH<sub>2</sub>PO<sub>4</sub>, 0.12 M MgSO<sub>4</sub>, 0.13 M CaCl<sub>2</sub>, 2.5 M HEPES, pH 7.4)) that had been passed through a 0.22  $\mu$ m pore sterile-filter (Merck Millipore, USA) for sterilisation and remove possible contaminating serum extracellular vesicles. Cells were incubated on ice, in darkness, with CellTrace™ (1:1000; C34851, Invitrogen). After 10 min, leukocytes were centrifuged, resuspended in Krebs-BSA and left to settle on ice for 30 min. Prior to the attachment assay, the leukocyte suspension was diluted 1:10 with Krebs-BSA pre-warmed to 37 °C.

### 2.4. Leukocyte attachment assay

Serum-free M199 (150  $\mu$ L) containing  $1.2 \pm 0.4 \times 10^{10}$  (Control),  $2.5 \pm 0.4 \times 10^{10}$  (TNF- $\alpha$ ),  $8.6 \pm 0.8 \times 10^{10}$  (Hypoxia) and  $1.2 \pm 0.3 \times 10^{11}$  (TNF- $\alpha$  & Hypoxia) adipocyte EV/mL (mean  $\pm$  SEM; N = 3, n = 9), was added to individual wells of a 96-well plate containing a confluent monolayer of HUVECs. Wells were also used to determine basal leukocyte attachment (no EVs; negative control) or 100 ng/mL TNF- $\alpha$  to increase HUVEC VCAM-1 production (no EVs; positive control). After 6 h, EVs were removed with three Krebs washes before 150  $\mu$ L of the fluorescently-labelled leukocyte suspension was added to all wells. Plates were incubated for 30 min before non-adherent cells were removed with three Krebs washes. Three images from around the centre of each well were captured using an inverted fluorescence microscope. The percentage of total image area covered by leukocytes was determined using Image J software (1.49v; National Institutes of Health, USA). Incubation timings were based on published protocols [20]. Three separate sets of adipocyte-derived EVs were used in this experiment and each set of EVs was tested on three HUVEC samples (N = 3, n = 9).

### 2.5. Effect of adipocyte EVs on HUVEC adhesion molecule production

Serum-free M199 (1.5 mL) containing  $1.2 \pm 0.4 \times 10^{10}$  (Control),  $2.5 \pm 0.4 \times 10^{10}$  (TNF- $\alpha$ ),  $8.6 \pm 0.8 \times 10^{10}$  (Hypoxia) and  $1.2 \pm 0.3 \times 10^{11}$  (TNF- $\alpha$  & Hypoxia) adipocyte EV/mL (mean  $\pm$  SEM, N = 3, n = 3), was added to individual wells of a 6-well

plate of HUVECs for 6 h. Another well was used for the determination of basal protein production (no EVs; negative control). EVs were removed with three  $1 \times$  PBS washes before HUVECs were lysed using radioimmunoprecipitation assay buffer (Invitrogen) containing protease inhibitors (Roche, Switzerland). Lysates were centrifuged (12,000 g, 10 min,  $4^\circ\text{C}$ ) and supernatants frozen until required. The protein concentration of each sample was determined using a NanoDrop spectrophotometer and samples analysed by Western blotting. Lysates were mixed with sample buffer and reducing agent (Invitrogen) and heated ( $95^\circ\text{C}$ , 10 min). Ten  $\mu\text{g}$  of total protein from each sample was loaded onto 4–12% Bis-Tris gels (Novex™, ThermoFisher) and SDS-PAGE performed (180 V, 1 h). Resolved proteins were transferred to polyvinylidene difluoride membranes (40 V, 75 min). Membranes were blocked using 5% (w/v) non-fat dried milk in  $1 \times$  Tris-buffered saline containing 0.05% Tween-20 (TBST) (1 h). Primary antibodies directed at proteins of interest (vascular cell adhesion molecule (VCAM-1; ab134047), E-selectin (ab18981), P-selectin (ab59738), platelet endothelial cell adhesion molecule (PECAM; ab28364), vascular endothelial (VE)-cadherin (ab33168), endothelial nitric oxide synthase (eNOS; ab76198) (all Abcam, Cambridge, UK)) were diluted 1:500 in blocking buffer and incubated with membranes overnight. Membranes were then incubated with horseradish peroxidase (HRP)-conjugated anti-rabbit IgG (1:1000; 10794347, ThermoFisher) (2 h). Protein bands developed on enhanced chemiluminescence film (10607665, Fisher Scientific). Membranes were re-probed for  $\beta$ -actin (1:2000; 4970S, Cell Signalling Technology (CST), Netherlands) to confirm equal loading. Band densitometry was performed using Image J. Three sets of adipocyte-derived EVs were used in this experiment with each set tested on one HUVEC sample ( $N = 3$ ,  $n = 3$ ).

## 2.6. Effect of adipocyte EVs on the plasma membrane expression of HUVEC adhesion molecules

HUVECs were treated with 3T3-L1 EVs from each condition (control, TNF- $\alpha$ , hypoxia, TNF- $\alpha$  + hypoxia) as described above, then washed twice and detached from the culture plate using  $200 \mu\text{L}$   $1 \times$  PBS. HUVECs were collected by centrifugation (300 g, 5 min) and then resuspended in ice-cold FACS Buffer (1% BSA (v/v) in  $1 \times$  PBS). Flow cytometry was used to assess the surface adhesion molecule profile of HUVECs. Antibodies used for flow cytometric analysis were obtained from Biolegend® (BioLegend, San Diego, CA, USA). They include; phycoerythrin (PE) anti-human CD144, allophycocyanin (APC) anti-human VCAM-1, Alexafluor 647 anti-human PECAM-1, PE anti human E-Selectin, and PECy7 anti-human ICAM-1. Cells were incubated with antibodies for 30 min at  $4^\circ\text{C}$  in darkness before being analysed on an FC500 MPL flow cytometer (Beckman Coulter equipped with 488 nm and 633 nm lasers) and data captured on MPX Cytometer List Mode Data Acquisition and Analysis Software (version 2.2). Acquisition was terminated upon recording 10,000 events, and cells gated based on their forward scatter and side scatter characteristics. Fluorescence minus one (FMO) stains were used to set the positive gates for each antibody (See supplemental information and [supplementary Fig. 3](#) for gating strategy). Three separate sets of adipocyte-derived EVs were used in this experiment and each set of EVs was tested on one individual HUVEC sample ( $N = 3$ ,  $n = 3$ ). Mean Fluorescence Intensity (MFI) was used to compare surface marker expression between treatments.

## 2.7. TNF- $\alpha$ neutralisation assay

A TNF- $\alpha$  neutralising antibody was used to determine whether EVs derived from adipocytes treated with TNF- $\alpha$ , mediated VCAM-1 upregulation in HUVECs. To neutralise any pre-existing TNF- $\alpha$ , HUVECs and adipocyte-derived EVs were incubated separately in serum-free culture medium containing 0, 3, 10, 30, 100 or 300 ng/mL TNF- $\alpha$  neutralising antibody. After 2 h, HUVEC media were removed and the EV-containing media were added to the corresponding wells. After 6 h, HUVECs were

lysed and VCAM-1 levels assessed by Western blotting, as previously described. Four sets of adipocyte-derived EVs were used in this experiment with each set tested on one HUVEC sample ( $N = 4$ ,  $n = 4$ ).

## 2.8. Analysis of 3T3-L1 cell and EV proteins

Adipocyte cell and EV lysates were assessed by Western blotting as previously described. Ten  $\mu\text{g}$  of total protein was loaded per lane for cell lysates, and 5  $\mu\text{g}$  of total protein for EV lysates. Primary antibodies directed at fatty acid binding protein-4 (FABP4; 3544S, CST), adiponectin (2789S, CST), peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ; 2443S, CST) and perilipin (9349S, Cell Signalling Technology), were used at a 1:500 dilution, and HRP-conjugated anti-rabbit IgG was used at a 1:1000 dilution. Three sets of adipocyte cell and EV lysates were used ( $N = 3$ ,  $n = 3$ ).

## 2.9. Statistical analyses

Data are presented as mean or mode  $\pm$  SEM. A one-way ANOVA with Tucky's Multiple Comparison Test was used to analyse differences. Data were analysed using GraphPad Prism (version 6; GraphPad Software Inc., USA) and  $p$ -values  $< 0.05$  were considered significant [20].

## 3. Results

### 3.1. Effect of adipocyte EVs on leukocyte-to-endothelial cell attachment


HUVECs pre-treated with TNF $\alpha$  + hypoxia-derived adipocyte EVs, increased leukocyte attachment to the same extent as the positive control (Fig. 1A). These increases were significant when compared to those observed for HUVECs not treated with EVs and HUVECs pre-treated with control adipocyte EVs (Fig. 1A). Leukocyte attachment was also greater following treatment with TNF $\alpha$  + hypoxia-derived EVs compared to hypoxia-derived adipocyte EVs. Pre-treating HUVECs with TNF- $\alpha$ -derived adipocyte EVs also increased leukocyte attachment compared to HUVECs not treated with EVs. No other differences were observed in leukocyte attachment to HUVECs after pre-treatment with adipocyte EVs. Using size-exclusion chromatography, we confirm that TNF- $\alpha$  is associated with EVs; there is no free TNF- $\alpha$  (Supplementary Fig. 1).

### 3.2. Effect of adipocyte EVs on HUVEC protein production

Following treatment of HUVECs with adipocyte EVs, Western blotting was used to examine the production of proteins involved in leukocyte attachment (Fig. 2). VCAM-1 production was increased in HUVECs treated with TNF- $\alpha$  EVs and TNF- $\alpha$  + hypoxia EVs compared to that of untreated HUVECs. No change was observed with control EVs nor hypoxia EVs (Fig. 2A). The production of other adhesion proteins including E-selectin (Fig. 2B), P-selectin (Fig. 2C) and PECAM (Fig. 2D) were unaffected by treatment with adipocyte EVs. In addition, no effect on the production levels of the vascular endothelial cell marker proteins VE-Cadherin (Fig. 2E) and eNOS (Fig. 2F) were identified.

### 3.3. Effect of adipocyte EVs on the plasma membrane expression of HUVEC adhesion molecules

Flow cytometry confirmed the expression of VCAM-1, E-selectin, PECAM-1 and ICAM-1 on HUVEC plasma membranes (Fig. 3). However, there were no statistically significant differences in the mean fluorescence intensity for each protein between any of the treatment groups. TNF- $\alpha$  treatment (no EVs) of HUVECs was used as a positive control in all experiments and showed positive expression of adhesion markers, other than for PECAM-1, which showed high expression at baseline and following all treatment conditions.



**Fig. 1.** Leukocyte attachment to HUVECs pre-treated with and without adipocyte-derived EVs.

(A) Ratio of percentage area covered by leukocytes to a confluent monolayer of HUVECs pre-treated with (C) control, (T) TNF- $\alpha$ , (H) hypoxia, and (TH) TNF- $\alpha$ +hypoxia-derived adipocyte EVs relative to that of (B; Basal) HUVECs not treated with EVs (negative control). Direct treatment with TNF- $\alpha$  acted as a positive control (mean  $\pm$  SEM; N = 3, n = 9; \*\*\* $p$  < 0.001). (B) Representative fluorescent images used to determine the percentage area of HUVECs covered by leukocytes. White dots are leukocytes stained with CellTrace™. Scale bar = 200  $\mu$ m.

### 3.4. TNF- $\alpha$ neutralisation prevents VCAM-1 upregulation in HUVEC

To determine if the upregulation of VCAM-1 in HUVECs seen following treatment with TNF- $\alpha$  and TNF- $\alpha$ +hypoxia adipocyte EVs (Fig. 2) is due to EV-associated TNF- $\alpha$ , a TNF- $\alpha$  neutralising antibody was used. Compared to untreated HUVECs, TNF- $\alpha$ +hypoxia EVs increased VCAM-1 production (Fig. 4). Compared to the level of VCAM-1 production achieved following treatment with TNF- $\alpha$ +hypoxia EVs, co-treatment with 100 ng/mL and 300 ng/mL neutralising antibody completely inhibited VCAM-1 upregulation (Fig. 4).

### 3.5. Adipokine production in adipocytes and adipocyte-derived EVs

No effect on the production of FABP4 in cell lysates due to any treatment was identified (Fig. 5A). FABP4 in was increased in EVs lysates of cells treated with TNF- $\alpha$  regardless of normoxia or hypoxia (Fig. 5B). Adiponectin production was decreased in cells treated with TNF- $\alpha$ +hypoxia compared to that of control cells, but individually, TNF- $\alpha$  and hypoxia had no effect (Fig. 5C). All treatments appeared to decrease adiponectin in EV lysates compared to that of control cells (Fig. 5D). PPAR $\gamma$  production in cell lysates decreased in response to hypoxia and TNF- $\alpha$ +hypoxia (Fig. 5E). Conversely, hypoxia- and TNF- $\alpha$ +hypoxia-derived EVs contained more PPAR $\gamma$  (Fig. 5F). Compared to that of control cells, perilipin production decreased in response to hypoxia and TNF- $\alpha$ +hypoxia (Fig. 5G). No differences in the production of perilipin in EVs were identified (Fig. 5H).

### 3.6. Effect of inflammatory and hypoxic stimuli on adipocyte EV yield and size


Compared to control adipocytes, treatment with TNF- $\alpha$ , hypoxia

and TNF- $\alpha$ +hypoxia all increased the number of EVs produced per cell (Fig. 6A). Treatment with TNF- $\alpha$ , hypoxia, or TNF- $\alpha$ +hypoxia all decreased EV size compared to control adipocytes (Fig. 6B).

## 4. Discussion

Increased adiposity has long been recognised as a risk factor for cardiovascular disease development [1]. In obesity, adipose tissue inflammation is associated with vascular inflammation [21] and one of the primary mediators of this process is the inflammatory cytokine, TNF- $\alpha$ . Whilst TNF- $\alpha$  is secreted by adipocytes themselves, levels in obese adipose tissue are predominantly raised by activated macrophages [22]. TNF- $\alpha$  desensitizes adipocytes to insulin [23], but distally and in terms of effects on vascular endothelial cells, plasma TNF- $\alpha$  levels positively correlate with adhesion molecule production, disrupted eNOS activity and oxidative stress [24]. We are the first to show that EVs derived from adipocytes residing within a hypoxic and inflammatory, TNF- $\alpha$ -containing environment (modelling the pathophysiological inflammatory nature of obese adipose tissue *in vivo*), are implicated in the onset of vascular disease by increasing vascular endothelial cell adhesion molecule production and thereby promoting leukocyte attachment. In addition, we show that inflammatory and hypoxic stimuli affect the content of adipocyte-derived EVs, as well as their yield and size.

Leukocyte-to-endothelial cell attachment was increased following treatment of HUVECs with TNF- $\alpha$  and TNF- $\alpha$ +hypoxia derived adipocyte EVs when compared to untreated HUVECs. However, when compared to the level of leukocyte attachment following treatment of HUVECs with control EVs, attachment was only increased following treatment with TNF- $\alpha$ +hypoxia EVs. This suggests that a factor(s) conferred by hypoxia, can elicit a functional effect on vascular



**Fig. 2.** Expression of adhesion and endothelial marker proteins in HUVECs following treatment with adipocyte EVs. Densitometry graphs and representative Western blots showing the ratio of (A) VCAM-1 (110 kDa), (B) E-Selectin (80 kDa), (C) P-Selectin (140 kDa), (D) PECAM (130 kDa), (E) VE-Cadherin (115 kDa) and (F) eNOS (140 kDa) expression in HUVECs following treatment with (C) control, (T) TNF- $\alpha$ , (H) hypoxia and (TH) TNF- $\alpha$  + hypoxia derived adipocyte EVs, relative to that of (B; basal) untreated HUVECs (mean  $\pm$  SEM; N = 3, n = 3; \*\*\*p < 0.001).  $\beta$ -actin confirms equal loading (bottom blots). Ten  $\mu$ g total protein per lane.


endothelial cells that exacerbates leukocyte adhesion in the presence of inflammation, and this is worthy of future investigation. The data do however, indicate that inflammatory EVs, regardless of whether they were produced under normoxic or hypoxic conditions, prime endothelial cells for subsequent leukocyte attachment.

Whilst flow cytometry confirmed the expression of the adhesion molecules VCAM-1, E-selectin, PECAM-1 and ICAM-1 on the surface of HUVEC plasma membranes, no differences between EV treatment groups were observed. Although not significant VCAM-1 surface expression did, however, show a trend to increase in cells treated with TNF- $\alpha$  and TNF- $\alpha$  + hypoxia derived adipocyte EVs and mirrored the increase in VCAM-1 detected by Western blotting. We hypothesize that in order to detect subtle changes in surface marker expression between treatment groups will likely require a very high number of HUVEC/experimental replicates and it is acknowledged further studies will be required in order to investigate differences in surface expression.


The fact that inflammatory adipocyte EVs increase HUVEC VCAM-1 production, does however, offer a mechanism through which leukocyte-to-endothelial cell attachment is achieved. TNFR1 receptors have been shown to induce VCAM-1 mRNA and protein via a pathway mediated by NF- $\kappa$ B but not ERK, p38MAPK or JNK kinase [25]. Future experiments aim to determine if the TNF- $\alpha$  delivered to HUVECs by 3T3-L1 EVs activates TNFR1 receptors and mediates VCAM-1 upregulation via

a similar mechanism. As TNF- $\alpha$  neutralisation prevents VCAM-1 upregulation, we hypothesize that this would also prevent the increase in leukocyte-to-endothelial cell attachment following incubation of HUVEC with TNF- $\alpha$  and TNF- $\alpha$  + hypoxia derived adipocyte EVs. *In vivo*, VCAM-1 is involved in the firm attachment of leukocytes to endothelial cells prior to their transmigration through the vessel wall [26,27], and whilst enhanced VCAM-1 production accompanied by leukocyte attachment is known to contribute to the progression of atherosclerosis [28,29], we can now suggest that this is at least partly mediated by EVs derived from hypoxic and inflamed adipocytes.

In this study, EVs were not processed further to remove possible contaminating “free” TNF- $\alpha$ . However, we do provide evidence that TNF- $\alpha$  co-elutes with CD63 and FABP-4 following size exclusion chromatography of EV samples and that no “free” TNF- $\alpha$  elutes in later fractions. This indicates that TNF- $\alpha$  within EV samples is EV-associated (Supplementary Fig. 1). In addition, it is important to note that the leukocytes for the leukocyte-to-endothelial attachment assay were isolated from heparin-anticoagulated blood. Heparin may activate platelets [30] causing them to expose P-selectin. P-selectin can sequentially bind to P-selectin glycoprotein on leukocytes, and this may have affected leukocyte attachment [31]. However, this effect would be equal across all experimental conditions, and as such, the results observed in this study can be attributed to 3T3-L1 EVs. It is also important



**Fig. 3.** Surface expression of HUVEC adhesion proteins following treatment with adipocyte EVs. Mean fluorescence intensity graphs showing the ratio of (A) VCAM-1 (-APC), (B) E-Selectin (-PE), (C) PECAM-1 (-AlexoFluor) and (D) ICAM-1 (-PE/Cy7) expressed on the plasma membrane of HUVECs following treatment with (C) control, (T) TNF- $\alpha$ , (H) hypoxia and (TH) TNF- $\alpha$  + hypoxia derived adipocyte EVs, relative to that of (B; basal) untreated HUVECs. Direct treatment with TNF- $\alpha$  was used as a positive control (mean  $\pm$  SEM; N = 3, n = 3 for all except ICAM-1 where N = 2, n = 2).



**Fig. 4.** TNF- $\alpha$  neutralisation inhibits endothelial VCAM upregulation following treatment with TNF- $\alpha$  + hypoxia adipocyte EVs. (A) Western blotting densitometry of VCAM-1 expression in control HUVEC lysates, and lysates of HUVECs treated with TNF- $\alpha$  + hypoxia EVs (denoted by +) in the presence of 0, 3, 10, 30, 100 and 300 ng/ml TNF- $\alpha$  neutralising antibody (mean  $\pm$  SEM; N = 4, n = 4; \*\* $p$  < 0.01). (B) Representative VCAM-1 (110 kDa; top) and  $\beta$ -actin (40 kDa; bottom) blots. Twenty  $\mu$ g total protein per lane.

to note that once isolated from blood, the leukocytes were resuspended in 0.22  $\mu$ m sterile-filtered Krebs-BSA. As such, there is a chance that bovine serum EVs smaller than 0.22  $\mu$ m were present in this buffer and that they too could have affected leukocyte adhesion. Likewise, this effect would have been equal across all conditions and consequentially the levels of leukocyte attachment observed across all experimental conditions would remain in proportion to each other.

Our results add to the growing body of literature confirming that adipocytes not only release EVs [15,16,32], but that the information conveyed through their content has a functional effect on other cell types. As such, we investigated the effect of inflammatory and hypoxic stimuli on the production of several proteins (FABP4, adiponectin, PPAR $\gamma$  and perilipin) linked to cardiovascular disease in both adipocyte cell and EV lysates. We hypothesise that differences in the content of these proteins within EVs derived from inflammatory and/or hypoxic adipocytes may confer mechanistic influences on leukocyte attachment to vascular endothelial cells. Future studies will seek to identify and elucidate such mechanisms.

Plasma levels of FABP4, a fatty-acid chaperone protein [33], have been shown to increase in obesity [34] and are associated with vascular endothelial cell dysfunction [35]. We observed no change in the FABP4 content of adipocyte cell lysates in response to inflammatory or hypoxic stimuli. This result supports a previous study whereby FABP4 production in adipocyte cell lysates was also shown not to change in response to hypoxia [36]. What is interesting, however, is that the authors of this study did identify an increased level of FABP4 in adipocyte culture media in response to hypoxia. In our study, hypoxia did not increase the FABP4 content of EVs suggesting that perhaps FABP4 is released by adipocytes in an EV-independent mechanism in this setting. However, we did observe an increase in FABP4 in the lysates of EVs derived from TNF- $\alpha$  treated adipocytes (regardless of normoxia or hypoxia). As such,




Fig. 5. FABP4, adiponectin, PPAR $\gamma$  and perilipin expression in adipocyte and adipocyte-derived EV lysates.

Densitometry graphs and representative Western blots for FABP4 and adiponectin in cell and EV lysates following (C) control, (T) TNF- $\alpha$ , (H) hypoxia and (TH) TNF- $\alpha$  + hypoxia treatment of adipocytes. FABP4 expression in (A) cell lysates (mean  $\pm$  SEM; N = 6, n = 6) and (B) EV lysates (mean  $\pm$  SEM; N = 4, n = 4) (15 kDa). Adiponectin expression in (C) cell lysates (mean  $\pm$  SEM; N = 5, n = 5; \* $p$  < 0.05) and (D) EV lysates (mean  $\pm$  SEM; N = 2, n = 2) (30 kDa). PPAR $\gamma$  expression in (E) cell lysates (mean  $\pm$  SEM; N = 6, n = 6; \*\*\* $p$  < 0.001) and (F) EV lysates (mean  $\pm$  SEM; N = 3, n = 3; \*\* $p$  < 0.01) (53 and 57 kDa). Perilipin expression in (G) cell lysates (mean  $\pm$  SEM; N = 5, n = 5; \*\*\* $p$  < 0.001) and (H) EV lysates (mean  $\pm$  SEM; N = 3, n = 3) (56 kDa). Twenty  $\mu$ g total protein per lane of cell lysates, and 5  $\mu$ g total protein per lane of EV lysates.

future experiments will aim to elucidate the influence of FABP4 within inflamed adipocyte-derived EVs on vascular endothelial dysfunction.


Adiponectin is abundant within the circulation of healthy people and exerts regulatory insulin-sensitising and anti-inflammatory effects [37,38]. Plasma adiponectin levels negatively correlate with degree of obesity [39] and therefore also correlate with increased obesity-related co-morbidities including insulin resistance, type II diabetes and cardiovascular disease [40]. We found that a combined inflammatory and hypoxic stimulus decreased adiponectin in adipocyte cell lysates. EV-associated adiponectin also appeared to decrease compared to control in response to all treatments, but particularly hypoxia. Using the same adipocyte cell model, Chen et al., showed that hypoxia prevents soluble adiponectin secretion [40], complementing our finding. Our data add to this by suggesting that at least a proportion of this reduction in adiponectin secretion under hypoxic conditions is because it is not being packaged into EVs. Reduced adiponectin within adipocyte EVs may contribute to the decrease in plasma adiponectin during obesity and thereby may contribute to the development of cardiovascular disease.

PPAR $\gamma$  is a nuclear receptor regulating the transcription of genes central to fatty acid and energy metabolism [41]. We found that hypoxia decreased PPAR $\gamma$  production in adipocyte cell lysates. Down-regulation of intracellular PPAR $\gamma$  in response to hypoxia is well documented [16,42] and is mediated by inhibition of its transcription by hypoxia inducible factor-1 (HIF-1 $\alpha$ ). Interestingly, in our study we also observed a simultaneous increase of PPAR $\gamma$  within the lysates of EVs derived from hypoxic adipocytes, confirming that adipocyte EVs are selectively packaged depending on environmental cues. We hypothesize that HIF-1 $\alpha$  may also be implicated in the selective packaging of PPAR $\gamma$  into EVs destined for release, as a further means to reduce intracellular

PPAR $\gamma$  levels. Overall, this finding highlights that EVs do not necessarily possess the same content as their cell-of-origin, but rather that their content reflects the cell-of-origin's current state.

Perilipin is a protein localised to the surface of lipid droplets and is involved in lipolysis [43]. In our study, hypoxia decreased perilipin production within adipocyte cell lysates when normalised to total protein. Wang et al. [44], previously showed no difference in total perilipin production per individual adipocyte sampled from lean and obese adipose tissue but when normalised to total protein or fat cell surface area, perilipin levels were also significantly lower in obese samples. We propose that hypoxia restricts perilipin production in adipose tissue in obesity *in vivo*. In addition, whilst perilipin levels are increased in the circulation in obese mice and humans [15] this may be because it is released in to the circulation from adipocytes as a free protein rather than being associated with EVs. This hypothesis fits with our results as we identified no difference in the perilipin content of adipocyte EVs in response to inflammatory and hypoxic stimuli.

In terms of specific EV character, our group has previously shown that EVs isolated from adipocytes express the vesicular proteins CD9, CD63, Alix, tumour susceptibility gene (TSG101)<sup>16</sup>, and now show that they also exhibit characteristic cup-shape morphology by electron microscopy (Supplementary Fig. 1). Here we show the yield and size of adipocyte-derived EVs can be modulated independently by external stimuli. TNF- $\alpha$ , hypoxia and TNF- $\alpha$  + hypoxia increased the yield of EVs/cell whilst decreasing their size. These physical changes may themselves also confer distinct functional effects. However, it is important to note that the size data presented is based solely on raw NTA values; no refractive index [45] nor mathematical modelling [46] was applied.



**Fig. 6.** Effect of inflammatory and hypoxic stimuli on adipocyte EV yield and size.

The effect of (C) control, (T) TNF- $\alpha$ , (H) hypoxia and (TH) TNF- $\alpha$ +hypoxia treatments on (A) EV yield per cell (mean  $\pm$  SEM; n = 3, n = 9; \*\*\*p < 0.001) and (B) EV size (mode  $\pm$  SEM; n = 3, n = 9; \*\*\*p < 0.001).

In summary, we provide evidence that adipocytes residing in a hypoxic and inflammatory environment produce EVs capable of inducing VCAM-1 production in vascular endothelial cells, and that this effect promotes leukocyte attachment. We also show that both inflammatory and hypoxic stimuli not only influence the adipokine content of adipocytes and their EVs, but also effect EV yield and size. Future studies will further explore the functional impact inflammatory and hypoxic adipocyte EVs have on cardiovascular disease. Indeed, the functional effects of adipocyte EVs in the setting of obesity, are likely to be not solely limited to vascular endothelial cells. We anticipate that a better understanding of the mechanisms through which adipocyte EVs negatively impact the vascular endothelium will open the door for the development of novel therapies for preventing and treating obesity-driven cardiovascular disease.

#### Conflicts of interest

The authors declared they do not have anything to disclose regarding conflict of interest with respect to this manuscript.

#### Financial support

This work was supported by the British Heart Foundation (grant reference: PG/14/51/30686).

#### Author contributions

Rebecca M. Wadey performed the experiments. Katherine D. Connolly performed pilot experiments and manuscript editing. Donna Mathew and Gareth Walters performed the flow cytometry

experiments. D. Aled Rees and Philip E. James were the Principal Investigators leading the project.

#### Acknowledgments

We would like to thank the clinical staff at University Hospital Wales for assisting with the collection of umbilical cords, the volunteers who donated blood samples, and Dr Justyna Witczak and Mrs Margaret Munnery for performing phlebotomy. We also thank Dr Errin Johnson (EM facility, Sir William Dunn School of Pathology, University of Oxford) for performing the transmission electron microscopy.

#### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.atherosclerosis.2019.01.013>.

#### References

- [1] H.B. Hubert, M. Feinleib, P.M. McNamara, W.P. Castelli, Obesity as an independent risk factor for cardiovascular disease: a 26-year follow-up of participants in the framingham heart study, *Circulation* 67 (1983) 968–977.
- [2] F.B. Ortega, D.C. Lee, P.T. Katzmarzyk, J.R. Ruiz, X. Sui, T.S. Church, S.N. Blair, The intriguing metabolically healthy but obese phenotype: cardiovascular prognosis and role of fitness, *Eur. Heart J.* 34 (2013) 389–397.
- [3] J. Kaur, A comprehensive review on metabolic syndrome, *Cardiol. Res. Pract.* 2014 (2014) 943162.
- [4] J. Ye, Z. Gao, J. Yin, Q. He, Hypoxia is a potential risk factor for chronic inflammation and adiponectin reduction in adipose tissue of ob/ob and dietary obese mice, *Am. J. Physiol. Endocrinol. Metab.* 293 (2007) E1118–E1128.
- [5] J. Ye, Emerging role of adipose tissue hypoxia in obesity and insulin resistance, *Int. J. Obes.* 33 (2009) 54–66.
- [6] M. Pasarica, J. Rood, E. Ravussin, J.M. Schwarz, S.R. Smith, L.M. Redman, Reduced oxygenation in human obese adipose tissue is associated with impaired insulin suppression of lipolysis, *J. Clin. Endocrinol. Metab.* 95 (2010) 4052–4055.
- [7] N. Hosogai, A. Fukuhara, K. Oshima, Y. Miyata, S. Tanaka, K. Segawa, S. Furukawa, Y. Tochino, R. Komuro, M. Matsuda, I. Shimomura, Adipose tissue hypoxia in obesity and its impact on adipocytokine dysregulation, *Diabetes* 56 (2007) 901–911.
- [8] P. Trayhurn, B. Wang, I.S. Wood, Hypoxia and the endocrine and signalling role of white adipose tissue, *Arch. Physiol. Biochem.* 114 (2008) 267–276.
- [9] N. Ouchi, J.L. Parker, J.J. Lugus, K. Walsh, Adipokines in inflammation and metabolic disease, *Nat. Rev. Immunol.* 11 (2011) 85–97.
- [10] M. Bulló, J. Salas-Salvadó, P. Garcia-Lorda, Adiponectin expression and adipose tissue lipolytic activity in lean and obese women, *Obes. Surg.* 15 (2005) 382–386.
- [11] N. Ouchi, S. Kihara, Y. Arita, K. Maeda, H. Kuriyama, Y. Okamoto, K. Hotta, M. Nishida, M. Takahashi, T. Nakamura, S. Yamashita, T. Funahashi, Y. Matsuzawa, Novel modulator for endothelial adhesion molecules: adipocyte-derived plasma protein adiponectin, *Circulation* 100 (1999) 2473–2476.
- [12] R. Ouedraogo, Y. Gong, B. Berzins, X. Wu, K. Mahadev, K. Hough, L. Chan, B.J. Goldstein, R. Scalia, Adiponectin deficiency increases leukocyte-endothelium interactions via upregulation of endothelial cell adhesion molecules in vivo, *J. Clin. Invest.* 117 (2007) 1718–1726.
- [13] A. Bakhai, Adipokines—targeting a root cause of cardiometabolic risk, *QJM* 101 (2008) 767–776.
- [14] P.D. Robbins, A.E. Morelli, Regulation of immune responses by extracellular vesicles, *Nat. Rev. Immunol.* 14 (2014) 195–208.
- [15] A. Eguchi, M. Lasic, A.M. Armando, S.A. Phillips, R. Katebian, S. Maraka, O. Quehenberger, D.D. Sears, A.E. Feldstein, Circulating adipocyte-derived extracellular vesicles are novel markers of metabolic stress, *J. Mol. Med. (Berl.)* 94 (2016) 1241–1253.
- [16] K.D. Connolly, I.A. Guschina, V. Yeung, A. Clayton, M.S. Draman, C. Von Ruhland, M. Ludgate, P.E. James, D.A. Rees, Characterisation of adipocyte-derived extracellular vesicles released pre- and post-adipogenesis, *J. Extracell. Vesicles* 4 (2015) 29159.
- [17] N. Aoki, S. Jin-no, Y. Nakagawa, N. Asai, E. Arakawa, N. Tamura, T. Tamura, T. Matsuda, Identification and characterization of microvesicles secreted by 3t3-l1 adipocytes: redox- and hormone-dependent induction of milk fat globule-epidermal growth factor 8-associated microvesicles, *Endocrinology* 148 (2007) 3850–3862.
- [18] S. Kralisch, T. Ebert, U. Lossner, B. Jessnitzer, M. Stumvoll, M. Fasshauer, Adipocyte fatty acid-binding protein is released from adipocytes by a non-conventional mechanism, *Int. J. Obes.* 38 (2014) 1251–1254.
- [19] M.E. Kranendonk, F.L. Visseren, B.W. van Balkom, E.N. Nolte-t Hoen, J.A. van Herwaarden, W. de Jager, H.S. Schipper, A.B. Brenkman, M.C. Verhaar, M.H. Wauben, E. Kalkhoven, Human adipocyte extracellular vesicles in reciprocal signaling between adipocytes and macrophages, *Obesity* 22 (2014) 1296–1308.
- [20] R. Lever, M.J. Rose, E.A. McKenzie, C.P. Page, Heparanase induces inflammatory cell recruitment in vivo by promoting adhesion to vascular endothelium, *Am. J. Physiol. Cell Physiol.* 306 (2014) C1184–C1190.
- [21] S. Hagita, M. Osaka, K. Shimokado, M. Yoshida, Adipose inflammation initiates

- recruitment of leukocytes to mouse femoral artery: role of adipo-vascular axis in chronic inflammation, *PLoS One* 6 (2011) e19871.
- [22] S.P. Weisberg, D. McCann, M. Desai, M. Rosenbaum, R.L. Leibel, A.W. Ferrante, Obesity is associated with macrophage accumulation in adipose tissue, *J. Clin. Invest.* 112 (2003) 1796–1808.
- [23] G.S. Hotamisligil, N.S. Shargill, B.M. Spiegelman, Adipose expression of tumor necrosis factor- $\alpha$ : direct role in obesity-linked insulin resistance, *Science* 259 (1993) 87–91.
- [24] H. Zhang, J. Zhang, Z. Ungvari, C. Zhang, Resveratrol improves endothelial function: role of  $\text{tnf}(\alpha)$  and vascular oxidative stress, *Arterioscler. Thromb. Vasc. Biol.* 29 (2009) 1164–1171.
- [25] Z. Zhou, M.C. Connell, J.D. MacEwan, TNFR1-induced NF- $\kappa$ B, but not ERK, p38MAPK or JNK activation, mediates TNF-induced ICAM-1 and VCAM-1 expression on endothelial cells, *Cell. Signal.* 19 (2007) 1238–1248.
- [26] D. Granger, E. Senchenkova, *Inflammation and the Microcirculation*, 1 edition, Morgan & Claypool Life Sciences, San Rafael, California, 2010.
- [27] L.A. Robinson, L. Tu, D.A. Steeber, O. Preis, J.L. Platt, T.F. Tedder, The role of adhesion molecules in human leukocyte attachment to porcine vascular endothelium: implications for xenotransplantation, *J. Immunol.* 161 (1990) 6931–6938.
- [28] D. Onat, D. Brillon, P.C. Colombo, A.M. Schmidt, Human vascular endothelial cells: a model system for studying vascular inflammation in diabetes and atherosclerosis, *Curr. Diabetes Rep.* 11 (2011) 193–202.
- [29] M.I. Cybulsky, K. Iiyama, H. Li, S. Zhu, M. Chen, M. Iiyama, V. Davis, J.C. Gutierrez-Ramos, P.W. Connelly, D.S. Milstone, A major role for VCAM-1, but not ICAM-1, in early atherosclerosis, *J. Clin. Invest.* 107 (2001) 1255–1262.
- [30] M. Durcin, A. Fleury, E. Taillebois, G. Hilairiet, Z. Krupova, C. Henry, S. Truchet, M. Trötz Müller, H. Köfeler, G. Mabileau, O. Hue, R. Andriantsitohaina, P. Martin, S. Le Lay, Characterisation of adipocyte-derived extracellular vesicle subtypes identifies distinct protein and lipid signatures for large and small extracellular vesicles, *J. Extracell. Vesicles* 6 (2017) 1305677.
- [31] C. Gao, B. Boylan, J. Fang, D.A. Wilcox, D.K. Newman, P.J. Newman, Heparin promotes platelet responsiveness by potentiating  $\alpha$ IIb $\beta$ 3-mediated outside-in signaling, *Blood* 117 (2011) 4946–4952.
- [32] S. Falati, Q. Liu, P. Gross, G. Merrill-Skoloff, J. Chou, E. Vandendries, A. Celi, K. Croce, B.C. Furie, B. Furie, Accumulation of tissue factor into developing thrombi in vivo is dependent upon microparticle P-selectin glycoprotein ligand 1 and platelet P-selectin, *JEM* 197 (2003) 1585–1598.
- [33] M. Furuhashi, G.S. Hotamisligil, Fatty acid-binding proteins: role in metabolic diseases and potential as drug targets, *Nat. Rev. Drug Discov.* 7 (2008) 489–503.
- [34] A. Xu, Y. Wang, J.Y. Xu, D. Stejskal, S. Tam, J. Zhang, N.M. Wat, W.K. Wong, K.S. Lam, Adipocyte fatty acid-binding protein is a plasma biomarker closely associated with obesity and metabolic syndrome, *Clin. Chem.* 52 (2006) 405–413.
- [35] G. Aragonès, P. Saavedra, M. Heras, A. Cabré, J. Girona, L. Masana, Fatty acid-binding protein 4 impairs the insulin-dependent nitric oxide pathway in vascular endothelial cells, *Cardiovasc. Diabetol.* 11 (2012) 72.
- [36] L.E. Wu, D. Samocha-Bonet, P.T. Whitworth, D.J. Fazakerley, N. Turner, T.J. Biden, D.E. James, J. Cantley, Identification of fatty acid binding protein 4 as an adipokine that regulates insulin secretion during obesity, *Mol. Metab.* 3 (2014) 465–473.
- [37] K. Makki, P. Froguel, I. Wolowczuk, Adipose tissue in obesity-related inflammation and insulin resistance: cells, cytokines, and chemokines, *ISRN Inflamm.* 2013 (2013) 139239.
- [38] Y. Arita, S. Kihara, N. Ouchi, M. Takahashi, K. Maeda, J. Miyagawa, K. Hotta, I. Shimomura, T. Nakamura, K. Miyaoka, H. Kuriyama, M. Nishida, S. Yamashita, K. Okubo, K. Matsubara, M. Muraguchi, Y. Ohmoto, T. Funahashi, Y. Matsuzawa, Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity, *Biochem. Biophys. Res. Commun.* 257 (1999) 79–83.
- [39] T.R. Aprahamian, F. Sam, Adiponectin in cardiovascular inflammation and obesity, *Int. J. Inflamm.* 2011 (2011) 376909.
- [40] B. Chen, K.S. Lam, Y. Wang, D. Wu, M.C. Lam, J. Shen, L. Wong, R.L. Hoo, J. Zhang, A. Xu, Hypoxia dysregulates the production of adiponectin and plasminogen activator inhibitor-1 independent of reactive oxygen species in adipocytes, *Biochem. Biophys. Res. Commun.* 341 (2006) 549–556.
- [41] E.A. Ivanova, A. Parolari, V. Myasoedova, A.A. Melnichenko, Y.V. Bobryshev, A.N. Orekhov, Peroxisome proliferator-activated receptor (ppar) gamma in cardiovascular disorders and cardiovascular surgery, *J. Cardiol.* 66 (2015) 271–278.
- [42] Z. Yun, H.L. Maecker, R.S. Johnson, A.J. Giaccia, Inhibition of ppar gamma 2 gene expression by the hif-1-regulated gene *dec1/stra13*: a mechanism for regulation of adipogenesis by hypoxia, *Dev. Cell* 2 (2002) 331–341.
- [43] P.A. Kern, G. Di Gregorio, T. Lu, N. Rassouli, G. Ranganathan, Perilipin expression in human adipose tissue is elevated with obesity, *J. Clin. Endocrinol. Metab.* 89 (2004) 1352–1358.
- [44] Y. Wang, S. Sullivan, M. Trujillo, M.J. Lee, S.H. Schneider, R.E. Brolin, Y.H. Kang, Y. Werber, A.S. Greenberg, S.K. Fried, Perilipin expression in human adipose tissues: effects of severe obesity, gender, and depot, *Obes. Res.* 11 (2003) 930–936.
- [45] C. Gardiner, M. Shaw, P. Hole, J. Smith, D. Tannetta, C.W. Redman, I.L. Sargent, Measurement of refractive index by nanoparticle tracking analysis reveals heterogeneity in extracellular vesicles, *J. Extracell. Vesicles* 24 (2014) 25361.
- [46] E. van der Pol, F.A.W. Coumans, A.E. Grootemaat, C. Gardiner, I.L. Sargent, P. Harrison, A. Sturk, T.G. van Leeuwen, R. Nieuwland, Particle size distribution of exosomes and microvesicles determined by transmission electron microscopy, flow cytometry, nanoparticle tracking analysis, and resistive pulse sensing, *JTH* 12 (2014) 1182–1192.