

References

1. Baggen, J. *et al.* (2018) The life cycle of non-polio enteroviruses and how to target it. *Nat. Rev. Microbiol.* 16, 368–381
2. Tapparel, C. *et al.* (2013) Picornavirus and enterovirus diversity with associated human diseases. *Infect. Genet. Evol.* 14, 282–293
3. Muehlenbachs, A. *et al.* (2015) Tissue tropism, pathology and pathogenesis of enterovirus infection. *J. Pathol.* 235, 217–228
4. Lin, J.Y. and Shih, S.R. (2014) Cell and tissue tropism of enterovirus 71 and other enteroviruses infections. *J. Biomed. Sci.* 21, 18–0127
5. Drummond, C.G. *et al.* (2017) Enteroviruses infect human enteroids and induce antiviral signaling in a cell lineage-specific manner. *Proc. Natl. Acad. Sci. U. S. A.* 114, 1672–1677
6. Ward, T. *et al.* (1998) Role for beta2-microglobulin in echovirus infection of rhabdomyosarcoma cells. *J. Virol.* 72, 5360–5365
7. Marjomaki, V. *et al.* (2002) Internalization of echovirus 1 in caveolae. *J. Virol.* 76, 1856–1865
8. Morosky, S. *et al.* (2019) The neonatal Fc receptor is a pan-echovirus receptor. *Proc. Natl. Acad. Sci. U. S. A.* 116, 3758–3763
9. Pyzik, M. *et al.* (2015) FcRn: The architect behind the immune and nonimmune functions of IgG and albumin. *J. Immunol.* 194, 4595–4603
10. Abzug, M.J. (2004) Presentation, diagnosis, and management of enterovirus infections in neonates. *Paediatr. Drugs* 6, 1–10

Forum

Influenza
Neuraminidase:
Underrated Role in
Receptor BindingFeng Wen¹ and Xiu-Feng Wan^{1,*}

Influenza virus neuraminidase cleaves sialic acid groups from cell glycoproteins, enabling release of the virus from host cells. Neuraminidase also contributes to virus binding to the sialic acid groups of cell glycoproteins, which could complement the receptor-binding function of hemagglutinin, enhancing enzymatic activities of neuraminidase, and facilitate virus infection.

Influenza A Tropism through
Binding Affinity

Influenza A viruses (IAVs) infect humans and a wide range of animal species,

including birds, pigs, horses, dogs, and marine mammals (e.g., seals and whales). The initial step in viral infection is the binding to sialic acid receptors, typically avian-like receptors (sialic acid- α 2,3-galactose linkage, SA2,3Gal) and human-like receptors (sialic acid- α 2,6-galactose linkage, SA2,6Gal) on the host cell surface glycoproteins. The last step in viral infection is the cleavage of sialic acid receptors from glycoproteins and glycolipids by neuraminidase (NA) and release of virus particles after budding from the host cell plasma membrane. Gaining binding affinity to human-like SA2,6Gal receptors is considered a key prerequisite for avian IAVs that cause pandemics or epidemics among humans. The IAV glycoprotein hemagglutinin (HA) attaches to host receptors during virus entry, and the receptor-binding specificity of HA is well documented to affect IAV host tropism. We review recent studies demonstrating that NA, the other IAV glycoprotein, contributes to virus binding to the sialic acid receptors at host cell glycoproteins and that, such receptor-binding contributions can complement those of HA, enhance NA enzymatic activities, and facilitate virus infection and host adaptation.

Receptor Binding of IAV NA

The NA of IAVs belongs to the exosialidase family of enzymes which cleave the α -ketosidic linkage between synthetic substrates and the terminal sialic acid residues in oligosaccharides, glycoproteins, glycolipids, or colominic acids. It is generally accepted that the functions of HA and NA are, to some extent, contrary: HA binds to sialic acid receptors on host cells and mediates virus attachment and entry, whereas NA cleaves the α -ketosidic linkage between the terminal sialic acid residues and the remaining substructures. However, it has been reported that, in addition to their enzymatic functions, NA proteins, such as N1, N2, and N9, also have sialic acid receptor-binding functions. For example, subtype N9 from A/tern/

Australia/G70C/75(H11N9) alone hemagglutinates animal erythrocytes by using a second receptor-binding site (SRBS), which is also referred to as the hemadsorption site, that is distinct from the NA catalytic site [1].

Structural analyses suggested that the SRBS of NA consists of three loops (367–372, 400–403, and 432) that interact with the Neu5Ac moiety in chair conformation, which is similar to that of the receptor binding site (RBS) of HA [2]. In contrast, the NA catalytic site interacts with the sialic acid receptor in a twisted boat conformation [2]. Sequence analyses showed that these residues in the SRBS of NA are mostly conserved among avian IAVs but not in human and swine IAVs [2]. Of interest, the sialic acid binding function of the SRBS of NA has been so far identified primarily in avian IAVs [3]. On the other hand, sequence diversity in the SRBS across different NA subtypes, or even within the same NA subtype, is well documented to affect the receptor-binding ability of NA. For example, exchange of amino acids in the loops at positions 368–370 and positions 399–403 of N9 into those corresponding positions in the N2 by site-directed mutagenesis made N2 acquire the hemagglutination activity as shown in N9 [4]. The pandemic viruses A/Singapore/1/57(H2N2) and A/Brevig Mission/1/18 (H1N1) differed from their avian counterpart precursors with abolished or reduced hemadsorption activity due to variations in the SRBS of NA [3].

Current studies suggest that, through its glycan binding, the SRBS of NA can increase the enzymatic activities on multivalent substrates but not monovalent substrates [3]. The NA of a human H7N9 virus enhanced overall virus binding to SA2,6Gal but not to SA2,3Gal via the SRBS of NA when the whole virus is used in the assays [5]. Interestingly, another study, when using recombinant NA proteins, showed that the NA of human

H7N9 virus, with a mutation A401T in the SRBS of N9, increased virus binding and cleavage activities to SA_{2,3}Gal, and none of these wild type or mutant NA recombinant proteins had significant binding affinities to SA_{2,6}Gal [6]. Interestingly, the T401A substitution of N9 protein was acquired prior to the other two mutations, G186V and Q226L, in the RBS of H7 protein, which altered the receptor-binding properties of the novel H7N9 virus. Of note, the binding specificity was determined by biolayer interferometry in [5] whereas it was determined by glycan array in [6]. The conflicting results in these studies could be affected by experimental conditions (i.e., whole virus particle or recombinant proteins) and/or the glycan binding detection methods. Although the role of SRBS of NA in the interaction between influenza virus and the receptors of the host cells during virus infection remains largely unknown, these findings suggest that mutations in the SRBS of NA could affect the HA–NA balance and facilitate crossing of host species barriers by H7N9 avian IAVs.

While receptor binding via the SRBS of NA enhances NA enzymatic activity, receptor binding via the catalytic site of NA displaying low enzymatic activity may complement HA receptor binding and serve as a receptor-binding protein during virus entry. In 2005, the N2 protein of human seasonal H3N2 virus gained a D151G substitution at the NA catalytic site, enabling the viruses to agglutinate erythrocytes in an oseltamivir-sensitive manner, increasing binding affinities to 3'-sialyllactosamine and 6'-sialyllactosamine higher than the corresponding HA affinities to those sialosides [7]. The introduction of mutation D151G into N1 and N2 reduces NA enzymatic activity and enables preferential binding to SA_{2,3}Gal [7,8]. Of note, the entry of human H3N2 viruses with mutation D151G into Madin–Darby canine kidney (MDCK) cells was inhibited by oseltamivir, which can block the enzymatic

site of NA [8]. Sequence comparison between clinical samples and virus isolates suggested that D151G was acquired during virus isolation and propagation for these recent human H3N2 viruses, which replicate poorly in MDCK cells, and the increased receptor binding due to D151G at the catalytic site of NA could help cell adaptation and facilitate viral infection [9]. Of note, the mutation D151G was not identified in the isolates recovered using MDCK-siat1 cells, which overexpress human SA_{2,6}Gal. On the other hand, H150R at the catalytic site of NA enabled hemagglutination ability for H3N2 viruses, and H150R was demonstrated as not a cell-adapted mutation (like D151G) but a natural mutation that emerged in 2005 and has become predominant in recent H3N2 viruses [10]. For human H1N1 viruses, G147R in the NA catalytic site of N1, which occurs at low frequencies in human pandemic and seasonal H1N1 viruses, enables an HA binding-deficient mutant to use N1 as the receptor-binding protein [11]. Guo *et al.* [12] showed, by using a biolayer interferometry assay, that a low-activity NA can contribute to virion-receptor binding by its catalytic site in a NA inhibitor-sensitive manner. Nevertheless, the function of these NA substitutions in the process of virus entry remains largely unknown.

Concluding Remarks

In summary, recent studies suggest that NA plays important roles in virus host adaptation and evolution by serving as a receptor-binding protein. Increasing evidence suggests that possibly all avian NAs have an SRBS distinct from, but close to the catalytic site. However, the receptor-binding function of NA in IAV biology is underrated and needs further research. First, the roles of NA receptor binding in host/tissue/cell tropism is far from being understood; the relationship between receptor binding properties of NA and those of HA during virus entry is not clear. Prior studies suggested that the

affinity to certain types of sialic acid receptors might be higher for NA protein than for HA protein, and further studies are needed to determine whether NA could affect the spectrum of receptors to which the viruses can bind. In addition, the role of the SRBS of NA in the interaction between influenza virus and its host receptors during virus infection is unclear, although the receptor-binding activities through the SRBS of NA can enhance the enzymatic activities of NA over multivalent substrates. Finally, the effect(s) of the balance between HA and NA functional activities during virus evolution (e.g., compatibility between HA and NA during genetic reassortment) is still not fully understood.

Acknowledgments

This study was supported by the National Institutes of Health (grant number R01AI116744).

¹Department of Basic Sciences, College of Veterinary Medicine, Mississippi State University, Mississippi State, MS, USA

*Correspondence:
wan@cvm.msstate.edu (X.-F. Wan).
<https://doi.org/10.1016/j.tim.2019.03.001>

© 2019 Elsevier Ltd. All rights reserved.



References

1. Webster, R.G. *et al.* (1987) Antigenic structure and variation in an influenza virus N9 neuraminidase. *J. Virol.* 61, 2910–2916
2. Varghese, J.N. *et al.* (1997) Structural evidence for a second sialic acid binding site in avian influenza virus neuraminidases. *Proc. Natl. Acad. Sci. U. S. A.* 94, 11808–11812
3. Uhlenndorff, J. *et al.* (2009) Functional significance of the hemadsorption activity of influenza virus neuraminidase and its alteration in pandemic viruses. *Arch. Virol.* 154, 945–957
4. Nuss, J.M. and Air, G.M. (1991) Transfer of the hemagglutinin activity of influenza virus neuraminidase subtype N9 into an N2 neuraminidase background. *Virology* 183, 496–504
5. Benton, D.J. *et al.* (2017) Role of neuraminidase in influenza A(H7N9) virus receptor binding. *J. Virol.* 91, e02293–16
6. Dai, M. *et al.* (2017) Mutation of the second sialic acid-binding site, resulting in reduced neuraminidase activity, preceded the emergence of H7N9 influenza A virus. *J. Virol.* 91, e00049–17
7. Zhu, X. *et al.* (2012) Influenza virus neuraminidases with reduced enzymatic activity that avidly bind sialic acid receptors. *J. Virol.* 86, 13371–13383
8. Gulati, S. *et al.* (2013) Human H3N2 influenza viruses isolated from 1968 to 2012 show varying preference for receptor substructures with no apparent consequences for disease or spread. *PLoS One* 8, e66325
9. Xue, K.S. *et al.* (2018) Cooperating H3N2 influenza virus variants are not detectable in primary clinical samples. *mSphere* 3, e00552–17

10. Mogling, R. *et al.* (2017) Neuraminidase-mediated haemagglutination of recent human influenza A(H3N2) viruses is determined by arginine 150 flanking the neuraminidase catalytic site. *J. Gen. Virol.* 98, 1274–1281
11. Hooper, K.A. and Bloom, J.D. (2013) A mutant influenza virus that uses an N1 neuraminidase as the receptor-binding protein. *J. Virol.* 87, 12531–12540
12. Guo, H. *et al.* (2018) Kinetic analysis of the influenza A virus HA/NA balance reveals contribution of NA to virus-receptor binding and NA-dependent rolling on receptor-containing surfaces. *PLoS Pathog.* 14, e1007233