



Technical Notes & Surgical Techniques

Techniques of intracranial aneurysm wall biopsy[☆]Sajjad Muhammad (MD, PhD)^{*}, Mika Niemelä (MD, PhD)

Department of Neurosurgery, University of Helsinki and Helsinki University Hospital, Helsinki, Finland



ARTICLE INFO

Keywords:

Intracranial aneurysm biopsy
Video demonstration
Technical aspects
Aneurysm wall research

ABSTRACT

Current treatment modalities for the treatment of intracranial aneurysms including surgical clipping and endovascular coiling are invasive and have some treatment risks. Since not all aneurysms rupture, it is critical to detect rupture prone aneurysms. Molecular and cellular analysis of aneurysm tissue may provide understanding about pathobiology of aneurysm rupture and to develop imaging techniques to detect rupture prone aneurysms. For more than 15-years we have collected samples to identify pathological processes in the aneurysm wall itself predisposing to rupture. This has opened a new field of research leading to novel findings and multiple scientific publications. Surgical techniques of sampling the aneurysm dome have never been demonstrated so therefore many neurosurgeons are reluctant to take biopsies for research. Now we demonstrate with an intraoperative video the techniques of sampling the aneurysm dome after clipping an incidental right-sided 5-mm unruptured MCA aneurysm in a 58-year-old hypertensive male with past long history of smoking through lateral supraorbital approach. A focused opening of the Sylvian fissure was performed and the aneurysm was clipped using standard techniques. After placement of a titanium clip, ICG and Doppler were performed to ensure patency of both M2 vessels and the aneurysm was punctured. The aneurysm dome was then held in place with the suction and cut with microscissors for research purposes. Another titanium clip was placed (Video 1). The clinical course was uneventful. This technical note will help young neurosurgeons to contribute actively in aneurysm research also potentially to find non-invasive methods to prevent aneurysms from rupture.

1. Introduction

Ruptured intracranial aneurysm (ICA) leading to subarachnoid hemorrhage (SAH) is a devastating disease killing around 50% of patients. Current treatment modalities including clipping and coiling are invasive and have some treatment risks. Since not all aneurysms rupture, it is critical to detect rupture prone aneurysms. State of the art research shows that Inflammation plays critical role in formation and rupture of intracranial aneurysms as well after the SAH [1–5,7]. Multiple publications have reported a critical role of inflammation and inflammatory cells in rupture of intracranial aneurysms [6,8–10,12–18]. Molecular and cellular analysis of aneurysm tissue may provide better understanding about pathobiology of aneurysm rupture and to develop imaging techniques to detect rupture prone aneurysms [11,16,17]. Sampling of aneurysm tissue during microsurgical clipping is the key to perform molecular and pathological investigations. Surgical technique of sampling the aneurysm dome has never been demonstrated. Due to lack of demonstrative publications many neurosurgeons are reluctant to take biopsy for research in

addition to the preference of endovascular therapy in many centres making aneurysm wall biopsies practically almost impossible.

2. Methods

Using intraoperative video material, we demonstrate the technique of sampling the aneurysm after clipping in a case of a hypertensive 58-year-old man who has smoked previously for decades and harboring incidental right middle cerebral artery aneurysm. We performed surgical clipping and aneurysm biopsy through lateral supraorbital approach using OPMI PENTERO 900 microscope (Car-Zeiss, Germany) using standard instruments.

3. Results and discussion

To expose the MCA aneurysm, a focused opening of Sylvian fissure was performed and aneurysm dome was dissected from the surrounded tissue. The aneurysm was clipped using standard tools. After placement of clip we performed Doppler US and video ICG to ensure patency of

Abbreviation: ICA, Intracranial aneurysm

[☆] Conflict of interest: None.

^{*} Corresponding author.

E-mail address: ext-sajjad.muhammad@hus.fi (S. Muhammad).

<https://doi.org/10.1016/j.inat.2019.01.022>

Received 16 January 2019; Accepted 28 January 2019

2214-7519/© 2019 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

both M2 vessels. Then the aneurysm was punctured using a 23G needle. Microscope was then focused on the aneurysm dome which was held with the suction at very low pressure and half of the dome was cut with micro-scissors. The suction was then removed and the aneurysm dome was held with long micro tweezer and cut further to complete the biopsy (Video 1).

This video demonstration showing technique of sampling will help young neurosurgeons to contribute actively to take more aneurysm biopsies and contribute in aneurysm research. Hence our article may help to enhance aneurysm wall research and open novel directions of research.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.inat.2019.01.022>.

Disclosures

The authors have no conflict of interest and have nothing to disclose.

References

- [1] S.R. Chaudhry, A. Guresir, B. Stoffel-Wagner, R. Fimmers, T.M. Kinfe, D. Dietrich, A. Lamprecht, H. Vatter, E. Guresir, S. Muhammad, Systemic high-mobility group box-1: a novel predictive biomarker for cerebral vasospasm in aneurysmal subarachnoid hemorrhage, *Crit. Care Med.* 46 (11) (2018) e1023–e1028.
- [2] S.R. Chaudhry, E. Guresir, H. Vatter, T.M. Kinfe, D. Dietrich, A. Lamprecht, S. Muhammad, Aneurysmal subarachnoid hemorrhage lead to systemic upregulation of IL-23/IL-17 inflammatory axis, *Cytokine* 97 (2017) 96–103.
- [3] S.R. Chaudhry, A. Hafez, B. Rezai Jahromi, T.M. Kinfe, A. Lamprecht, M. Niemela, S. Muhammad, Role of damage associated molecular pattern molecules (DAMPs) in aneurysmal subarachnoid hemorrhage (aSAH), *Int. J. Mol. Sci.* 19 (2018).
- [4] S.R. Chaudhry, B. Stoffel-Wagner, T.M. Kinfe, E. Guresir, H. Vatter, D. Dietrich, A. Lamprecht, S. Muhammad, Elevated systemic IL-6 levels in patients with aneurysmal subarachnoid hemorrhage is an unspecific marker for post-SAH complications, *Int. J. Mol. Sci.* 18 (2017).
- [5] J. Frosen, A. Piippo, A. Paetau, M. Kangasniemi, M. Niemela, J. Hernesniemi, J. Jaaskelainen, Remodeling of saccular cerebral artery aneurysm wall is associated with rupture: histological analysis of 24 unruptured and 42 ruptured cases, *Stroke* 35 (2004) 2287–2293.
- [6] J. Frosen, A. Piippo, A. Paetau, M. Kangasniemi, M. Niemela, J. Hernesniemi, J. Jaaskelainen, Growth factor receptor expression and remodeling of saccular cerebral artery aneurysm walls: implications for biological therapy preventing rupture, *Neurosurgery* 58 (2006) 534–541 discussion 534–541.
- [7] J. Frosen, R. Tulamo, A. Paetau, E. Laaksamo, M. Korja, A. Laakso, M. Niemela, J. Hernesniemi, Saccular intracranial aneurysm: pathology and mechanisms, *Acta Neuropathol.* 123 (2012) 773–786.
- [8] E.I. Gaal, T. Tammela, A. Anisimov, S. Marbacher, P. Honkanen, G. Zarkada, V.M. Leppanen, T. Tatlisumak, J. Hernesniemi, M. Niemela, K. Alitalo, Comparison of vascular growth factors in the murine brain reveals placenta growth factor as prime candidate for CNS revascularization, *Blood* 122 (2013) 658–665.
- [9] E. Laaksamo, R. Tulamo, A. Liiman, M. Baumann, R.M. Friedlander, J. Hernesniemi, M. Kangasniemi, M. Niemela, A. Laakso, J. Frosen, Oxidative stress is associated with cell death, wall degradation, and increased risk of rupture of the intracranial aneurysm wall, *Neurosurgery* 72 (2012) 109–117.
- [10] S. Marbacher, J. Marjamaa, K. Bradacova, M. von Gunten, P. Honkanen, U. Abo-Ramadan, J. Hernesniemi, M. Niemela, J. Frosen, Loss of mural cells leads to wall degeneration, aneurysm growth, and eventual rupture in a rat aneurysm model, *Stroke* 45 (2014) 248–254.
- [11] J. Marjamaa, R. Tulamo, J. Frosen, U. Abo-Ramadan, J.A. Hernesniemi, M.R. Niemela, M. Kangasniemi, Occlusion of neck remnant in experimental rat aneurysms after treatment with platinum- or polyglycolic-poly(lactic acid)-coated coils, *Surg. Neurol.* 71 (2009) 458–465 discussion 465.
- [12] E. Ollikainen, R. Tulamo, J. Frosen, S. Lehti, P. Honkanen, J. Hernesniemi, M. Niemela, P.T. Kovanen, Mast cells, neovascularization, and microhemorrhages are associated with saccular intracranial artery aneurysm wall remodeling, *J. Neuropathol. Exp. Neurol.* 73 (2014) 855–864.
- [13] E. Ollikainen, R. Tulamo, S. Kaitainen, P. Honkanen, S. Lehti, T. Liimatainen, J. Hernesniemi, M. Niemela, P.T. Kovanen, J. Frosen, Macrophage infiltration in the saccular intracranial aneurysm wall as a response to locally lysed erythrocytes that promote degeneration, *J. Neuropathol. Exp. Neurol.* 77 (2017) 890–903.
- [14] E. Ollikainen, R. Tulamo, S. Kaitainen, P. Honkanen, S. Lehti, T. Liimatainen, J. Hernesniemi, M. Niemela, P.T. Kovanen, J. Frosen, Macrophage infiltration in the saccular intracranial aneurysm wall as a response to locally lysed erythrocytes that promote degeneration, *J. Neuropathol. Exp. Neurol.* 77 (2018) 890–903.
- [15] E. Ollikainen, R. Tulamo, S. Lehti, J. Hernesniemi, M. Niemela, P.T. Kovanen, J. Frosen, Myeloperoxidase associates with degenerative remodeling and rupture of the saccular intracranial aneurysm wall, *J. Neuropathol. Exp. Neurol.* 77 (2018) 461–468.
- [16] R. Tulamo, J. Frosen, J. Hernesniemi, M. Niemela, Inflammatory changes in the aneurysm wall: a review, *J. Neurointerv. Surg.* 2 (2010) 120–130.
- [17] R. Tulamo, J. Frosen, J. Hernesniemi, M. Niemela, Inflammatory changes in the aneurysm wall: a review, *J. Neurointerv. Surg.* 10 (2018) i58–i67.
- [18] R. Tulamo, J. Frosen, S. Junnikkala, A. Paetau, J. Pitkaniemi, M. Kangasniemi, M. Niemela, J. Jaaskelainen, E. Jokitalo, A. Karatas, J. Hernesniemi, S. Meri, Complement activation associates with saccular cerebral artery aneurysm wall degeneration and rupture, *Neurosurgery* 59 (2006) 1069–1076 discussion 1076–1067.