



Original Article

Is There a Role for an 18F-fluorodeoxyglucose-derived Biological Boost in Squamous Cell Anal Cancer?



A. Sabbagh^{*}, C. Jacobs^{*}, R. Cooke^{*†}, K.-Y. Chu^{*†}, S.M. Ng[‡], V.Y. Strauss[§], P.S. Virdee[§], M.A. Hawkins[†], M.C. Aznar[¶], R. Muirhead^{*}

^{*} Department of Oncology, Oxford University Hospitals Trust, Oxford, UK

[†] CRUK MRC Oxford Institute for Radiation Oncology, University of Oxford, Oxford, UK

[‡] Oncology Clinical Trials Office, Department of Oncology, University of Oxford, Oxford, UK

[§] Centre for Statistics in Medicine, NDORMS, University of Oxford, Oxford, UK

[¶] Manchester Cancer Research Centre, Division of Cancer Sciences, School of Medical Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester, UK

Received 16 August 2018; received in revised form 1 November 2018; accepted 7 November 2018

Abstract

Aims: To investigate the potential role for a biological boost in anal cancer by assessing whether subvolumes of high 18F-fluorodeoxyglucose (FDG) avidity, identified at outset, are spatially consistent during a course of chemoradiotherapy (CRT).

Materials and methods: FDG-positron emission tomography (FDG-PET) scans from 21 patients enrolled into the ART study (NCT02145416) were retrospectively analysed. In total, 29 volumes including both primary tumours and involved nodes >2 cm were identified. FDG-PET scans were carried out before treatment and on day 8 or 9 of CRT. FDG subvolumes were created using a percentage of maximum FDG avidity at thresholds of 34%, 40%, 50%, on the pre-treatment scans, and 70% and 80% on the subsequent scans. Both FDG-PET scans were deformably registered to the planning computed tomography scan. The overlap fraction and the vector distance were calculated to assess spatial consistency. FDG subvolumes for further investigation had an overlap fraction >0.7, as this has been defined in previous publications as a 'good' correlation.

Results: The median overlap fractions between the diagnostic FDG-PET subvolumes 34%, 40% and 50% of maximum standardised uptake value (SUV_{max}) and subsequent FDG-PET subvolumes of 70% of SUV_{max} were 0.97, 0.92 and 0.81. The median overlap fraction between the diagnostic FDG-PET subvolumes 34%, 40% and 50% and subsequent FDG-PET subvolumes of 80% were 1.00, 1.00 and 0.92. The median (range) vector distance values between diagnostic FDG-PET subvolumes 34%, 40% and 50% and subsequent FDG-PET subvolumes of 80% were 0.74 mm (0.19–2.94) 0.74 mm (0.19–3.39) and 0.71 mm (0.2–3.29), respectively. Twenty of 29 volumes (69.0%) achieved a threshold > 0.7 between the FDG 50% subvolume on the diagnostic scan and the FDG 80% subvolume on the subsequent scan.

Conclusion: FDG-avid subvolumes identified at baseline were spatially consistent during a course of CRT treatment. The subvolume of 50% of SUV_{max} on the pre-treatment scan could be considered as a potential target for dose escalation.

Crown Copyright © 2018 Published by Elsevier Ltd on behalf of The Royal College of Radiologists. All rights reserved.

Keywords: Anal cancer; biological boost; FDG-PET scan; radiotherapy

Introduction

Radical chemoradiotherapy (CRT) is the established standard of care for localised squamous cell carcinoma of the anus, delivering doses up to 66 Gy in 1.8–2.2 Gy/fraction

concurrently with chemotherapy [1]. CRT provides 3-year progression-free survival of 73% [2]. However, despite the widespread use of intensity-modulated radiotherapy, acute and chronic toxicity remain significant. Acute toxicities include radiation dermatitis, pain, gastrointestinal and genitourinary disturbance; these usually resolve 2–3 months after CRT. Late toxicities, however, can be lifelong and significantly affect quality of life; these include diarrhoea, faecal incontinence, dyspareunia and erectile dysfunction [3–6].

Author for correspondence: R. Muirhead, Level 2 Cancer Offices, Department of Oncology, Cancer and Haematology Centre, Churchill Hospital, Oxford OX3 7LE, UK.

E-mail address: rebecca.muirhead@oncology.ox.ac.uk (R. Muirhead).

<https://doi.org/10.1016/j.clon.2018.11.034>

0936-6555/Crown Copyright © 2018 Published by Elsevier Ltd on behalf of The Royal College of Radiologists. All rights reserved.

Treatment failure occurs most often in patients with locally advanced disease, with about 75% of relapses occurring at the site of the original tumour [7–10]. A tumour control probability model has suggested that dose escalation may improve tumour control by potentially reducing local relapse in these patients [11]. PLATO (Personalising Anal cancer RadioTherapy dose, ISRCTN88455282), the international anal cancer trial funded by Cancer Research UK, is investigating the role of dose escalation to the primary site of gross tumour in locally advanced disease [12]. Dose escalation to the whole gross tumour would probably add acute and late toxicity to the treatment; so the use of functional imaging to identify a smaller biological boost is of interest, as smaller volumes are recognised to minimise toxicity [13].

Over 90% of squamous cell anal tumours are 18F-fluorodeoxyglucose (FDG) avid on positron emission tomography (PET) scan [14] and there has been significant interest in the role of FDG-PET in anal cancer management. Trautmann and Zuger [15] reported in a study of 21 patients, that the diagnostic FDG-PET identified additional nodes and metastasis not seen on magnetic resonance imaging and computed tomography (CT), having implication for prognosis and radiotherapy planning. The metabolically active volume on diagnostic images has also correlated with local recurrence in small series of 23 and 45 patients [16,17]. Metabolic tumour volume probably correlates with lesion size, therefore this finding is in keeping with the phase III data highlighting that larger lesions do worse [18]. However, Kidd *et al.* [19] identified a new finding that increased standardised uptake value (SUV) at outset correlated with disease-free survival and the presence of persistence or recurrence on subsequent FDG-PETs in 77 patients. A number of single-centre publications have reviewed between 23 and 110 patients with diagnostic FDG-PET and repeat FDG-PET completion of CRT at time points ranging from 3 to 36 weeks [20–22]. In these studies, the metabolic response correlated with one or more of: local control rate, progression-free survival, cancer-specific survival or overall survival. Finally, Hong *et al.* [16] looked at 23 patients with FDG-PET at outset and after 30 Gy of radiotherapy. They found that the interim scan metabolic tumour volume and total lesion glycolysis correlated with freedom from local progression.

All of the FDG-PET data in anal cancer to date have looked at the predictive value of diagnostic or repeat FDG-PET. The authors are unaware of any other publications investigating the role of a biological boost in anal carcinoma. In other tumour types treated by concomitant CRT, such as non-small cell lung carcinoma (NSCLC), head and neck squamous cell carcinomas and pancreatic adenocarcinoma, FDG-PET/CT scans have been reported to identify tumour subvolumes at greater risk of relapse [23–29]. There are ongoing trials investigating the use of biological boosts in head and neck squamous cell carcinomas and NSCLC [30–32].

The ART (Anal squamous cell carcinoma: investigation of functional imaging during chemoradiotherapy) study (NCT02145416) is a feasibility trial investigating the use of

functional imaging to predict response in anal cancer. All patients receive baseline FDG-PET/CT scans and further scans at fractions 8–10. We carried out a post-hoc analysis on images obtained in this trial. Our aim was to investigate the feasibility of defining a biological boost in anal squamous cell carcinoma by assessing whether FDG-avid subvolumes, identified at baseline, are spatially consistent in the second week of a course of CRT.

Materials and Methods

Patient Population

We identified 21 patients with paired scans carried out at diagnosis and at fraction 8 or 9 of CRT, obtained from the ART study. Twenty-nine patients were recruited however, four patients withdrew consent before any imaging and a further four patients withdrew consent after their initial scans due to ongoing social and medical issues. The ART study is a single-arm, single-centre imaging study of patients receiving radical CRT for anal cancer within Oxford University Hospitals, evaluating changes in functional imaging during CRT treatment. ART recruited patients who were aged 18 years or above, had histologically confirmed invasive primary squamous carcinoma of the anus, had any tumour stage \geq T2N0 and were fit to receive radical CRT with curative intent. All patients had given informed consent in accordance with the Code of Ethics of the World Medical Association (Declaration of Helsinki). Ethical approval incorporated post-hoc analysis of images to answer additional questions.

Chemoradiotherapy

CRT was delivered according to UK-based guidance [33]. In summary, patients received 50.4 or 53.2 Gy to the primary tumour depending on TNM stage, 50.4 Gy to involved nodes and 40 Gy prophylactic dose to non-involved nodes, all in 28 fractions using a simultaneous integrated boost. Three patients were treated within the PLATO study, two with a dose of 61.6 Gy to the primary tumour while maintaining 40 Gy to the prophylactic nodes; the other patient was treated with a reduced dose of 41.4 Gy to the primary tumour and 34.5 Gy to elective nodes in 23 fractions with simultaneous integrated boost. Treatment was delivered using intensity-modulated radiotherapy or volumetric-modulated arc therapy, calculated using a type B algorithm, prescribing 100% to the median dose of the planned target volume boost. Chemotherapy was mitomycin 12 mg/m² day 1 and capecitabine 825 mg/m² orally twice a day on radiotherapy treatment days.

FDG-PET/CT Scanning

All patients had a routine, full-body, diagnostic FDG-PET/CT scan carried out at baseline. As part of the ART study, 18 patients underwent a second FDG-PET/CT on fraction 8 and three on fraction 9 of their CRT.

All scans were carried out using GE Discovery 690 scanners (GE Healthcare, Buckinghamshire, UK). Patients were positioned supine with arms above the head, on a flat-top couch using radiotherapy immobilisation equipment to replicate their treatment position. After fasting for 6 h and ensuring that the blood glucose was <10 mmol/l, FDG was injected at a dose of 4 MBq/kg (up to 600 MBq). FDG-PET/CT acquisition started after an uptake time of 75 min. The initial FDG-PET/CT scanned the whole body. For the second FDG-PET/CT, eight patients underwent an image limited to the pelvis, 13 patients had a further full-body image.

Scans were carried out in three-dimensions with a scan time of 4 min at each bed position. For the CT phase, 120 kV automA (maximum 250 mA), noise index 25.0 0.5 s/rotation, pitch 0.984:1, 3.75 mm slice width were used. Attenuation corrected FDG-PET/CT images were used in the analysis.

Image Analysis

The paired FDG-PET/CT images and the planning CT scans were imported into Mirada (Mirada Medical, Build 1.2.0.39, Oxford, UK). All data analysis was carried out and checked by AS and RM. The planning CT scan and CT component of each FDG-PET/CT scan underwent a rigid registration followed by a deformable registration. The quality of both registrations was assessed visually by an experienced clinician and if the registration was judged poor, due to the FDG avidity lying outside the gross tumour, a rectangular region of interest (ROI) was drawn around the volume incorporating surrounding muscles and bones. The registration was repeated using that ROI. As the FDG-PET and CT components of the FDG-PET/CT scan share the same intrinsic frame of reference, this registered each FDG-PET image with the planning CT scan. To create the subvolumes, an elliptical ROI was drawn solely around the area of increased uptake within the anus on each FDG-PET scan and tumour segmentation was carried out as a percentage of the SUV_{max} within the elliptical ROI.

The method used to assess spatial consistency of volumes has been previously published by groups performing similar work in different tumour types [23,24]. On the baseline FDG-PET, tumour segmentation was carried out at 34, 40, 50, 60 and 70% of the SUV_{max} (Pre34%, Pre40%, Pre50%, Pre60% and Pre70%, respectively). On the subsequent scans, subvolumes of 70 and 80% of the SUV_{max} were used (Sub70% and Sub80%, respectively). The overlap fraction between the baseline and the second scan was calculated and expressed as a fraction of the volume of the second scan. Based on previous publications of this method, a mean overlap fraction >0.7 supported the use of a threshold as a target for a biological boost [23,24]. Following this, the planning scan with attached subvolumes was loaded into Eclipse software (version 13.0, Varian Medical Systems, Palo Alto, CA, USA). The centre of mass shifts were used to calculate a vector distance, which was calculated using the following formula:

$$\text{vector distance} = \sqrt{(\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2}.$$

Figure 1 shows the planning scan of patient 14 with two volumes (Pre50% and Sub70%) highlighted. The Dice similarity coefficient (DSC) was calculated for completeness. However, limitations of this method of analysis in this setting are highlighted in the discussion.

Statistics

Overlap fraction, vector distance and DSC between subvolumes of the baseline and subsequent scans were described using medians (interquartile range (IQR)) and means (standard deviation (SD)).

Results

Twenty-one patients underwent two FDG-PET/CT scans between January 2015 and November 2017 at the two time points within the ART study; these images were used for this analysis. Areas of gross tumour of primary disease as well as nodal disease >2 cm in the longest diameter were analysed, resulting in 29 volumes in total.

Patient Characteristics

Table 1 describes the patient and tumour demographics.

Imaging Results

The gross tumour volumes (GTVs), Pre50% and Sub70% volumes in cm^3 are given in Table 2. In terms of overlap fraction, the median overlap fraction of the Pre34%, Pre40% and Pre50% with the Sub70% subvolumes were all above 0.8. The median overlap fraction of the Pre34%, Pre40% and Pre50% subvolumes with the Sub80% subvolumes were all above 0.9. The mean overlap fraction between the Pre34%, Pre40%, Pre50%, Pre60% and Pre70% and the Sub80% fell as the volumes reduced in size; from 0.86 with Pre34%, 0.79 with Pre40%, 0.70 with Pre50%, with Pre60% and Pre70% both below 0.7. The median vector distance values between Pre34%, Pre40% and Pre50% and Sub80% were 0.74, 0.74 and 0.71 mm, respectively. Figure 2 illustrates the median overlap fraction and vector distance for the different volumes. Full details on overlap fractions and vector distances between volumes of high FDG avidity on the pre-treatment scan and the subsequent interim scan are shown in Tables 3 and 4, respectively. As expected, DSC values were poor. Figure 3 illustrates the median DSC for the different volumes assessed. DSC for all the subvolumes per node per patient are available in supplementary material (see Appendix A).

Discussion

FDG uptake within tumours is heterogeneous. The first steps to delivering an FDG-derived biological boost is to



Fig 1. Axial (A) and sagittal (B) slices through a planning computed tomography scan with the Pre50% (black) and the Sub70% (white) volumes illustrated.

Table 1

Patient and tumour demographics

Characteristic	n (%); median (IQR)
Gender	
Male	2 (10%)
Female	19 (90%)
Age (years)	60 (51,70)
T Stage	
T2	16 (76%)
T3	2 (10%)
T4	3 (14%)
N Stage	
N0	10 (48%)
N1	7 (33%)
N3	4 (19%)
M Stage	
M0	21 (100%)

IQR, interquartile range.

determine whether the areas of high avidity are spatially consistent during a course of radiotherapy. We report that in most patients studied, the areas identified at baseline were indeed spatially consistent at fraction 8–9. This introduces the possibility of selectively boosting those areas at highest risk of relapse to improve local control. This has the potential to improve overall survival in this rare tumour type, where dose escalation to the whole tumour is in use and under investigation, but acute and late toxicities may be dose limiting.

We have shown that areas of increased FDG uptake remain reasonably consistent between baseline and in the second week of radiotherapy treatment. Subvolumes of high FDG uptake before and during CRT showed good agreement. Both the mean and medians of the overlap fraction between the diagnostic FDG-PET subvolumes of 34, 40 and 50% of SUV_{max} and subsequent FDG-PET subvolumes of 70 and 80% were all overlap fraction >0.7 . It was notable that agreement between those volumes was less successful in nodal disease compared with primary disease (mean overlap fractions between the diagnostic FDG-PET subvolumes of 34 and 40% and subsequent FDG-PET subvolumes of 80% were 0.95 and 0.65 for nodal volumes but 1.0 for all primary tumours). This may be due to smaller volumes of nodal disease, where a slight inaccuracy in

registration can have a significant effect on the results. Therefore, dose escalation to nodal sites might only be feasible with advanced image guidance or treatment adaptation.

Our approach to assessing volume overlap is based on a publication in NSCLC [23]. Aerts *et al.* [34] recommended a FDG-PET subvolume of 50% of SUV_{max} on the diagnostic FDG-PET/CT as a target for biological boost. This was based on overlap between the diagnostic scans and post-treatment residual disease. Therefore, we acknowledge that our suggested boost volume shows reproducibility rather than site of residual tumour or relapse, as our second scan point is the second week of treatment. A study of pancreatic cancer [24] found that a threshold of 40% of the SUV_{max} on baseline FDG-PET/CT identifies areas of residual metabolic activity seen on a post-CRT FDG-PET/CT and this threshold could aid in the definition of a biological target volume.

Wilson *et al.* [24] and Aerts *et al.* [34], detailed above, proposed that a mean overlap fraction > 0.7 supported the use of a threshold as a target for a biological boost. We have presented the median as the outcomes of interests are normally distributed. However, in order to allow comparison with previous literature we have also reported the mean (standard deviation). Using a mean overlap fraction >0.7 , defined in previous publications as a ‘good’ correlation, the diagnostic FDG-PET subvolumes showed generally satisfactory results. When comparing the FDG-PET subvolumes of 34, 40, 50, 60 and 70% of the SUV_{max} with the subsequent subvolumes of 80%; 24 of 29 (83%), 23 (79%), 20 (69%), 15 (52%) and 11 (38%) volumes achieved that threshold. Therefore, it would be reasonable to consider the diagnostic FDG-PET threshold of 50% of SUV_{max} as a possible target for dose escalation based on pre-treatment imaging, as it achieved a mean overlap fraction of 0.7 used in other papers and was reasonably consistent, with $>70\%$ of cases showing good overlapping agreement. As the diagnostic FDG-PET threshold of 50% represented a mean of only 34% of the GTV, the reduction to volume would be substantial. Therefore, boosting this reduced volume to a higher radiation dose will probably result in reduced normal tissue toxicity than boosting the whole GTV, although this would require confirmation.

Table 2

List of lesions with site, gross tumour volume (GTV) and 50% standardised uptake value (SUV) volume size

	Site of lesion	GTV (cm ³)	Pre50% SUV volume (cm ³)	Sub70% SUV volume (cm ³)
Patient 1	Primary	34.7	5.6	0.7
Patient 2	Primary	29.8	2.7	1.8
Patient 3	Primary	14	5.3	0.5
Patient 4	Primary	69.8	36.9	5.8
Patient 4, Node 1	Right inguinal node	23.1	8.3	3.2
Patient 4, Node 2	Right external iliac node	6.2	3.5	1.2
Patient 4, Node 3	Left inguinal node	7.2	3.4	2.7
Patient 5	Primary	22.2	7.7	3.8
Patient 6	Primary	14	3.4	2
Patient 6, Node 1	Left inguinal node	18.3	5.8	0.3
Patient 6, Node 2	Left internal iliac node	13.3	5.5	2
Patient 6, Node 3	Left external iliac node	13.6	0.2	0.9
Patient 7	Primary	28.7	8.9	3
Patient 8	Primary	13.5	5.3	3.1
Patient 9	Primary	16.9	11.7	1.1
Patient 10	Primary	10.6	2.2	0.2
Patient 10, Node	Right mesorectal node	11.1	1.2	0.1
Patient 11	Primary	14.4	4.1	0.8
Patient 12	Primary	16.4	7	1.4
Patient 13	Primary	4.9	2.3	0.3
Patient 14	Primary	154.2	79.2	3
Patient 15	Primary	109.1	58	29
Patient 15, Node	Right inguinal node	5.8	4.7	1.3
Patient 16	Primary	23.1	8.9	1.7
Patient 17	Primary	32.6	6.9	2.1
Patient 18	Primary	9.7	0.5	0.5
Patient 19	Primary	17.5	4.6	1
Patient 20	Primary	6.4	1	0.7
Patient 21	Primary	63.5	14.4	6.2

Pre50% SUV, volume created from 50% of SUV_{max} on initial positron emission tomography/computed tomography (PET/CT) scan; Sub70% SUV, volume created from 70% of SUV_{max} on PET/CT scan taken at fraction 8 or 9 of radiotherapy.

Our study has some limitations; the fact that the study incorporated small numbers is representative of a rare disease. However, even with small numbers, our study serves to show the feasibility of defining a biological boost in anal cancer and suggests possible boost volumes for further investigation. There are multiple methods of reporting differences between two volumes [35]. We have primarily used the method previously published by Aerts *et al.* [34], as they compared spatial position in volumes

with changing sizes. Our assessment of the volumes using DSC suggests less spatial correlation. However, these contrasting results between our primary methods of comparison and DSC were also seen in the study carried out by Calais *et al.* [19], who highlighted that two volumes with a good superimposition but a large difference in size will yield low Dice and Jaccard indices. DSC values have been calculated for completeness, but the poor DSC correlation should be taken in the context of its limitations when


**Fig 2.** Illustration of median overlap fraction and vector distances between the different volumes assessed.

Table 3

Overlapping fractions between volumes on the pre-treatment scan (Pre34%, Pre40%, Pre50%, Pre60% and Pre70% of the SUV max) and the subsequent interim scans (Sub70% and Sub80% of the SUV max).

Patient	Sub70%					Sub80%				
	Pre34%	Pre40%	Pre50%	Pre60%	Pre70%	Pre34%	Pre40%	Pre50%	Pre60%	Pre70%
Patient 1	1	1	0.86	0.71	0.43	1	1	1	1	1
Patient 2	0.50	0.28	0.11	0.56	0	0.60	0.40	0.20	0	0
Patient 3	1	0.80	0.60	0.40	0.20	1	1	0.50	0.50	0
Patient 4	1	1	1	0.95	0.66	1	1	1	1	1
Patient 4, Node 1	0.66	0.63	0.56	0.47	0.38	0.50	0.40	0.40	0.40	0.40
Patient 4, Node 2	0.33	0.25	0.17	0.83	0	0.33	0	0	0	0
Patient 4, Node 3	0.89	0.81	0.67	0.52	0.33	0.91	0.91	0.73	0.64	0.36
Patient 5	0.97	0.95	0.82	0.63	0.42	1	1	0.95	0.79	0.47
Patient 6	0.95	0.90	0.80	0.60	0.30	1	1	0.90	0.70	0.30
Patient 6, Node 1	1	1	1	1	0.67	1	1	1	1	1
Patient 6, Node 2	1	1	1	0.95	0.75	1	1	1	1	1
Patient 6, Node 3	0.22	0.11	0	0	0	0	0	0	0	0
Patient 7	0.97	0.97	0.93	0.83	0.83	0.92	0.92	0.92	0.92	0.92
Patient 8	0.84	0.77	0.65	0.52	0.35	0.93	0.86	0.71	0.57	0.43
Patient 9	1	1	1	1	0.91	1	1	1	1	1
Patient 10	1	1	1	1	1	1	1	1	1	1
Patient 10, Node	0	0	0	0	0	1	0	0	0	0
Patient 11	1	1	1	1	0.88	1	1	1	1	1
Patient 12	1	1	1	0.93	0.86	1	1	1	0.83	0.83
Patient 13	0	0	0	0	0	0	0	0	0	0
Patient 14	1	1	0.90	0.30	0	1	1	0.93	0.29	0.00
Patient 15	0.98	0.95	0.89	0.80	0.65	1	1	0.94	0.85	0.66
Patient 15, Node	0.92	0.92	0.85	0.77	0.69	1	1	1	1	0.75
Patient 16	1	1	1	0.94	0.94	1	1	1	1	1
Patient 17	0.76	0.71	0.67	0.57	0.38	0.80	0.80	0.70	0.70	0.40
Patient 18	0.60	0.40	0.20	0.20	0	1	1	0	0	0
Patient 19	0.90	0.80	0.60	0.30	0.10	1	0.75	0.75	0.50	0
Patient 20	1	0.86	0.71	0.43	0.14	1	1	0.67	0.33	0.33
Patient 21	0.98	0.97	0.81	0.39	0.07	1	1	0.94	0.50	0.06
Median (IQR)	0.97 (0.76, 1)	0.92 (0.71, 1)	0.81 (0.60, 1)	0.60 (0.40, 0.93)	0.38 (0.07, 0.69)	1.00 (0.92, 1)	1.00 (0.80, 1)	0.92 (0.52, 1)	0.70 (0.33, 1)	0.40 (0, 1)
Mean (SD)	0.81 (0.31)	0.76 (0.33)	0.68 (0.34)	0.61 (0.32)	0.41 (0.35)	0.86 (0.29)	0.79 (0.36)	0.70 (0.38)	0.60 (0.38)	0.48 (0.42)

comparing different-sized lesions. In our study, it was sometimes challenging to separate the tumour FDG activity from the background physiological uptake. All the subvolumes in our study were checked visually and areas of overlap with normal organs were removed. This may also be responsible for the only moderate overlap seen when like-for-like subvolumes were compared (Pre70% versus Sub70%), as such a comparison would leave no margin for small errors in registration and variations in background physiological FDG uptake. Results must be interpreted with caution in view of the potential for errors with deformable registration. The differing field of view in the second FDG-PET/CT scans may raise some concerns. This occurred due to an amendment to the trial protocol in order to incorporate a further tertiary end point investigating the bone marrow toxicity of CRT. However, SUV thresholds are calculated independently of non-pelvic organs using the patients' weight and activity in the primary tumour. Therefore, this will not have affected results. Finally, the subvolumes used within this study are based on previous

literature and we acknowledge that the SUV threshold levels are somewhat arbitrary. The authors have based the study and the methods on best available evidence and we anticipate that further investigation of biological boost volumes in a larger prospective format would be required to validate these methods, to guide patient selection and quantify the reduction in toxicity with smaller boost volumes in comparison with whole GTV.

Previous publications confirmed that higher SUV correlates with worse outcome [16,17,19]. Therefore, the next question that must be addressed is the hypothesis – Does local relapse occur at sites consistent with the high FDG-avid subvolumes identified on diagnostic PET? Unfortunately, as FDG-PET/CT has only recently been incorporated into staging of anal cancer in the UK, and there is a relatively low locoregional relapse rate, there are not sufficient numbers of patients with baseline and subsequent FDG-PET/CT to carry out this analysis. Therefore, we must be clear that the proposed subvolumes are currently based on identifying a reproducible target; further work is

Table 4

Vector Distances between volumes on the pre-treatment scan (Pre34%, Pre40%, Pre50%, Pre60% and Pre70% of the SUV max) and the subsequent interim scans (Sub70% and Sub80% of the SUV max).

Patient	Sub70%					Sub80%				
	Pre34%	Pre40%	Pre50%	Pre60%	Pre70%	Pre34%	Pre40%	Pre50%	Pre60%	Pre70%
Patient 1	0.88	1.04	0.74	0.63	0.55	1.00	0.98	0.86	0.74	0.61
Patient 2	0.96	1.02	1.11	1.14	1.08	0.81	0.85	0.92	0.95	0.90
Patient 3	0.77	0.82	0.86	0.94	0.98	0.85	0.89	0.94	1.03	0.34
Patient 4	0.90	0.34	0.36	0.36	0.38	0.33	0.88	0.90	0.92	0.92
Patient 4, Node 1	0.81	0.73	0.62	0.44	0.42	1.22	1.14	1.02	0.86	0.82
Patient 4, Node 2	0.91	0.93	0.97	0.96	0.96	0.93	0.93	0.95	0.94	0.91
Patient 4, Node 3	0.50	0.53	0.51	0.51	0.55	0.55	0.57	0.55	0.55	0.58
Patient 5	0.33	0.31	0.28	0.30	0.32	0.36	0.34	0.32	0.36	0.39
Patient 6	0.95	0.93	0.89	1.02	1.08	1.00	0.99	0.95	1.08	1.13
Patient 6, Node 1	0.34	0.33	0.31	0.25	0.11	0.47	0.48	0.45	0.39	0.24
Patient 6, Node 2	1.47	1.93	2.03	1.46	1.94	2.94	3.39	3.49	3.51	3.41
Patient 6, Node 3	0.56	0.49	0.38	0.39	0.39	0.67	0.59	0.38	0.31	0.31
Patient 7	0.54	0.53	0.51	0.50	0.55	0.65	0.64	0.61	0.57	0.57
Patient 8	0.51	0.50	0.46	0.41	0.32	0.75	0.74	0.71	0.66	0.56
Patient 9	0.15	0.16	0.20	0.16	0.20	0.19	0.19	0.16	0.21	0.24
Patient 10	0.80	0.76	0.69	0.66	0.68	0.82	0.77	0.71	0.67	0.71
Patient 10, Node	0.31	0.31	0.33	0.33	0.36	0.37	0.37	0.40	0.39	0.42
Patient 11	0.67	0.62	0.60	0.60	0.59	0.63	0.58	0.54	0.51	0.51
Patient 12	1.12	1.16	1.23	1.29	1.39	1.13	1.17	1.24	1.30	1.36
Patient 13	0.55	2.43	2.49	2.85	3.28	2.54	2.51	2.57	2.93	3.36
Patient 14	0.25	0.55	0.56	0.59	0.80	1.57	1.57	1.57	1.60	1.69
Patient 15	0.63	0.23	0.19	0.19	0.21	0.51	0.47	0.42	0.36	0.33
Patient 15, Node	0.25	0.23	0.19	0.19	0.21	0.51	0.47	0.42	0.36	0.33
Patient 16	0.51	0.47	0.42	0.37	0.37	0.57	0.54	0.51	0.47	0.47
Patient 17	0.75	0.75	0.76	0.77	0.77	0.74	0.74	0.74	0.79	0.79
Patient 18	0.65	0.56	0.31	0.41	0.25	0.71	0.63	0.31	0.35	0.31
Patient 19	0.19	0.65	0.72	0.83	0.96	0.62	0.61	0.68	0.78	0.92
Patient 20	0.63	0.21	0.26	0.31	0.36	0.26	0.25	0.24	0.25	0.30
Patient 21	2.43	0.64	0.57	0.60	1.87	0.97	0.98	0.92	0.91	2.17
Median (IQR)	0.65 (0.51, 0.88)	0.62 (0.47, 0.82)	0.60 (0.36, 0.86)	0.59 (0.37, 0.83)	0.55 (0.36, 0.96)	0.74 (0.55, 0.97)	0.74 (0.57, 0.98)	0.71 (0.45, 0.94)	0.74 (0.39, 0.94)	0.61 (0.39, 0.92)
Mean (SD)	0.72 (0.44)	0.71 (0.49)	0.72 (0.51)	0.69 (0.53)	0.78 (0.67)	0.86 (0.60)	0.88 (0.66)	0.86 (0.69)	0.87 (0.73)	0.90 (0.82)




Fig 3. Illustration of the median Dice coefficient of the different volumes assessed.

required to confirm that the proposed subvolumes are appropriate to cover sites of subsequent relapse using scans following treatment. We plan to carry out this analysis initially on the ART cohort in due course and ideally subsequently in a larger number of patients as part of a prospective trial.

Finally, investigation to date regarding the role of functional imaging to predict response is detailed in the introduction. The primary end point of the ART study is in fact based on the hypothesis that we can use functional imaging as an early predictor of response in order to perform treatment intensification, perhaps with a radiotherapy boost, on a selected group identified as poor responders. However, the ability to select patients for dose escalation does not deter from the potential to use an FDG-derived boost to these selected patients as a method of improving local control while limiting toxicity.

In conclusion, on the basis of these findings we conclude that areas of high FDG avidity on diagnostic FDG-PET scans

remain spatially consistent during a course of treatment. A subvolume of 50% of SUV_{max} on the diagnostic scan could be considered as the target for dose escalation. Despite the limitations of this small study it shows the potential of using a biological boost rather than whole GTV dose escalation, which may facilitate improved outcomes while minimising the significant side-effects of whole GTV dose escalation. This study suggests an exciting avenue for further investigation.

Conflicts of interest

None.

Acknowledgements

The authors would like to thank the patients who took part in the ART study and the OUH NHS Foundation Trust. This work was supported by the CRUK EPSRC Cancer Imaging Centre in Oxford, the CRUK Oxford Centre and the NIHR. The trial was sponsored by the University of Oxford and was managed by the Oncology Clinical Trials Office, which is part of the UKCRC-registered Oxford Clinical Trials Research Unit. M. Hawkins is supported by Medical Research Council grant MC_UU_00001/2.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.clon.2018.11.034>.

References

- [1] Glynne-Jones R, Nilsson PJ, Aschele C, Goh V, Peiffert D, Cervantes A, et al. Anal cancer: ESMO-ESSO-ESTRO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2014;25(Suppl. 3):iii10–iii20.
- [2] James RD, Glynne-Jones R, Meadows HM, Cunningham D, Myint AS, Saunders MP, et al. Mitomycin or cisplatin chemoradiation with or without maintenance chemotherapy for treatment of squamous-cell carcinoma of the anus (ACT II): a randomised, phase 3, open-label, 2×2 factorial trial. *Lancet Oncol* 2013;14:516–524.
- [3] Muirhead R, Drinkwater K, O’Cathail SM, Adams R, Glynne-Jones R, Harrison M, et al. Initial results from the Royal College of Radiologists’ UK National Audit of Anal Cancer Radiotherapy 2015. *Clin Oncol* 2017;29:188–197.
- [4] Joseph K, Vos LJ, Warkentin H, Paulson K, Polkosnik LA, Usmani N, et al. Patient reported quality of life after helical IMRT based concurrent chemoradiation of locally advanced anal cancer. *Radiother Oncol* 2016;120:228–233.
- [5] Han K, Cummings BJ, Lindsay P, Skliarenko J, Craig T, Le LW, et al. Prospective evaluation of acute toxicity and quality of life after IMRT and concurrent chemotherapy for anal canal and perianal cancer. *Int J Radiat Oncol Biol Phys* 2014;90:587–594.
- [6] De Francesco I, Thomas K, Wedlake L, Tait D. Intensity-modulated radiotherapy and anal cancer: clinical outcome and late toxicity assessment. *Clin Oncol* 2016;28:604–610.
- [7] Bentzen AG, Guren MG, Wanderas EH, Frykholm G, Tveit KM, Wilsgaard T, et al. Chemoradiotherapy of anal carcinoma: survival and recurrence in an unselected national cohort. *Int J Radiat Oncol Biol Phys* 2012;83:e173–e180.
- [8] Das P, Bhatia S, Eng C, Ajani JA, Skibber JM, Rodriguez-Bigas MA, et al. Predictors and patterns of recurrence after definitive chemoradiation for anal cancer. *Int J Radiat Oncol Biol Phys* 2007;68:794–800.
- [9] Tomaszewski JM, Link E, Leong T, Heriot A, Vazquez M, Chander S, et al. Twenty-five-year experience with radical chemoradiation for anal cancer. *Int J Radiat Oncol Biol Phys* 2012;83:552–558.
- [10] Wright JL, Patil SM, Temple LK, Minsky BD, Saltz LB, Goodman KA. Squamous cell carcinoma of the anal canal: patterns and predictors of failure and implications for intensity-modulated radiation treatment planning. *Int J Radiat Oncol Biol Phys* 2010;78:1064–1072.
- [11] Muirhead R, Partridge M, Hawkins MA. A tumor control probability model for anal squamous cell carcinoma. *Radiother Oncol* 2015;116:192–196.
- [12] Sebag-Montefiore D, Adams R, Bell S, Berkman L, Gilbert DC, Glynne-Jones R, et al. The development of an umbrella trial (PLATO) to address radiation therapy dose questions in the locoregional management of squamous cell carcinoma of the anus. *Int J Radiat Oncol Biol Phys* 2016;96(2S):E164–E165.
- [13] Chen YJ, Suh S, Nelson RA, Liu A, Pezner RD, Wong JY. Setup variations in radiotherapy of anal cancer: advantages of target volume reduction using image-guided radiation treatment. *Int J Radiat Oncol Biol Phys* 2012;84:289–295.
- [14] Cotter SE, Grigsby PW, Siegel BA, Dehdashti F, Malyapa RS, Fleshman JW, et al. FDG-PET/CT in the evaluation of anal carcinoma. *Int J Radiat Oncol Biol Phys* 2006;65:720–725.
- [15] Trautmann TG, Zuger JH. Positron emission tomography for pretreatment staging and posttreatment evaluation in cancer of the anal canal. *Mol Imaging Biol* 2005;7:309–313.
- [16] Hong JC, Cui Y, Patel BN, Rushing CN, Faught A, Eng J, et al. Association of interim FDG-PET imaging during chemoradiation for squamous anal canal carcinoma with recurrence. *Int J Radiat Oncol Biol Phys* 2018;102(4):1046–1051.
- [17] Mohammadkhani Shali S, Schmitt V, Behrendt FF, Winz OH, Heinzel A, Mottaghy FM, et al. Metabolic tumour volume of anal carcinoma on (18)FDG PET/CT before combined radiochemotherapy is the only independent determinant of recurrence free survival. *Eur J Radiol* 2016;85:1390–1394.
- [18] Ajani JA, Winter KA, Gunderson LL, Pedersen J, Benson 3rd AB, Thomas Jr CR, et al. Prognostic factors derived from a prospective database dictate clinical biology of anal cancer: the intergroup trial (RTOG 98-11). *Cancer* 2010;116:4007–4013.
- [19] Kidd EA, Dehdashti F, Siegel BA, Grigsby PW. Anal cancer maximum F-18 fluorodeoxyglucose uptake on positron emission tomography is correlated with prognosis. *Radiother Oncol* 2010;95:288–291.
- [20] Cardenas ML, Spencer CR, Markovina S, DeWees TA, Mazur TR, Weiner AA, et al. Quantitative FDG-PET/CT predicts local recurrence and survival for squamous cell carcinoma of the anus. *Adv Radiat Oncol* 2017;2:281–287.
- [21] Day FL, Link E, Ngan S, Leong T, Moodie K, Lynch C, et al. FDG-PET metabolic response predicts outcomes in anal cancer managed with chemoradiotherapy. *Br J Cancer* 2011;105:498–504.
- [22] Schwarz JK, Siegel BA, Dehdashti F, Myerson RJ, Fleshman JW, Grigsby PW. Tumor response and survival predicted by posttherapy FDG-PET/CT in anal cancer. *Int J Radiat Oncol Biol Phys* 2008;71:180–186.
- [23] Aerts HJ, Bussink J, Oyen WJ, van Elmpt W, Folgering AM, Emans D, et al. Identification of residual metabolic-active

- areas within NSCLC tumours using a pre-radiotherapy FDG-PET-CT scan: a prospective validation. *Lung Cancer* 2012;75:73–76.
- [24] Wilson JM, Mukherjee S, Chu KY, Brunner TB, Partridge M, Hawkins M. Challenges in using (1)(8)F-fluorodeoxyglucose-PET-CT to define a biological radiotherapy boost volume in locally advanced pancreatic cancer. *Radiat Oncol* 2014;9:146.
- [25] Calais J, Thureau S, Dubray B, Modzelewski R, Thiberville L, Gardin I, et al. Areas of high 18F-FDG uptake on pre-radiotherapy PET/CT identify preferential sites of local relapse after chemoradiotherapy for non-small cell lung cancer. *J Nucl Med* 2015;56:196–203.
- [26] Yu W, Cai XW, Liu Q, Zhu ZF, Feng W, Zhang Q, et al. Safety of dose escalation by simultaneous integrated boosting radiation dose within the primary tumor guided by (18)FDG-PET/CT for esophageal cancer. *Radiother Oncol* 2015;114:195–200.
- [27] Even AJ, van der Stoep J, Zegers CM, Reymen B, Troost EG, Lambin P, et al. PET-based dose painting in non-small cell lung cancer: comparing uniform dose escalation with boosting hypoxic and metabolically active sub-volumes. *Radiother Oncol* 2015;116:281–286.
- [28] Shusharina N, Cho J, Sharp GC, Choi NC. Correlation of (18)F-FDG avid volumes on pre-radiation therapy and post-radiation therapy FDG PET scans in recurrent lung cancer. *Int J Radiat Oncol Biol Phys* 2014;89:137–144.
- [29] van den Bogaard J, Janssen MH, Janssens G, Buijsen J, Reniers B, Lambin P, et al. Residual metabolic tumor activity after chemo-radiotherapy is mainly located in initially high FDG uptake areas in rectal cancer. *Radiother Oncol* 2011;99:137–141.
- [30] Berwouts D, Olteanu LA, Duprez F, Vercauteren T, De Gerssem W, De Neve W, et al. Three-phase adaptive dose-painting-by-numbers for head-and-neck cancer: initial results of the phase I clinical trial. *Radiother Oncol* 2013;107:310–316.
- [31] Rasmussen JH, Hakansson K, Vogelius IR, Aznar MC, Fischer BM, Friborg J, et al. Phase I trial of 18F-fluorodeoxyglucose based radiation dose painting with concomitant cisplatin in head and neck cancer. *Radiother Oncol* 2016;120:76–80.
- [32] van Elmpt W, De Ruyscher D, van der Salm A, Lakeman A, van der Stoep J, Emans D, et al. The PET-boost randomised phase II dose-escalation trial in non-small cell lung cancer. *Radiother Oncol* 2012;104:67–71.
- [33] Muirhead R, Adams RA, Gilbert DC, Glynne-Jones R, Harrison M, Sebag-Montefiore D, et al. Anal cancer: developing an intensity-modulated radiotherapy solution for ACT2 fractionation. *Clin Oncol* 2014;26:720–721.
- [34] Aerts HJ, van Baardwijk AA, Petit SF, Offermann C, Loon J, Houben R, et al. Identification of residual metabolic-active areas within individual NSCLC tumours using a pre-radiotherapy (18)Fluorodeoxyglucose-PET-CT scan. *Radiother Oncol* 2009;91:386–392.
- [35] Hanna GG, Hounsell AR, O'Sullivan JM. Geometrical analysis of radiotherapy target volume delineation: a systematic review of reported comparison methods. *Clin Oncol* 2010;22:515–525.