

First record of a ‘fish’ blood fluke (Digenea: Aporocotylidae) from a marine mammal: *Cardicola dhangali* n. sp.

Kate Suzanne Hutson^{a,b,*}, David Brendan Vaughan^a, David Blair^a

^a Marine Parasitology Laboratory, College of Science and Engineering, Centre for Sustainable Fisheries and Aquaculture, James Cook University, Douglas, Townsville, 4811, Australia

^b Cawthron Institute, 98 Halifax Street East, Nelson, 7010, New Zealand

ARTICLE INFO

Keywords:

Aporocotylidae
Schistosomidae
Spirorchiidae
Blood fluke
Cardicola
Dugong

ABSTRACT

We describe the first known blood fluke from a marine mammal, the dugong, *Dugong dugon* (Sirenia: Dugongidae), which represents a new species of aporocotylid, *Cardicola dhangali* n. sp. (Digenea: Aporocotylidae). Eggs presumed to be of blood flukes have been previously reported from dugongs. This exciting discovery raises questions regarding evolution and host-switching in the Aporocotylidae, which prior to this study were only known to infect actinopterygian and chondrichthyan fishes. The new species has male and female genital pores opening on the right side of the body, with the male genital pore opening posterior to the entire reproductive system and the testis is extra-caecal. The uterus is highly convoluted, and the ovary is irregularly lobate. These features, together with the size and number of the tegumental spines per row, easily distinguish the new species from the most similar congeners *Cardicola aurata* Holzer et al., 2008, *Cardicola chaetodontis* Yamaguti, 1970, *Cardicola currani* Bullard and Overstreet, 2004, *Cardicola forsteri* Cribb et al., 2000, *C. jiunguru* Yong et al., 2016, and *Cardicola palmeri* Bullard and Overstreet, 2004, all of which infect actinopterygian fishes. Given that *Cardicola* is the most diverse and least host-specific of the marine aporocotylid genera, it seems credible that a successful host-switch has occurred from an actinopterygian to *D. dugon*. Further sampling of sirenians and other marine mammals is warranted to gain a more comprehensive understanding of the evolutionary biology and biodiversity of the blood flukes (superfamily Schistosomatoidea Stiles and Hassall, 1898), but presents a substantial challenge with respect to their conservation status and large size.

1. Introduction

Blood flukes (Platyhelminthes: Digenea: Schistosomatoidea) are aquatic parasites that infect the cardiovascular system of their definitive host. Historically, blood flukes have been assigned to three families corresponding to the vertebrate definitive host lineages they infect; the Schistosomatidae, which infect birds and terrestrial mammals, the Spirorchiidae, which infect reptiles and the Aporocotylidae, which infect fishes. The schistosomatids are dioecious while the aporocotylids and paraphyletic spirorchiids (Snyder, 2004; Oréllis-Ribeiro et al., 2014) are hermaphrodites (with the exception of the dioecious ‘spirorchiid’ infecting crocodiles, *Griffithsillharzia amoena* Platt et al., 1991; see Brant and Loker, 2005; Platt et al., 1991). Blood flukes exhibit a two-host life cycle with asexual reproduction occurring in an invertebrate host and direct penetration of the skin of the definitive host. Schistosomes and most spirorchiids utilize gastropods as intermediate hosts (e.g. Brant

et al., 2006; Cribb et al., 2017a; see de Buron et al., 2018 for evidence of spirorchiids using a polychaete intermediate host), while aporocotylids of actinopterygians exploit freshwater gastropods (e.g. Evans and Heckmann, 1973; Schell, 1974; Kirk and Lewis, 1993) and marine terebellid polychaetes (e.g. Køie, 1982; Cribb et al., 2011; Sugihara et al., 2014; Shirakashi et al., 2012). Recently, Cribb et al. (2017b) provided compelling evidence that bivalves serve as distinct intermediate hosts for aporocotylids of chondrichthyan fishes. The rate of proposal of new genera and species of Aporocotylidae suggests that a potentially high diversity remains to be discovered in this family (Cribb and Bray, 2011).

To date, blood flukes have not been recorded from marine mammals. The first indication that marine mammals may be susceptible was reported by Marsh et al. (1984) who documented trematode parasite eggs in ovary sections of the dugong, *Dugong dugon* (Müller, 1776; Sirenia: Dugongidae). These eggs exhibited ‘distinct black paired spots’

* Corresponding author. Marine Parasitology Laboratory, College of Science and Engineering, Centre for Sustainable Fisheries and Aquaculture, James Cook University, Douglas, Townsville, 4811, Australia.

E-mail address: kate.hutson@cawthron.org.nz (K.S. Hutson).

<https://doi.org/10.1016/j.ijppaw.2019.06.009>

Received 4 April 2019; Received in revised form 18 June 2019; Accepted 19 June 2019

2213-2244/ © 2019 Published by Elsevier Ltd on behalf of Australian Society for Parasitology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

which is consistent with the description of dark bodies within developing *Cardicola* miracidia (McVay et al., 2011). Later, one of the authors (DB) recovered a single aporocotyloid blood fluke of suitable quality for description from a stranded dugong in Townsville, Queensland. Subsequent efforts by the authors to recover additional specimens were unsuccessful due to a combination of the rarity of opportunities to necropsy fresh dugongs and the rapid decomposition of the host and soft-bodied parasites in tropical environments (Eros et al., 2007). Indeed, opportunities to conduct dugong necropsies are few because they are protected in Australia under the Environment Protection and Biodiversity Act (1999).

Herein we describe the new species of aporocotyloid discovered by DB from *Dugong dugon*. This is timely given current global research momentum in documenting species richness within the Aporocotylidae. This novel species represents an important discovery that warrants description despite the absence of molecular data. Future material and molecular analyses could permit new inferences into the potential for evolutionary expansion of the Aporocotylidae into marine mammals.

2. Materials and methods

A single male dugong, *Dugong dugon* (273 cm total length; tail width 90 cm; collectors' ID MM313) originating from the Strand, Townsville, Queensland (19°15'00.0"S 146°49'59.9"E), in post-mortem condition D2 ("carcass in good condition, fresh/edible"; Eros et al., 2007) was dissected by DB on the 8th September 1992. A single aporocotyloid was recovered from heart washings. The specimen was not apparently alive but was in good condition. It was fixed beneath a coverslip, with a small dab of Vaseline® jelly under each corner (to prevent excess flattening) while 10% formalin was drawn under the coverslip using absorbent paper, prior to being stained with Gower's carmine, and mounted in Canada balsam on a glass slide under a cover slip. A fragment of a second specimen was noticed in heart washings but was lost. Examination of the heart, a common site for spirorchiid and aporocotyloid blood flukes, had been prompted by the report in Marsh et al. (1984) of probable blood-fluke eggs in the ovaries of several dugongs. Animal ethics statement: note that the animal was dead and found washed up on a beach and therefore did not require specific animal ethics approval.

The parasite specimen was observed using an Olympus BX53 compound light microscope fitted with direct interference contrast optics, or a Leica DM LS2 compound light microscope fitted with phase-contrast optics. Drawings were made using a drawing tube connected to the Leica DM LS2. All measurements are given in microns. Measurements were taken using Olympus LabSens® image analysis software. Where specifically indicated, measurements of curved structures following the curvature were taken using the polyline function of the software. In the interest of accessibility to the scientific community, images through the focal plane of this rare specimen were prepared and made available as supplementary video files: Hutson, K. (2019). Z-stack videos files for: First record of a 'fish' blood fluke (Digenea: Aporocotylidae) from a marine mammal: *Cardicola dhangali* n. sp. James Cook University. (dataset). <https://doi.org/10.25903/5c527ca5c8b6d> Digital Object Identifier (DOI):10.25903/5c527ca5c8b6d.

The following specimens were viewed for comparative purposes: *Cardicola brasiliensis* Knoff and Amato, 1992 (University of Nebraska State Museum 31717, 31718 paratypes; two slides), *Cardicola whitteni* Manter, 1954 (KSH personal collection, voucher, one slide), *Paradeontacylix sanguinicoides* McIntosh, 1934 (South Australian Museum Australian Helminth Collection [SAMA AHC] vouchers AHC 28909, 28910; two slides), *Paradeontacylix* sp. (SAMA AHC vouchers AHC 28911, 28912; two slides).

3. Results

Family Aporocotylidae Odhner, 1912




Fig. 1. *Cardicola dhangali* n. sp. whole mount ventral view. Abbreviations: ac, anterior caecal branch; alp, antero-lateral projection of posterior caecal branch; fgp, female genital pore; me, Mehlis' glands; o, ovary; oes, oesophagus; oöt, oötype; pc, posterior caecal branch; sv, seminal vesicle; t, testicular field; ut, uterus; vd, vas deferens; vitf, vitelline follicle. Scale bar = 1000 µm.

Genus *Cardicola* Short, 1953

Cardicola dhangali n. sp.

3.1. Description (Figs. 1–3)

[Based on single whole-mounted holotype specimen] Body lanceolate (Fig. 1), 3690 in total length; 1105 wide at widest point; 3.3 times longer than wide. Between 2 (anterior-most), and 20 lanceolate




Fig. 2. *Cardicola dhangali* n. sp. anterior region detail. Abbreviations: dnc, dorsal nerve commissure; m, mouth; ts, tegumental spines. Other abbreviations as for Fig. 1. Scale bar = 100 µm.

tegumental spines (Fig. 2) 7–8 long arranged marginally in ventro-lateral rows initially 7–8 apart anteriorly, increasing to 11–14 apart at body mid region, decreasing to 9–10 posteriorly. Tegumental bristles or papillae not observed. Dorsal nerve commissure 18 thick, 186 from anterior extremity. No discernible oral sucker present. Mouth small, aperture 7 in diameter, immediately subterminal, ventral, slightly raised from body surface and weakly domed (Fig. 2). Oesophagus 1071 long, occupying ~29% of body length (Fig. 1). Caecum roughly H-shaped (Fig. 1). Anterior caecal branches slightly curved, terminally lobate (Fig. 1). Right anterior branch 359 long, left anterior branch 379 long, following curvature. Anterior caecal branches 796 and 828 from anterior body extremity, respectively. Posterior caecal branches sigmoidally convoluted (Fig. 1), each initially forming short antero-laterally directed protuberance post-bifurcation (Fig. 1). Right protuberance 59 long, left protuberance 55 long. Right posterior caecal branch 3055 long, left caecal branch 2757 long (following curvature), terminating non-confluently dorsal to ovary (Fig. 1). Caecal extent occupies ~58% of body length.

Testis (Figs. 1), 2388 long, 674 wide, not easily discernible, extra-caecal, extending anteriorly, medially beyond anterior caecal branches, but not reaching dorsal nerve commissure; extending posteriorly to level of ovary and immediately anterior to uterus. Vas deferens dorsal to ovary, arising medially from posterior portion of testis, initially broad, narrowing distally and turning right, ventrally over proximal uterine fold (Fig. 1), then posteriorly to meet curved seminal vesicle on right side of body, posterior to remainder of reproductive system (Figs. 1 and 3). Seminal vesicle curved to right, 194 long (following curvature), 50 wide. Short, terminally folded ejaculatory duct present (Figs. 1 and 3). Male genital pore difficult to discern, opens dorsally on right of body, posterior to remainder of reproductive system in immediate vicinity of short ejaculatory duct.

Ovary distinct, irregularly lobate, 327 long, 504 wide, positioned medially and left in posterior body portion (Figs. 1 and 3). Oviduct exits

ovary posteriorly, left of body mid-line, travelling posteriorly before opening into empty oviductal chamber 464 long, 147 wide. Duct leading from oviductal chamber curves right to enter oötype (Figs. 1 and 3). Mehli's gland present (Figs. 1 and 3). Vitelline duct not observed. Proximal portion of uterus broad, convoluted, travels anteriorly, right of vas deferens to right posterior portion of ovary, narrows slightly, becoming increasingly convoluted, extending right, curving posteriorly to distal portion opening dorsally in prominent, muscular female genital pore 256 from right body margin, in line with oötype, well anterior to male genital pore, 487 from posterior body extremity (Figs. 1 and 3). Vitelline follicles conspicuous along lateral outer margin of lateral nerve to level of ovary on left of body, and anterior portion of uterus on right. Vitelline follicles also present in anterior portion, anterior to dorsal nerve commissure. Excretory vesicle not observed.

Type host: *Dugong dugon* (Müller, 1776).

Host details: adult male, unknown age, D2-condition, 273 cm; tail width 90 cm (specimen number MM313).

Site in host: heart washings.

Type locality: The Strand, Townsville, Queensland, Australia (19.25 S, 146.8333 E).

Type material: holotype (Queensland Museum G237841).

ZooBank registration: To comply with the regulations set out in article 8.5 of the amended 2012 version of the International Code of Zoological Nomenclature (ICZN, 1999), details of the new species have been submitted to ZooBank. The Life Science Identifier (LSID). The LSID for *Cardicola dhangali* n. sp. is <http://zoobank.org/urn:lsid:zoobank.org:pub:FC2C3CA7-503A-4846-AA28-D56E9B00B7B>.

Etymology: the new species name is derived from the traditional language (Kala Lagaw Ya) of the Western and Central islands of the Torres Strait for dugong: 'Dhangal'.

3.2. Remarks

Cardicola dhangali n. sp. differs from all other known *Cardicola* species by its parasitism of the only known mammalian host for the genus, *Dugong dugon*, its highly convoluted uterus (Figs. 1 and 3), and the possession of antero-lateral projections on the posterior caecal branches immediately post-bifurcation (Fig. 1). It is similar in morphological aspects to *Cardicola chaetodontis* Yamaguti (1970) found in the gills and heart of butterflyfishes, *Cardicola aurata* Holzer et al. (2008) from *Sparus aurata*, *Cardicola currani* Bullard and Overstreet (2004) from *Sciaenops ocellatus*, *Cardicola forsteri* Cribb et al. (2000), *Cardicola jiugurru* Yong et al. (2016) from *Chanos chanos*, and *Cardicola palmeri* Bullard and Overstreet (2004) from *Pogonias cromis*. The majority of *Cardicola* species possess female and male genital pores positioned to the left of the body mid-line. The genital pores of *C. chaetodontis* (sensu Yamaguti, 1970) and the new species are positioned to the right of the body mid-line. Of the male and female genital pores in *Cardicola* species, the male genital pore alone of *C. dhangali* n. sp., *C. aurata*, *C. currani*, *C. forsteri*, *C. jiugurru*, *C. palmeri* opens posterior to the remainder of the reproductive system (see Cribb et al., 2000; Bullard and Overstreet, 2004; Holzer et al., 2008; Yong et al., 2016).

Cardicola dhangali n. sp. can be distinguished from *C. chaetodontis* (Yamaguti, 1970; Nolan and Cribb, 2006) by the position of the male genital pore (not posterior to the reproductive system in the latter), the morphology and position of the ovary, which is bi-lobed and medial in *C. chaetodontis*, but irregularly lobate and positioned medial to left of the body-mid line in the new species, and the extent and pathway of the posterior caecal branches. The posterior caecal branches terminate overlapping with the ovary of *Cardicola dhangali* n. sp. (Fig. 1), yet terminate short of the ovary in *C. chaetodontis* (see Yamaguti, 1970; Nolan and Cribb, 2006). The posterior caecal branches are characteristically sigmoidally convoluted in *Cardicola dhangali* n. sp. (Fig. 1), yet relatively straight in *C. chaetodontis* (see Yamaguti, 1970; Nolan and Cribb, 2006).

Cardicola dhangali n. sp. differs from *C. aurata*, *C. currani*, *C. forsteri*,




Fig. 3. *Cardicola dhangali* n. sp. reproductive system (excluding testis). Abbreviations: du; distal portion of uterus; ed, ejaculatory duct; oc, oviductal chamber; od, oviduct; pu, proximal portion of uterus. Other abbreviations as for Fig. 1. Scale bar = 150 µm.

Table 1
Comparison between *Cardicola dhangali* n. sp. and its most morphologically similar congeners; percentages calculated from total body length, following (in part) Nolan et al. (2014). Measurements shown in μm .

Species	Body	Body length/width	Spine length	Spines per row	Oral sucker	Oesophagus %	Anterior caeca length %	Posterior caeca/ anterior caeca	Testis length/width	Testis length %	Testis width %	Ovary length %	Ovary position	Male genital pores position	Female genital pore position
<i>C. dhangali</i> n. sp.	3690 $\times$ 1105	3.3	7–8	7–14	Absent	29.02	9.73, 10.27	6.79, 8.50	3.54	64.71	18.26	8.86	Medial and left	Right	Right
<i>C. aurata</i> ^a	1093–1321 $\times$ 248–284	3.8–4.9	4–5	4–11	Present	–	–	3.10–3.60	–	–	–	–	–	Left-medially	Left
^b <i>C. chaetodontis</i> ^b	1150–1850 $\times$ 180–200	–	5–7	–	Absent	–	–	–	–	–	–	–	–	Right	Right
^b <i>C. chaetodontis</i> ^c	899–1396 $\times$ 119–257	4.3–7.0	–	5–6	Present	33–37	2–5	5.4–12.5	3.1–5.5	36–38	41–66	7–16	Right to medial	Left	Left
<i>C. currani</i> ^d	1375–2853 $\times$ 442–663	2.6–5.0	19–25	2–4	Absent	31–40	9–18	1.8–4.2	–	24–37	25–59	–	Medial	Left	Left
<i>C. forsteri</i> ^{e, f}	2512–4670 $\times$ 570–1070 ⁽ⁱ⁾	4.34–5.79 ^(e)	–	–	Present	29–33 ⁽ⁱ⁾	–	4–7 ⁽ⁱ⁾	–	72–75 ⁽ⁱ⁾	–	–	–	Left	Left
<i>C. jiugurru</i> ^g	1785–2505 $\times$ 132–160	13.2–15.7	< 1	–	Present	28.2–36.3	15.9–22.0	2.2–4.1	–	–	–	–	–	Left	Left
<i>C. palmeri</i> ^d	1449–2357 $\times$ 867–1105	1.9–2.9	33–38	3–5	Absent	44–52	19–26	1.0–1.8	–	14–21	42–61	–	Medial	Left	Left

^aHolzer et al. (2008); ^bYamaguti (1970); ^cNolan and Overstreet (2004); ^dShirakashi et al. (2012); ^eNolan et al. (2014); ^fYong et al. (2016); ^g*Cardicola chaetodontis* considered here as two potentially separate species based on the morphological differences in Yamaguti (1970) and Nolan and Cribb, (2006).

C. jiugurru, and *C. palmeri* in the position of the male genital pore on the right side of the body. In addition, *Cardicola dhangali* n. sp. differs from *C. aurata* in the size and shape of the seminal vesicle, which is much larger in the latter, the morphology of the ejaculatory duct, which is armed in *C. aurata*, and there are fewer tegumental spines per row (4–11, and 11–14, respectively) in *C. aurata* and *C. forsteri*. *Cardicola dhangali* n. sp. differs further from *C. currani*, *C. jiugurru*, and *C. palmeri* by the size, or size and shape of its tegumental spines. These are shortest in *C. jiugurru* (< 1 μm long); much larger and re-curved in *C. currani* and *C. palmeri*. These three species are also further differentiated from *Cardicola dhangali* n. sp. by the position or morphology of the ovary, which is medial in *C. currani* and *C. palmeri*, rather than medial to left of the body in the new species, and is clearly bi-lobed in *C. jiugurru*. The testis of *C. aurata*, *C. chaetodontis*, *C. currani*, and *C. jiugurru* is inter-caecal, whereas the testis of the new species extends extra-caecally. For additional morphological comparisons see Table 1.

4. Discussion

Infection of dugong (*Dugong dugon*) with an aporocotyloid blood fluke likely represents a host-switch event from an actinopterygian fish. While this new host record presents an exciting discovery, it is less surprising that the new species was morphologically attributed to *Cardicola*, because it is the richest and least host-specific of the marine aporocotyloid genera. *Cardicola* is unusual compared to most other aporocotyloid genera in that it exhibits lower host-specificity; thirty-five *Cardicola* species have been described from 14 actinopterygian fish families, in contrast to its sister taxon, *Paradeontacylix* (see Holzer et al., 2008; Yong et al., 2016, 2018), of which eight (of the nine described) species only infect carangid fishes (e.g. Hutson and Whittington, 2006; Repullés-Albelda et al., 2008; WoRMS Editorial Board, 2018). *Cardicola* continues to exhibit a remarkable rate of discovery, with 24 new species attributed to this genus since 2000 (Yong et al., 2018; WoRMS Editorial Board, 2018). This is likely a consequence of multiple factors, including the circulatory system historically being overlooked in routine fish dissections (Cribb and Bray, 2011), increased dedicated taxonomic research effort in the northern (e.g., Bullard and Overstreet, 2004; Bullard et al., 2012; Bullard, 2010, 2013) and southern hemispheres (e.g., Nolan and Cribb, 2006, 2014; Yong et al., 2016, 2018), and importance in aquaculture (e.g., Cribb et al., 2000; Ogawa et al., 2010). Recently, Nolan et al. (2014) predicted that continued discovery of new *Cardicola* species would likely reveal a series of radiations in association with particular fish taxa, as well as evidence of host-switching.

Sirenian ancestry is remote from that of cetaceans or pinnipeds and sirenians re-evolved an aquatic lifestyle independently of, although simultaneously with, cetaceans (Domning, 2009). Thus, it is unclear whether dedicated parasitological necropsies will reveal more aporocotyloid richness in non-sirenian marine mammal groups. Indeed, unlike most other taxa of marine mammals, modern sirenians, and especially dugongs, inhabit warm, shallow waters where many aporocotyloid species occur in fish hosts. Given this, host-switching of aporocotyloids into marine mammals is more likely to involve sirenians than cetaceans or pinnipeds. Alternatively, given the low host specificity of *Cardicola* and the shared habitat with dugongs, it is plausible that this discovery may be a case of accidental infection. Parasitological surveys of a broad range of marine mammalian taxa are probably not feasible because traditional approaches to study parasites are impractical for large marine mammals, most of which are protected (Hermosilla et al., 2015). Thus, rare opportunities to acquire quality morphological and molecular material, such as those from fresh strandings, must be seized to enable the elucidation of the likelihood and frequency of host-switching events and co-evolution in the Aporocotyloidea.

The location of blood fluke eggs in the definitive host influences transmission and pathogenesis and could potentially assist with screening marine mammals for blood flukes. Successful parasite transmission requires eggs or miracidia to escape from the closed circulatory

system. Some venous-dwelling schistosome eggs concentrate in the nasal circulation of birds and terrestrial mammals, which are presumably released when the host submerges its head during drinking, feeding or diving (Platt and Brooks, 1997). Adults of *Schistosoma mansoni* Sambon, 1907 produce eggs that pass into the intestinal lumen for release into the environment, while many eggs are carried to the liver, where they become trapped, often causing severe pathology (Pearce and MacDonald, 2002). Disease manifestation from egg accumulation has led to the hypothesis that hosts that succumb to infection in aquatic environments could release eggs via predation, scavengers, or decomposition (Platt and Brooks, 1997). Indeed, the arterial-dwelling spirorchiids release eggs in the direction of blood flow, which results in the wide dissemination of eggs within the host (Platt and Brooks, 1997). Aporocotylid miracidia are believed to primarily escape to the external environment through the gills, while a few species' eggs probably traverse the gut or depend on the death of the host for release (Lester et al., 2009; Bray et al., 2012). There are numerous reports of aporocotylid eggs trapped in the heart of fish where they can become encapsulated and die (Overstreet and Thulin, 1989; Ogawa et al., 1989; Bullard and Overstreet, 2002, 2008; Lester et al., 2009; Yong et al., 2013) and endocarditis has been associated with infections in aquaculture (Warren et al., 2017). Plausible routes to the external environment for aporocotylids of dugongs could include the lungs, nostrils and excretory or reproductive systems. Aporocotylid eggs observed trapped in dugong ovaries (Marsh et al., 1984) indicate that host death may also be a feasible mechanism for release of eggs and transmission to susceptible intermediate hosts. If blood flukes utilize mammalian excretory systems for release of eggs like their schistosome relatives, non-invasive detection could be feasible from collection of fecal deposits and the application of environmental DNA techniques.

5. Conclusions

Infection of dugongs with the aporocotylid blood fluke, *Cardicola dhangali* n. sp., most likely represents a host-switch event from an actinopterygian fish. Given current research interest in aporocotylid discovery and the evolutionary history of the Schistosomatoidea, it is of intrinsic value to present this description of *Cardicola dhangali* n. sp. so that researchers can maximize future sampling opportunities of sirinians and other marine mammals that might act as hosts.

Conflicts of interest

All authors declare no conflicts of interest.

Acknowledgements

We thank RA Bray and GA Boxshall at the Natural History Museum for discussion and technical support. This research was supported through a James Cook University Special Studies Program (sabbatical) awarded to KSH. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jppaw.2019.06.009>.

References

Bray, R.A., Cribb, T.H., Littlewood, D.T.J., 2012. *Sasala nolani* gen. n., sp. n. (Digenea: Aporocotylidae) from the body-cavity of the guineafowl puffer fish *Arothron meleagris* (Lacepede) (Tetraodontiformes: tetraodontidae) from off moorea. Fr. Polyn. Zootaxa 3334, 29–41.

Brant, S.V., Loker, E.S., 2005. Can specialized pathogens colonize distantly related hosts? Schistosome evolution as a case study. PLoS Pathog. 1 (3), e38. <https://doi.org/10.1371/journal.ppat.0010038>.

Brant, S.V., Morgan, J.A.T., Mkoji, G.M., Snyder, S.D., Rajapakse, R.P.V.J., Eric, S., Loker, E.S., 2006. An approach to revealing blood fluke life cycles, taxonomy, and diversity: provision of key reference data including dna sequence from single life cycle stages. J. Parasitol. 92, 77–88.

Bullard, S.A., 2010. A new species of *Cardicola* Short, 1953 (Digenea: Aporocotylidae) from the heart and branchial vessels of two surfperches (Perciformes: embiotocidae) in the eastern Pacific Ocean off California. Am. Soc. Parasitol. 96, 382–388.

Bullard, S.A., 2013. *Cardicola langeli* sp. n. (Digenea: Aporocotylidae) from heart of sheephead, *Archosargus probatocephalus* (Actinopterygii: sparidae) in the Gulf of Mexico, with an updated list of hosts, infection sites and localities for *Cardicola* spp. Folia Parasitol. 60, 17–27.

Bullard, S.A., Baker, T., de Buron, I., 2012. New species of *Cardicola* (Digenea: Aporocotylidae) from heart of atlantic croaker, *Micropogonias undulatus* (Perciformes: Sciaenidae), of the south atlantic light. J. Parasitol. 98, 328–332.

Bullard, S.A., Overstreet, R.M., 2002. Potential pathological effects of blood flukes (Digenea: sanguinicolidae) on pen-reared marine fishes. Proc. Gulf Caribb. Fish. Inst. 52, 10–25.

Bullard, S.A., Overstreet, R.M., 2004. Two new species of *Cardicola* (Digenea: sanguinicolidae) in drums (sciaenidae) from Mississippi and Louisiana. J. Parasitol. 90, 128–136.

Bullard, S.A., Overstreet, R.M., 2008. Digeneans as enemies of fishes. In: In: Eiras, J.C., Segner, H., Wahli, T., Kapoor, B.G. (Eds.), Fish Diseases 2. Science Publishers, Enfield, pp. 817–976.

Cribb, T.H., Adlard, R.D., Hayward, C.J., Bott, N.J., Ellis, D., Evans, D., Nowak, B.F., 2011. The life cycle of *Cardicola forsteri* (Trematoda: Aporocotylidae), a pathogen of ranches southern bluefin tuna, *Thunnus maccoyi*. Int. J. Parasitol. 41, 861–870.

Cribb, T.H., Bray, R.A., 2011. Trematode families and genera: have we found them all? Trends Parasitol. 27, 149–154.

Cribb, T.H., Chick, R.C., O'Connor, W., O'Connor, S., Johnson, D., Sewell, K.B., Cutmore, S.C., 2017b. Evidence that blood flukes (Trematoda: Aporocotylidae) of chondrichthyan infect bivalves as intermediate hosts: indications of an ancient diversification of the Schistosomatoidea. Int. J. Parasitol. 47, 885–891.

Cribb, T.H., Crespo-Picazo, J.L., Cutmore, S.C., Stacy, B.A., Chapman, P.A., García-Párraga, D., 2017a. Elucidation of the first definitively identified life cycle for a marine turtle blood fluke (Trematoda: Spirorchiidae) enables informed control. Int. J. Parasitol. 47, 61–67.

Cribb, T.H., Daintith, M., Munday, B., 2000. A new blood-fluke, *Cardicola forsteri*, (Digenea: sanguinicolidae) of southern blue-fin tuna (*Thunnus maccoyi*) in aquaculture. T. Roy. Soc. South. Aust. 124, 117–120.

de Buron, I., Colon, B.L., Siegel, S.V., Oberstaller, J., Rivero, A., Kyle, D.E., 2018. First evidence of polychaete intermediate hosts for *Neosporichis* spp. marine turtle blood flukes (Trematoda: spirorchiidae). Int. J. Parasitol. <https://doi.org/10.1016/j.ijpara.2018.08.002>.

Domning, D.P., 2009. Sirenian evolution. In: Würsig, B., Perrin, W., Würsig, B., Thewissen, J.G.M. (Eds.), Encyclopedia of Marine Mammals, second ed. Academic Press, pp. 1016–1019.

Eros, C., Marsh, H., Bonde, R., O'Shea, T., Beck, C., Recchia, C., Dobbs, K., Turner, M., Lemm, S., Pears, R., Bowater, R., 2007. Procedures for the salvage and necropsy of the dugong (Dugong dugon). In: Townsville: Great Barrier Reef Marine Park Authority, second ed. Research Publication No. 85.

Evans, W.A., Heckmann, R.A., 1973. The life history of *Sanguinicola Klamathensis* life. Science 13, 1285–1291.

Hermosilla, C., Silva, L.M.R., Prieto, R., Kleinertz, S., Taubert, A., Silva, M.A., 2015. Endo- and ectoparasites of large whales (Cetartiodactyla: balenopteridae, Physeteridae): overcoming difficulties in obtaining appropriate samples by non- and minimally-invasive methods. Int. J. Parasitol. Parasites Wildl. 4, 414–420.

Holzer, A.S., Montero, F.E., Repullés, A., Nolan, M.J., Sitja-Bobadilla, A., Alvarez-Pellitero, P., Zarza, C., Raga, J.A., 2008. *Cardicola aurata* sp. n. (Digenea: sanguinicolidae) from mediterranean *Sparus aurata* L. (Teleostei: sparidae) and its unexpected phylogenetic relationship with *Paradeontacylix* McIntosh, 1934. Parasitol. Int. 57, 472–482.

Hutson, K.S., Whittington, I.D., 2006. *Paradeontacylix godfreyi* n. sp. (Digenea: sanguinicolidae) from the heart of wild *Seriola lalandi* (Perciformes: carangidae) in southern Australia. Zootaxa 1151, 55–68.

ICZN International Code of Zoological Nomenclature, 1999. The International Trust for Zoological Nomenclature, London, UK, Fourth Edition. 306 pp.

Kirk, R.S., Lewis, J.W., 1993. The life-cycle and morphology of *Sanguinicola inermis* plehn, 1905 (Digenea: sanguinicolidae). Syst. Parasitol. 25, 125–133.

Koie, M., 1982. The redia, cercaria and early stages of *Aporocotyle simplex* Odhner, 1900 (Sanguinicolidae) – a digenetic trematode which has a polychaete annelid as the only intermediate host. Ophelia 21, 115–145.

Lester, R.J.G., Rawlinson, S.E., Weaver, L.C., 2009. Movement of sea mullet *Mugil cephalus* as indicated by a parasite. Fish. Res. 96, 129–132.

Marsh, H., Heinsohn, G.E., Channells, P.W., 1984. Changes in the ovaries and uterus of the dugong, *Dugong dugon* (Sirenia: Dugongidae), with age and reproductive activity. Aust. J. Zool. 32, 743–766.

McVay, M.J., Bakenhaster, M., Bullard, S.A., 2011. *Cardicola laruei* Short, 1953 (Digenea: Aporocotylidae) from heart of seatrout, *Cynoscion* spp., (Perciformes: sciaenidae) in the Gulf of Mexico and Atlantic Ocean: taxonomic redescription, first observations of egg and miracidium, and comments on geographic distribution and host specificity. Comp. Parasitol. 78, 291–305.

Nolan, M.J., Cribb, T.H., 2006. *Cardicola* Short, 1953 and *Braya* n. gen. (Digenea: sanguinicolidae) from five families of tropical Indo-Pacific fishes. Zootaxa 1265, 1–80.

Nolan, M.J., Miller, T.L., Cutmore, S.C., Cantacessi, C., Cribb, T.H., 2014. *Cardicola beveridgei* n. sp. (Digenea: Aporocotylidae) from the mangrove jack, *Lutjanus argentimaculatus* (perciformes: lutjanidae), and *C. bullardi* n. sp. from the Australian spotted

- mackerel, *Scomberomorus munroi* (perciformes: scombridae), from the northern great barrier reef. *Parasitol. Int.* 63, 735–745.
- Ogawa, K., Hattori, K., Hatai, K., Kubota, S., 1989. Histopathology of cultured marine fish, *Seriola purpurascens* (Carangidae) infected with *Paradeontacylix* spp. (Trematoda: sanguinicolidae) in its vascular system. *Fish Pathol.* 24, 75–81.
- Ogawa, K., Tanaka, S., Sugihara, Y., Takami, I., 2010. A new blood fluke of the genus *Cardicola* (Trematoda: sanguinicolidae) from Pacific bluefin tuna *Thunnus orientalis* (Temminck and Schlegel, 1844) cultured in Japan. *Parasitol. Int.* 59, 44–48.
- Oréllis-Ribeiro, R., Arias, C.R., Halanych, K.M., Cribb, T.H., Bullard, S.A., 2014. Diversity and ancestry of flatworms infecting blood of the nontetrapod craniates “fishes”. *Adv. Parasitol.* 85, 1–64.
- Overstreet, R.M., Thulin, J., 1989. Response by *Plectropomus leopardus* and other serranid fishes to *Pearsonellum corventum* (Digenea: sanguinicolidae), including melanomacrophage centres in the heart. *Aust. J. Zool.* 37, 129–142.
- Pearce, E.J., MacDonald, A.S., 2002. The immunobiology of schistosomiasis. *Nat. Rev. Immunol.* 2, 499–511.
- Platt, T.R., Blair, D., Purdie, J., Melville, L., 1991. *Griphobilharzia amoena* n. gen. n. sp. (Digenea: Schistosomatidae), a parasite of the freshwater crocodile *Crocodylus johnstoni* (Reptilia: crocodylia) from Australia, with the erection of a new subfamily, *Griphobilharziinae*. *J. Parasitol.* 77, 65–68.
- Platt, T.R., Brooks, D.R., 1997. Evolution of the schistosomes (Digenea: Schistosomatoidea): the origin of dioecy and colonization of the venous system. *J. Parasitol.* 83, 1035–1044.
- Repullés-Albelda, A., Montero, F.E., Holzer, A.S., Ogawa, K., Hutson, K.S., Raga, J.A., 2008. Speciation of the *Paradeontacylix* spp. (sanguinicolidae) of *Seriola dumerili*. Two new species of the genus *Paradeontacylix* from the mediterranean. *Parasitol. Int.* 57, 405–414.
- Schell, S.C., 1974. The life history of *Sanguinicola idahoensis* sp. n. (Trematoda: sanguinicolidae), a blood parasite of steelhead trout *Salmo gairdneri* richardson. *J. Parasitol.* 60, 561–566.
- Shirakashi, S., Kishimoto, Y., Kinami, R., Katano, H., Ishimaru, K., Murata, O., Itoh, N., Ogawa, K., 2012. Morphology and distribution of blood fluke eggs and associated pathology in the gills of cultured Pacific bluefin tuna, *Thunnus orientalis*. *Parasitol. Int.* 61, 242–249.
- Snyder, S.D., 2004. Phylogeny and paraphyly among tetrapod blood flukes (Digenea: Schistosomatidae and spirorchhiidae). *Int. J. Parasitol.* 34, 1385–1392.
- Sugihara, Y., Yamada, T., Tamaki, A., Yamanishi, R., Kanai, K., 2014. Larval stages of the bluefin tuna blood fluke *Cardicola opisthorchis* (Trematoda: Aporocotylidae) found from *Terebella* sp. (Polychaeta: terebellidae). *Parasitol. Int.* 63, 295–299.
- Warren, M.B., Oréllis-Ribeiro, R., Ruiz, C.F., Dang, B.T., Arias, C.R., Bullard, S.A., 2017. Endocarditis associated with blood fluke infections (Digenea: Aporocotylidae: *Psettarium* cf. *anthicum*) among aquacultured cobia (*Rachycentron canadum*) from Nha Trang Bay. *Vietnam Aquacult.* 468, 549–557.
- WoRMS Editorial Board, 2018. World Register of Marine Species. Checklist dataset. <https://doi.org/10.14284/170> accessed via GBIF.org on 2018-10-26.
- Yamaguti, S., 1970. Digenetic Trematodes of Hawaiian Fishes, vol 436p Keigaku Pub. Co., Tokyo.
- Yong, R.Q.-Y., Cutmore, S.C., Miller, T.L., Adlard, R.D., Cribb, T.H., 2013. The ghost of parasites past: eggs of the blood fluke *Cardicola chaetodontis* (Aporocotylidae) trapped in the heart and gills of butterflyfishes (Perciformes: chaetodontidae) of the great barrier reef. *Parasitology* 140, 1186–1194.
- Yong, R.Q.-Y., Cutmore, S.C., Wee, N.Q.-X., Cribb, T.H., 2016. A complex of *Cardicola* (Digenea: Aporocotylidae) species infecting the milkfish, *Chanos chanos* (Gonorynchiformes), with descriptions of two new species. *Syst. Parasitol.* 93, 831–846.
- Yong, R.Q.-Y., Cutmore, S.C., Cribb, T.H., 2018. Two new species of *Cardicola* (trematoda: Aporocotylidae) from the damselfish *Abudefduf whiteleyi* (perciformes: pomacentridae) and the triggerfish *Sufflamen chrysopterum* (tetraodontiformes: balistidae). *Mar. Biodivers.* 1–11. <https://doi.org/10.1007/s12526-018-0895-4>.