



## T follicular helper cells restricted by IRF8 contribute to T cell-mediated inflammation

Ruihua Zhang<sup>a</sup>, Chen-feng Qi<sup>b</sup>, Yuan Hu<sup>a</sup>, Yanhong Shan<sup>a</sup>, Yuan-Pang Hsieh<sup>c</sup>, Feihong Xu<sup>a</sup>, Geming Lu<sup>a</sup>, Jun Dai<sup>a</sup>, Monica Gupta<sup>d</sup>, Miao Cui<sup>a</sup>, Liang Peng<sup>a</sup>, Jianjun Yang<sup>a</sup>, Qingjie Xue<sup>a</sup>, Ray Chen-Liang<sup>a</sup>, Kang Chen<sup>e</sup>, Yeyunfei Zhang<sup>a</sup>, Wai-Ping Fung-Leung<sup>f</sup>, J. Rodrigo Mora<sup>g</sup>, Liwu Li<sup>c</sup>, Herbert C. Morse III<sup>b</sup>, Keiko Ozato<sup>d</sup>, Peter S. Heeger<sup>h,\*</sup>, Huabao Xiong<sup>a,\*</sup>

<sup>a</sup> Department of Medicine, Precision Immunology Institute, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

<sup>b</sup> Laboratory of Immunogenetics, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20892, USA

<sup>c</sup> Department of Biological Sciences, College of Science, Virginia Tech, USA

<sup>d</sup> Programs in Genomics of Differentiation, Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, MD 20892, USA

<sup>e</sup> Department of Obstetrics and Gynecology, Wayne State University, Detroit, MI 48201, USA

<sup>f</sup> Janssen Research & Development, LLC., CA 92121, USA

<sup>g</sup> Janssen R&D, Spring House, PA 19477, USA

<sup>h</sup> Department of Medicine, Translational Transplant Research Center, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

### ARTICLE INFO

#### Keywords:

IRF8  
TFH  
Colitis  
IRF4

### ABSTRACT

The follicular helper T cell (T<sub>FH</sub>) are established regulators of germinal center (GC) B cells, whether T<sub>FH</sub> have pathogenic potential independent of B cells is unknown. Based on *in vitro* T<sub>FH</sub> cell differentiation, *in vivo* T cell transfer animal colitis model, and intestinal tissues of inflammatory bowel disease (IBD) patients, T<sub>FH</sub> and its functions in colitis development were analyzed by FACS, ChIP, ChIP-sequencing, WB, ELISA and PCR. Herein we demonstrate that intestinal tissues of patients and colon tissues obtained from *Rag1*<sup>-/-</sup> recipients of naïve CD4<sup>+</sup> T cells with colitis, each over-express T<sub>FH</sub>-associated gene products. Adoptive transfer of naïve *Bcl6*<sup>-/-</sup> CD4<sup>+</sup> T cells into *Rag1*<sup>-/-</sup> recipient mice abrogated development of colitis and limited T<sub>FH</sub> differentiation *in vivo*, demonstrating a mechanistic link. In contrast, T cell deficiency of interferon regulatory factor 8 (IRF8) resulted in augmentation of T<sub>FH</sub> induction *in vitro* and *in vivo*. Functional studies showed that adoptive transfer of IRF8 deficient CD4<sup>+</sup> T cells into *Rag1*<sup>-/-</sup> recipients exacerbated colitis development associated with increased gut T<sub>FH</sub>-related gene expression, while *Irf8*<sup>-/-</sup>/*Bcl6*<sup>-/-</sup> CD4<sup>+</sup> T cells abrogated colitis, together indicating that IRF8-regulated T<sub>FH</sub> can directly cause colon inflammation. Molecular analyses revealed that IRF8 suppresses T<sub>FH</sub> differentiation by inhibiting transcription and transactivation of the TF IRF4, which is also known to be essential for T<sub>FH</sub> induction. Our documentation showed that IRF8-regulated T<sub>FH</sub> can function as B-cell-independent, pathogenic, mediators of colitis suggests that targeting T<sub>FH</sub> could be effective for treatment of IBD.

### 1. Introduction

T cell help to B cells is a fundamental mechanism for the generation of protective humoral immunity, but over-activation of B cells by T cells can result in excessive humoral immune responses, pathologic inflammation, and autoimmunity [1–3]. A subset CD4<sup>+</sup> T cells, termed follicular helper T cells (T<sub>FH</sub>), are specialized regulators of T cell help to

B cells and are required for induction of germinal center (GC) B cell responses [4,5]. T<sub>FH</sub> stably express C-X-C chemokine receptor 5 (CXCR5), which mediates chemotaxis toward GCs upon ligation by C-X-C chemokine ligand 13 (CXCL13) expressed by follicular dendritic cells (fDCs) [6–8]. Expression of transcription factor (TF) B-cell lymphoma 6 (Bcl-6), among other TFs (e.g. IRF4, c-MAF, Ascl2), requisitely orchestrates T<sub>FH</sub> differentiation [9–16]. T<sub>FH</sub> also express an elaborate

\* Corresponding author. Precision Immunology Institute, Box 1630, Icahn School of Medicine at Mount Sinai, 1 Gustave L. Levy Place, New York, NY 10029-6574, USA.

\*\* Corresponding author. Department of Medicine, Translational Transplant Research Center, Icahn School of Medicine at Mount Sinai, 1 Gustave L. Levy Place, New York, NY 10029-6574, USA.

E-mail addresses: [peter.heeger@mssm.edu](mailto:peter.heeger@mssm.edu) (P.S. Heeger), [huabao.xiong@mssm.edu](mailto:huabao.xiong@mssm.edu) (H. Xiong).

<https://doi.org/10.1016/j.jaut.2018.09.001>

Received 24 May 2018; Received in revised form 6 September 2018; Accepted 7 September 2018

Available online 19 September 2018

0896-8411/ © 2018 Elsevier Ltd. All rights reserved.

network of cell surface molecules that promote T-B cell collaboration in GCs [7,8], including the co-stimulatory receptor inducible co-stimulatory molecule (ICOS) which engages ICOS ligand (ICOSL) on GC B cells [17,18]. ICOS expression is essential for the generation and maintenance of  $T_{FH}$  cells including production of the cytokine interleukin-21 (IL-21) [19,20], which promotes GC B-cell proliferation, class switch recombination (CSR), memory B cell formation and plasma cell differentiation [21,22]. Antigen-experienced  $T_{FH}$  also rapidly up-regulate the expression of CD154 (CD40L) [23,24], which ligates the B cell surface receptor CD40 to induce B cell activation, proliferation, somatic hypermutation (SHM) and class switch recombination (CSR). Engagement of GC B cell-expressed programmed cell death 1 ligand 1 (PD-L1) and/or PD-L2 [25,26] to PD-1 on  $T_{FH}$  cells negatively regulates the size and function of the induced  $T_{FH}$  response. Activated  $T_{FH}$  express the highest levels of CD40L, ICOS, PD-1 and IL-21 among T cell subsets [27–29].

Through decades of research by many investigators, T helper cells including  $T_H1$  and  $T_H2$  cells were initially implicated in the pathogenesis of inflammatory bowel diseases (IBD). Following the discovery of  $T_H17$  cells, which specifically produce  $T_H17$  family cytokines (e.g. IL-17A, IL-17F, and IL-22), investigators reconsidered the  $T_H1/T_H2$  cytokine balance hypothesis, and posited that  $T_H17$  cells are potentially instrumental in IBD pathogenesis. Despite the strong associative evidence between  $T_H17$ -associated genes/mutations and IBD from murine and human studies, clinical trials targeting IL-17A proved ineffective in Crohn's disease patients and in fact paradoxically worsened disease in a set of patients. In addition, naïve  $CD4^+$  T cells from IL-17 deficient mice induced more severe colitis in recipient mice. Thus, the mechanism through which T helper cells mediate inflammatory bowel diseases remains elusive. Interestingly,  $T_{FH}$  signature genes have been expressed in several inflammatory diseases including IBD, suggesting that  $T_{FH}$  may contribute to the development of inflammatory diseases.

Multiple transcription factors, including C-Maf, Batf, Irf4, STAT1, STAT3, and Ascl2, are actively involved in the development and function of  $T_{FH}$  [9–16], but maintenance and full differentiation of  $T_{FH}$  critically requires expression of Bcl-6 [9,10]. In addition, the  $T_{FH}$  differentiation pathway is also opposed by other factors including Blimp-1, Foxo1 and Foxp1 [30,31]. However, the molecular mechanism for the regulation of  $T_{FH}$  is incompletely understood, especially how  $T_{FH}$  are negatively regulated.

IRF8 is a member of the evolutionarily conserved IRF family of transcription factors with diverse and important regulatory roles in the growth, differentiation, and function of innate and adaptive immune cells. IRF8 is expressed by a wide spectrum of immature and mature hematopoietic cells including B cells, dendritic cells (DCs), macrophages, and activated T cells [32,33]. It has an N-terminal DNA-binding domain (DBD) and a C-terminal IRF association domain (IAD), the latter of which is responsible for heterodimerization with other transcription factors [33,34]. IRF8 can function as either a transcriptional repressor or an activator, depending on the specific heterodimeric DNA-binding complexes produced with its varied partners [35–41]. Previously published work showed that germline *Irf8*<sup>-/-</sup> mice develop a chronic myeloid leukemia-like syndrome with impaired  $T_H1$  immunity, but how IRF8 controls T cell function was seldom discussed [33–35,40]. We previously reported that IRF8 negatively regulates  $T_H17$  cell differentiation [42] raising the possibility that this TF could serve as a negative regulator of other  $T_H$  subsets including  $T_{FH}$ .

Herein, we provide new evidence that  $T_{FH}$  can mediate intestinal pathology independent of B cells and show that this pathogenic function is regulated by IRF8 inhibition of IRF4. In addition to providing paradigm shifting mechanistic insight into the functions of  $T_{FH}$  our new findings raise the possibility that  $CD4^+$  T cell-intrinsic IRF8 expression critically regulates other pathogenic autoimmune responses driven by  $T_{FH}$ .

## 2. Materials and methods

### 2.1. Mice

C57BL/6J (B6, stock#000664), *Rag1*<sup>-/-</sup> (on B6 background, B6.129S7-*Rag1*<sup>tm1Mom</sup>/J, stock#002216), *Bcl-6*<sup>-/-</sup> mice (B6.129S(FVB)-*Bcl6*<tm1.1Dent>/J, stock# 023727) were obtained from the Jackson laboratory. *Irf8*<sup>-/-</sup> and Lck-Cre<sup>+</sup>*Irf8*<sup>fl/fl</sup> were maintained in the barrier facility at the Icahn School of Medicine at Mount Sinai. The animal study protocols were approved by the Institutional Animal Care and Use Committees of Icahn School of Medicine at Mount Sinai.

### 2.2. Human colon tissue

Human colon tissues from Crohn's disease patients and control patients undergoing resection for cancer screening were obtained from the Mount Sinai Hospital with a protocol approved by the Institutional Review Board of Icahn School of Medicine at Mount Sinai.

### 2.3. Antibodies

The following antibodies against mouse antigens and conjugated to FITC, PE, PE-Cy5, PerCP-Cy5.5 or APC were purchased from BD Biosciences (CD4 (L3T4), eBioscience: biotin-rat anti-mouse CXCR5 (2G8), plus streptavidin-APC-eFluor 780 or eBioscience: PD-1 (J43), ICOS (7E17G9), Bcl-6 (mG1191E), IL-21 (mhalx21), CD8 (53–6.7), CD3ε (145-2C11), CD62L (MEL-14) and isotype controls.

### 2.4. Intracellular staining and flow cytometry

Naïve  $CD4^+$  T cells ( $CD62L^+CD44^{lo}$ ) were prepared by fluorescence-activated cell sorting (FACS) from spleens and lymph nodes of *Irf8*<sup>-/-</sup> mice or WT littermates. Anti-mouse CD4 microbeads (L3T4, Miltenyi Biotec) from spleen and lymph nodes of mice using AutoMACS separator (Miltenyi Biotec). The sorted cells were stimulated for 72 h with plate-bound anti-CD3 (1 µg/ml; 145-2C11; BD Biosciences) and soluble anti-CD28 (1 µg/ml; 37.51; BD Biosciences) with or without IL-21. The cells were then re-stimulated for 5 h with PMA and ionomycin in the presence of Brefeldin A, cells were fixed with IC Fixation Buffer (BD), incubated with permeabilization buffer, and stained with PE-, APC- or PE-Cy 5.5 anti-mouse antibodies and intracellular cytokines were measured by flow cytometry. Flow cytometry was performed on a FACSCalibur (BD). Cells stimulated under neutral conditions were defined as  $T_H0$  cells. Cells were stimulated to differentiate into  $T_{FH}$  cells by the supplementation with 10 ng/ml IL-21 (R&D Systems). Flow cytometry was performed on a FACSCalibur or LSR Fortessa analyzer (BD Biosciences).

### 2.5. T cell transfer model and histology

T cell transfer experiment was performed as previously described [42]. In brief, purified  $CD4^+CD45RB^{hi}$  T cells from WT, *Irf8*<sup>-/-</sup> and *Bcl6*<sup>-/-</sup> mice with or without  $CD19^+B220^+$  B cells ( $1 \times 10^6$  cells per mouse in 200 µl sterile PBS) from WT mice were transferred intraperitoneally into *Rag1*<sup>-/-</sup> recipients ( $6 \times 10^5$  cells per mouse in 200 µl sterile PBS). Mice were weighed every week throughout the course of experiments. After 5 weeks, mice were sacrificed, their spleens and mesenteric lymph nodes excised then analyzed by flow cytometry. Spleens and colon tissues from WT, *Irf8*<sup>-/-</sup> or *bcl6*<sup>-/-</sup> mice were fixed in 10% neutral buffered formalin and embedded in paraffin or in frozen section. 5 µm sections of tissue were stained with fluorochrome-conjugated antibodies. The degree of inflammation in the epithelium, submucosa and muscularis propria of colon tissue was

scored separately as described by Totsuka et al. [43].

## 2.6. Mice immunization

Mixing anti-mouse CD3 antibody (BD Biosciences, San Jose, CA) 5µg/mouse and anti-mouse CD28 antibody (BD Biosciences, San Jose, CA) 2µg/mouse in 100 µL PBS, injection intraperitoneally two times interval three days. NP(40)-OVA/alum was prepared by mixing NP(40)-OVA (Biosearch Technologies, Petaluma, CA) in PBS with alum (Pierce, Rockford, IL) at a 1:1 ratio. NP-OVA/alum immunizations consisted of 100 µg given intradermal injection (i.d) four times interval one week.

## 2.7. RNA isolation and quantitative real-time RT-PCR

Total RNA was extracted using an RNeasy plus kit (QIAGEN) and cDNA was transcribed using the Superscript II system (Invitrogen) with an oligo-dT primer followed by analysis using iCycler PCR with SYBR Green PCR master Mix (Applied Biosystems) using the primers in Tables S1 and S2. Results were normalized based on the expression of ubiquitin.

## 2.8. Immunoblot

Cells were washed with cold phosphate-buffered saline and lysed for 15 min on ice in 0.5 ml of lysis buffer (50 mM Tris-HCl, pH 8.0, 280 mM NaCl, 0.5% Nonidet P-40, 0.2 mM EDTA, 2 mM EGTA, 10% glycerol and 1 mM dithiothreitol) containing protease inhibitors. Cell lysates were performed for immunoblotting. Anti-IRF4 and anti-IRF8 (SantaCruz Biotechnology) and anti-β-actin (Sigma) antibodies were used according to the manufactures' instructions. Secondary antibodies were purchased from SantaCruz Biotechnology.

## 2.9. Chromatin immunoprecipitation (ChIP)

ChIP was performed using a kit following the manufacturers' instruction (Upstate Biotechnology). Briefly, cells were fixed by 1% formaldehyde for 10 min at 37 °C. Nuclei were purified and sonicated to obtain DNA fragments. Chromatin fractions were pre-cleared with protein A-conjugated agarose beads followed by immunoprecipitation overnight at 4 °C with 3 µg of anti-IRF8 or control antibody. Crosslinking was reversed at 65 °C for 4 h, followed by proteinase K digestion. DNA was purified and subjected to qPCR. The input DNA was diluted 200 times before PCR amplification. The input and immunoprecipitated DNAs were amplified by qPCR using the primers targeting.

## 2.10. Chromatin immunoprecipitation sequence (ChIP-Seq)

Purified CD4<sup>+</sup> T cells from WT mice were stimulated with or without anti-CD3 and anti-CD28 for 48 h in the presence of IL-21. The cell pellet was cross-linked and sonicated, the chromatin was ready for MOWChIP-seq. The mixed beads of protein A beads and protein G beads were incubated with IRF8 antibody. After washing, the IRF8-beads were loaded to chamber and incubate with sonicated chromatin samples. After ChIP, the washed immune complexes were collected and resuspended in the reverse crosslinking buffer to incubate. The DNA was extracted and precipitation. Sequencing libraries were prepared by Accel-NGS 2S Plus DNA Library Kit (Swift). The libraries were sequenced on an Illumina HiSeq 4000 with single-end 50-nt reads.

## 2.11. Statistical analysis

Statistical analysis was performed using Student's *t*-test for most of the experiments. *P* < 0.05 was considered statistically significant.

## 3. Results

### 3.1. T<sub>FH</sub> are critically involved in intestinal inflammation


While the primary known function of T<sub>FH</sub> is to provide helper signals that drive GB B cell differentiation [5–8], it is possible that T<sub>FH</sub> can directly function as pathogenic mediators of autoimmune injury independent of their helper functions. To investigate the possibility that T<sub>FH</sub> cells contribute to intestinal inflammation, we first examined the expression of T<sub>FH</sub> signature genes in the intestinal tissues of Crohn's disease patients. These analyses revealed that T<sub>FH</sub>-associated gene products including IL-21, CXCR5, ICOS, PD1 and Bcl-6 were significantly up-regulated in intestinal tissues of Crohn's disease patients compared to normal controls (Fig. S1). Similarly, when we analyzed the expression of T<sub>FH</sub> signature genes in the intestinal tissues of a murine model in which adoptive transfer of CD4<sup>+</sup> T cells together with or without B cells into syngeneic *Rag1*<sup>-/-</sup> hosts reproducibly results in colitis [42,43] (Fig. S2A and data not shown), we observed significantly higher expression of the same T<sub>FH</sub>-related gene products (Fig. S2B). Histological staining showed that Bcl-6 was highly expressed in the intestinal tissues with colitis (Fig. S2C) and flow cytometry analysis of spleen cells showed detectable CXCR5<sup>+</sup>ICOS<sup>+</sup>PD1<sup>+</sup>BCL-6<sup>+</sup> T<sub>FH</sub> only in *Rag1*<sup>-/-</sup> recipients of WT naïve CD4<sup>+</sup> T cells (Fig. S2D).

As Bcl-6 is one of several TFs required for T<sub>FH</sub> differentiation [9,10] we directly tested the pathogenicity of T<sub>FH</sub> by adoptively transferring naïve *Bcl-6*<sup>-/-</sup>CD4<sup>+</sup> T cells into *Rag1*<sup>-/-</sup> recipients (Fig. 1A–B). Distinct from recipients of WT CD4<sup>+</sup> T cells, recipients of *Bcl-6*<sup>-/-</sup> CD4<sup>+</sup> CD45Rb<sup>hi</sup> cells appeared phenotypically normal and maintained rather than lost weight. Histological analyses of the intestinal tissue 6 weeks post-transfer revealed less inflammatory cell infiltration and lower pathology scores in the *Rag1*<sup>-/-</sup> recipients of *Bcl-6*<sup>-/-</sup> CD4<sup>+</sup> cells (Fig. 1C). When we analyzed T cell phenotypes of mesenteric lymph nodes at 6 weeks we observed significantly fewer T<sub>FH</sub> in adoptive recipients of *Bcl-6*<sup>-/-</sup> CD4<sup>+</sup> cells (Fig. 1D). Taken together, the results support the conclusion that T<sub>FH</sub> rather than T<sub>H1</sub> or T<sub>H17</sub> cells (data not shown) are pathogenic mediators of intestinal inflammation in this model.

### 3.2. IRF8-deficient mice display enhanced T<sub>FH</sub> cell differentiation

Building upon our previous observation that IRF8 inhibits T<sub>H17</sub> differentiation [42] we tested the impact of IRF8 on differentiation of naïve CD4<sup>+</sup> T cells toward the T<sub>FH</sub> phenotype using an *in vitro* culture system (Fig. 2). Three days after stimulating purified naïve *Irf8*<sup>-/-</sup> or WT CD4<sup>+</sup> T cells with anti-CD3/CD28 under T<sub>FH</sub>-inducing conditions, we analyzed cytokine, surface marker and TF profiles in the responding T cells. These assays showed higher proportions of CXCR5<sup>+</sup>PD-1<sup>+</sup>, CXCR5<sup>+</sup>ICOS<sup>+</sup> and CXCR5<sup>+</sup>Bcl-6<sup>+</sup> cells (Fig. 2A–C) and more IL-21, within the *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells vs. WT controls (Fig. 2D and E). Quantitative PCR assays showed higher *Icos*, *Pdcd1*, *Cxcr5* and *Il21* gene expression in the *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells (Fig. 2F). Analysis of CD4<sup>+</sup> T cells obtained from mice with IRF8 deficiency restricted to T cells (Lck-Cre<sup>+</sup> *Irf8*<sup>fl/fl</sup>) confirmed enhanced T<sub>FH</sub> cell differentiation (Fig. 3A–C). Proliferative responses of WT and *IRF8*<sup>-/-</sup> CD4<sup>+</sup> T cells cultured under T<sub>FH</sub>-inducing conditions did not differ (Fig. S3), and IRF8 deficiency did not alter *in vitro* induction of IFNγ<sup>+</sup> T<sub>H1</sub>, IL-4<sup>+</sup> T<sub>H2</sub> or Foxp3<sup>+</sup> T<sub>reg</sub> cells (Fig. S4) [42], together demonstrating that IRF8 specifically inhibits the T<sub>FH</sub> differentiation program.

To begin to assess how the above observed effects of T cell IRF8 deficiency apply *in vivo*, we injected groups of Lck-Cre<sup>+</sup> *Irf8*<sup>fl/fl</sup> and Lck-Cre<sup>+</sup> *Irf8*<sup>wt/wt</sup> mice with anti-CD3 mAb (Fig. 4) and independently we immunized groups of animals with 4-hydroxy-3-nitrophenylacetyl-conjugated ovalbumin (NP-OVA, Fig. S5). Under both conditions, flow cytometry analyses revealed ~2-fold higher frequencies of T<sub>FH</sub> cells in the spleens of treated Lck-Cre<sup>+</sup> *Irf8*<sup>fl/fl</sup> mice (Fig. 4A and Fig. S5). Quantification of T<sub>H1</sub>, T<sub>H2</sub> and T<sub>reg</sub> did not differ between groups in



**Fig. 1.** T<sub>FH</sub> signature is up-regulated in autoimmune and inflammatory diseases. (A and B) Colon morphology and body weight of *Rag1*<sup>-/-</sup> recipient mice after receiving  $6 \times 10^5$  purified WT (n = 5) or *Bcl-6*<sup>-/-</sup> (n = 5) CD4<sup>+</sup> T cells. (C) H&E staining and histology score of colon tissues of *Rag1*<sup>-/-</sup> recipient mice after receiving  $6 \times 10^5$  purified WT (n = 5) or *Bcl-6*<sup>-/-</sup> (n = 5) CD4<sup>+</sup> T cells. (D) The percentages of CXCR5<sup>+</sup>ICOS<sup>+</sup> and CXCR5<sup>+</sup>PD-1<sup>+</sup> CD4<sup>+</sup> T cells were compared between *Rag1*<sup>-/-</sup> recipients of WT CD4<sup>+</sup> T cells (n = 5) and those of *Bcl-6*<sup>-/-</sup> CD4<sup>+</sup> T cells (n = 5).

either set of experiments (Fig. S4), validating that the *in vitro* findings (Fig. 2) apply *in vivo* and supporting the concept that IRF8 specifically regulates T<sub>FH</sub> differentiation [42].


### 3.3. IRF8-deficient T<sub>FH</sub> cells are pathogenic mediators of colitis

To directly test the hypothesis that IRF8 negatively regulates T<sub>FH</sub> capable of mediating colonic pathology, we adoptively transferred WT or *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells into groups of *Rag1*<sup>-/-</sup> mice and followed the animals for up to 6 weeks, analyzing and comparing clinical disease expression, colon histology and splenic T cell responses between groups. These analyses showed *Rag1*<sup>-/-</sup> recipients of *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells exhibited greater and more rapid weight loss with more severe histological changes in the colons (Fig. 5A and B) and higher frequencies of splenic T<sub>FH</sub> (Fig. 5C).

To confirm that absence of T cell IRF8 exacerbates colitis through a T<sub>FH</sub>-dependent mechanism, we adoptively transferred naive *Irf8*<sup>-/-</sup> *Bcl-6*<sup>-/-</sup> CD4<sup>+</sup> T cells into *Rag1*<sup>-/-</sup> recipients. These experiments showed diminished colitis severity (Fig. 6A–C), similar to that observed in the recipients of *Bcl-6*<sup>-/-</sup> CD4<sup>+</sup> T cells (Fig. 1B). Frequencies of splenic T<sub>FH</sub> were also reduced in *Irf8*<sup>-/-</sup> *Bcl-6*<sup>-/-</sup> recipient mice compared to WT controls (Fig. 6D). Taken together, the data support the conclusion that IRF8 suppresses inflammation via the inhibiting differentiation of pathogenic T<sub>FH</sub>.

### 3.4. Mice with T cell-specific IRF8 deficiency exhibit exaggerated B cell differentiation

The data above suggests that IRF8-regulated T<sub>FH</sub> cells play an important role in T cell-mediated inflammation. To determine whether the enhanced T<sub>FH</sub> signature functionally affects B cell development, we analyzed mice with germline (*Irf8*<sup>-/-</sup>) or T cell-specific IRF8 deficiency (*Lck-Cre*<sup>+</sup> *Irf8*<sup>fl/fl</sup>). These mice had enlarged spleens and lymph nodes (Fig. S6A and B) that harbored increased numbers of CD19<sup>+</sup>CD138<sup>+</sup> plasma cells (Fig. S6C) and germinal center B cells (Fig. S6D). Consistently, *Lck-Cre*<sup>+</sup> *Irf8*<sup>fl/fl</sup> mice had increased levels of serum IgG and IgM (Fig. S6E). The expansion of plasma cells in secondary lymphoid organs of *Irf8*<sup>-/-</sup> mice was not due to infection or autoimmunity, as CD11c<sup>+</sup> DCs were not increased in any immune organ or tissue analyzed (Fig. S7A), and there was no autoimmune kidney damage, such as cellular infiltration and glomerular crescents, in *Irf8*<sup>-/-</sup> mice (Fig. S7B). There is no significant expansion in population of T cells but as reported data that myeloid cells were obviously expanded (Fig. S7A). To determine whether *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells also induce more B cell proliferation *in vivo*, we co-transferred WT B cells with either WT or *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells into *Rag1*<sup>-/-</sup> mice. Splenomegaly after adoptive transfer was markedly increased in recipients of *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells (Fig. S6F), and elevated percentage of B cells was found in the spleen and lymph nodes (Fig. S6G). These results suggest that IRF8 controls the magnitude of humoral immunity by regulating the



**Fig. 2.** IRF8-deficient CD4<sup>+</sup> T cells show enhanced T<sub>FH</sub> polarization after *ex vivo* stimulation. Purified CD4<sup>+</sup> T cells from WT or *Irf8*<sup>-/-</sup> mice (A, B, and C) were stimulated with or without anti-CD3 and anti-CD28 for 48 h in the presence of IL-21. Cell surface expression of PD-1, ICOS, CXCR5 and intracellular expression of Bcl-6 were analyzed by flow cytometry. The percentages of CXCR5<sup>+</sup>PD1<sup>+</sup>, CXCR5<sup>+</sup>ICOS<sup>+</sup> and CXCR5<sup>+</sup>Bcl6<sup>+</sup> cells were compared between WT (n = 5) and *Irf8*<sup>-/-</sup> (n = 5) CD4<sup>+</sup> T cell cultures. (D) Flow cytometry evaluation of IL-21<sup>+</sup>CD4<sup>+</sup> T cells in WT and *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells under T<sub>FH</sub> condition (n = 5). (E) ELISA of IL-21 levels in culture supernatant of WT and *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells cultured under T<sub>FH</sub> condition (n = 4). (F) Quantitative real-time RT-PCR analysis of T<sub>FH</sub>-associated genes in WT and *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells under T<sub>FH</sub> condition (n = 5). The ubiquitin gene (*Ubi*) was used as an internal control. Results shown are representative of three independent experiments. Data are given as means ± SEM.


differentiation and function of T<sub>FH</sub> cells. In addition to regulating the function of CD4<sup>+</sup> T cells in GC B cell response, IRF8 expression in CD4<sup>+</sup> T cells may influence B cell differentiation prior to the GC stage. To test this possibility, we co-transferred C57BL/6 (H-2<sup>b</sup>) bone marrow cells depleted of T cells with either WT or *Irf8*<sup>-/-</sup> C57BL/6 (H-2<sup>b</sup>) CD4<sup>+</sup> T cells into irradiated allogeneic BALB/c (H-2<sup>d</sup>) recipients. Co-transferred *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells induced more B cell development from H-2<sup>b</sup> bone marrow cells than co-transferred WT CD4<sup>+</sup> T cells (Fig. S8). Therefore, IRF8 in CD4<sup>+</sup> T cells also restrains the ability of CD4<sup>+</sup> T cells to support B cell development *in vivo*.

### 3.5. IRF8 suppresses IRF4 expression in CD4<sup>+</sup> T cells

IRF4 is important for the differentiation of various T helper cell subsets, including T<sub>FH</sub>, T<sub>H2</sub>, T<sub>H9</sub> and T<sub>H17</sub> cells [11,14,34]. IRF4 and IRF8 represent immune-specific members of the interferon regulatory family and co-operate to play major roles in controlling the development and functioning of T cell subsets and other immune cells. To better understand the molecule mechanism of IRF8 in controlling of T<sub>FH</sub> cell function, we purified CD4<sup>+</sup> T cells from WT or *Irf8*<sup>-/-</sup> mice and

stimulated with or without anti-CD3 and anti-CD28 under T<sub>FH</sub> cells culture condition. The results clearly indicated that *Irf4* and *Il21* expression levels were significantly increased in *Irf8*<sup>-/-</sup> mice compared with WT mice by microarray analysis (Fig. 7A). Furthermore, we evaluated the *Irf4* gene by Quantitative real-time-PCR analysis (Fig. 7B) and protein expression levels of CXCR5, Bcl-6, IRF4 and IRF8 by western blotting (Fig. 7E) in WT and *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells under T<sub>FH</sub> condition for time course. The results showed that both the *Irf4* gene and IRF4 protein were significantly up-regulated in *Irf8*<sup>-/-</sup> mice compared with WT mice (Fig. 7A, B and E). We performed ChIP-seq assay to understand whether IRF8 binds to the promoter region of IRF4 gene in T<sub>FH</sub> cell condition. The result showed that IRF8 indeed bound to the promoter region of IRF4 gene (Fig. 7C). Furthermore, the ChIP assays also showed more IRF4 protein bound to the *Il21* promoter in CD4<sup>+</sup> T cells of *Irf8*<sup>-/-</sup> mice under T<sub>FH</sub> cell culture condition when compared with WT mice (Fig. 7D). The results suggested that IRF8 suppresses T<sub>FH</sub> differentiation by inhibiting the DNA binding activity of IRF4 to the promoter region of *Il21* gene.


We also sought to determine if IRF8 controls the expression of CD40 ligand (CD40L), a molecule crucial to the function of T<sub>FH</sub> cells.




**Fig. 3.** IRF8-deficient in T cells show enhanced T<sub>FH</sub> polarization after *ex vivo* stimulation. Purified CD4<sup>+</sup> T cells from *Irf8*<sup>wt/wt</sup> or *Irf8*<sup>lck/lck</sup> mice (A, B, and C) were stimulated with or without anti-CD3 and anti-CD28 for 48 h in the presence of IL-21. Cell surface expression of PD-1, ICOS, CXCR5 and intracellular expression of Bcl-6 were analyzed by flow cytometry. The percentages of CXCR5<sup>+</sup>PD-1<sup>+</sup>, CXCR5<sup>+</sup>ICOS<sup>+</sup> and CXCR5<sup>+</sup>Bcl6<sup>+</sup> cells were compared between *Irf8*<sup>wt/wt</sup> (n = 5) and *Irf8*<sup>lck/lck</sup> (n = 5) CD4<sup>+</sup> T cell cultures. Results shown are representative of three independent experiments. Data are given as means ± SEM.

Compared to WT CD4<sup>+</sup> T cells, *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells expressed higher protein and mRNA levels of CD40L following *ex vivo* stimulation in T<sub>FH</sub> conditions (Fig. 7F–H). The same results for protein expression were also seen in *ex vivo* stimulated CD4<sup>+</sup> T cells from Lck-Cre<sup>+</sup> *Irf8*<sup>fl/fl</sup> mice (Fig. 7I). Upon TCR stimulation by anti-CD3 Ab *in vivo*, CD40L

expression was higher on CD4<sup>+</sup> T cells of Lck-Cre<sup>+</sup> *Irf8*<sup>fl/fl</sup> mice than those of WT control mice in mesenteric lymph nodes (Fig. S9A). Considering that IRF8 can function as a transcriptional repressor, we performed a chromatin immunoprecipitation (ChIP) assay to determine if IRF8 can repress the *Cd40lg* gene in CD4<sup>+</sup> T cells. IRF8 bound to three



**Fig. 4.** T<sub>FH</sub>-associated signatures were significantly increased in IRF8-deficient T cells *in vivo*. Cell surface expression of ICOS, PD-1, CXCR5 and intracellular expression of Bcl-6 in CD4<sup>+</sup> T cells of Lck-Cre<sup>+</sup> *Irf8*<sup>wt/wt</sup> or Lck-Cre<sup>+</sup> *Irf8*<sup>fl/fl</sup> mice after anti-CD3 administration two times interval three days intraperitoneally, analyzed by flow cytometry. The percentages of CXCR5<sup>+</sup>ICOS<sup>+</sup>, CXCR5<sup>+</sup>PD-1<sup>+</sup> and CXCR5<sup>+</sup>Bcl6<sup>+</sup> cells in CD4<sup>+</sup> T cells of spleen were compared between Lck-Cre<sup>+</sup> *Irf8*<sup>wt/wt</sup> (n = 6) and Lck-Cre<sup>+</sup> *Irf8*<sup>fl/fl</sup> (n = 6) mice. Results shown are representative of three independent experiments. Data are given as means ± SEM.



**Fig. 5.** T<sub>FH</sub> signature is up-regulated in autoimmune and inflammatory diseases. (A) Body weight of *Rag1*<sup>-/-</sup> recipient mice after receiving  $6 \times 10^5$  purified CD4<sup>+</sup> T cells from Lck-Cre<sup>+</sup> *Irf8*<sup>wt/wt</sup> (n = 5) or Lck-Cre<sup>+</sup> *Irf8*<sup>fl/fl</sup> (n = 5) mice with CD19<sup>+</sup>B220<sup>+</sup> B cells from WT mice. (B) Histological staining of colon tissues and histology score of *Rag1*<sup>-/-</sup> recipients of Lck-Cre<sup>+</sup> *Irf8*<sup>wt/wt</sup> CD4<sup>+</sup> T cells (n = 5) and those of Lck-Cre<sup>+</sup> *Irf8*<sup>fl/fl</sup> CD4<sup>+</sup> T cells (n = 5). (C) The percentages of CXCR5<sup>+</sup>ICOS<sup>+</sup>, CXCR5<sup>+</sup>PD-1<sup>+</sup> and CXCR5<sup>+</sup>Bcl6<sup>+</sup> CD4<sup>+</sup> T cells were compared between *Rag1*<sup>-/-</sup> recipients of Lck-Cre<sup>+</sup> *Irf8*<sup>wt/wt</sup> CD4<sup>+</sup> T cells (n = 5) and those of Lck-Cre<sup>+</sup> *Irf8*<sup>fl/fl</sup> CD4<sup>+</sup> T cells (n = 5) and WT as control (n = 5). Results shown are representative of three independent experiments. Data are given as means  $\pm$  SD.


of four putative IRF8-binding sites in the *Cd40lg* promoter and one experiment of ChIP data shown as Fig. 7J. Consistently, IRF8 suppressed CD40L expression in CD4<sup>+</sup> T cells in a cell-intrinsic fashion, as evidenced by reduced cell surface CD40L levels in CD4<sup>+</sup> T cells transduced by an *Irf8*-encoding retrovirus (Fig. S9B), as compared to marked increases in CD40L staining in cells transfected with empty virus. Taken together, these results suggest that IRF8 represses CD40L expression at both the transcriptional and protein levels in T<sub>FH</sub> cells.

#### 4. Discussion

Our data newly and uniquely demonstrate a proinflammatory role for T<sub>FH</sub> as mediators of colitis independent of their ability to provide helper signals to B cells. We show that intestinal tissues of inflammatory bowel disease (IBD) patients and colon tissues obtained from *Rag1*<sup>-/-</sup> recipients of naive CD4<sup>+</sup> T cells with colitis, each over-express T<sub>FH</sub>-associated gene products. Adoptive transfer of naive *Bcl6*<sup>-/-</sup> CD4<sup>+</sup> T cells into *Rag1*<sup>-/-</sup> recipient mice abrogated development of colitis and limited T<sub>FH</sub> differentiation *in vivo*, demonstrating a mechanistic link. In contrast, T cell deficiency of interferon regulatory factor 8 (IRF8) resulted in augmentation of T<sub>FH</sub> induction *in vitro* and *in vivo*. Functional

studies showed that adoptive transfer of *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells into *Rag1*<sup>-/-</sup> recipients exacerbated colitis development associated with increased gut T<sub>FH</sub>-related gene expression, while *Irf8*<sup>-/-</sup>/*Bcl6*<sup>-/-</sup> CD4<sup>+</sup> T cells abrogated colitis, together indicating that IRF8-regulated T<sub>FH</sub> can directly cause colon inflammation. Molecular analyses revealed that IRF8 suppresses T<sub>FH</sub> differentiation by inhibiting transcription and transactivation of the TF IRF4, which is also known to be essential for T<sub>FH</sub> induction.

T<sub>FH</sub> cells are a distinct T helper cell subset shown to coordinate generation of the germinal center (GC) responses by initiating help for antigen specific B cells. Through this B cell help mechanism, T<sub>FH</sub> have been shown to be pathogenic in a number of autoimmune diseases. Increased frequencies of T<sub>FH</sub>-like cells in peripheral blood are observed in subsets of patients with Sjogren's syndrome, juvenile dermatomyositis, and systemic lupus erythematosus [44,45], each of which is associated extensive autoantibody production. However, several observations in the literature hint that T<sub>FH</sub> have other functions beyond providing helper signals for B cell differentiation and antibody switching. T<sub>FH</sub> are found within injured organs and tissues in subjects with lupus nephritis [46], multiple sclerosis [47], inflammatory arthritis, type 1 diabetes [48], and intestinal tissues of patients with IBD



**Fig. 6.**  $T_{FH}$  signature is up-regulated in autoimmune and inflammatory diseases. Colon morphology and body weight of  $Rag1^{-/-}$  recipient mice after receiving  $6 \times 10^5$  purified WT (n = 5) or  $Irf8^{-/-}Bcl6^{-/-}$  (n = 6)  $CD4^+$  T cells with  $CD19^+B220^+$  B cells from WT mice (A and B). (C) H&E staining of colon tissues of  $Rag1^{-/-}$  recipient mice after receiving  $6 \times 10^5$  purified WT (n = 5) or  $Irf8^{-/-}Bcl6^{-/-}$  (n = 6)  $CD4^+$  T cells. (D) The percentages of  $CXCR5^+ICOS^+$  and  $CXCR5^+PD-1^+$   $CD4^+$  T cells were compared between  $Rag1^{-/-}$  recipients of WT  $CD4^+$  T cells (n = 5) and those of  $Irf8^{-/-}Bcl6^{-/-}$   $CD4^+$  T cells (n = 6). Results shown are representative of five mice. Data are given as means  $\pm$  SD.


[49]. Our data using adoptive transfer of Bcl-6 deficient naïve  $CD4^+$  T cells into  $Rag1^{-/-}$  recipients provide direct evidence in support of this hypothesis linking  $T_{FH}$  to intestinal inflammation.

$T_{FH}$ -produced IL-21, which is the key cytokine shown to drive B cell differentiation in GCs [19–22]. Early studies revealed that IL-21 is markedly overproduced in inflamed guts of patients with IBD compared to non-inflamed controls [49]. Mice lacking IL-21 are protected against chemically induced colitis; moreover, wild type mice given a neutralizing IL-21R/Fc fusion protein exhibit less experimental colitis as compared to control mice. IL-21 has been shown to activate macrophages, participate in granuloma formation, and inhibit induction of regulatory T cells (Treg) in murine graft vs host diseases. These findings together raise the possibility that  $T_{FH}$ /IL-21 have pathogenic potential. In the present study, we demonstrate that IL-21 is highly expressed in intestinal tissues of IBD patients and also in the colon tissues of recipient mice transferred with naïve  $CD4^+$  T cells with colitis. Based these published findings and our new data, including the strong association of  $T_{FH}$ -related genes to human IBD (Fig. S1), we speculate that  $T_{FH}$ -produced IL-21 is one pathological mediator of intestinal inflammation.

IRF8 plays critical roles in the differentiation of myeloid cells, B cells, dendritic cells and T cells, and hence the regulation of both innate and adoptive immune responses [11,32–42]. A recent genome-wide

association (GWA) study has strongly implicated a variant near the IRF8 gene in SLE susceptibility in Europeans [50]. In addition, a different genetic variation in the IRF8 gene has been implicated in multiple sclerosis [51]. A GWA study identified the IRF8 gene as strongly associated with development of Crohn's disease [52], but provided no mechanistic explanation. Our findings in which we demonstrate that IRF8 deficiency a) favors T cell differentiation toward the  $T_{FH}$  lineage and b) exacerbates  $CD4^+$  T cell-mediated colitis provide a potential explanation to account for this GWAS association. The observations support the testable hypothesis that IRF8 mutations that diminish production or function of IRF8 lead to augmented  $T_{FH}$  differentiation which in turn contributes to the development of IBD.

Published evidence demonstrates that IRF8 cooperates with other TFs including Bcl-6, IRF4, and PU.1 to regulate immune cell function [11,33–42]. IRF8 and IRF4 are immune-specific members of the IRF TF family and have evolved not only to interact with specific members of the Ets superfamily, e.g. PU.1 and Spi-B, but also with particular members of the AP-1 superfamily, e.g. BATF-containing heterodimers [40,41]. IRF4 is important for the differentiation of various T helper cell subsets, including  $T_{FH}$ ,  $T_H2$ ,  $T_H9$  and  $T_H17$  cells [11,14,34]. Adding to this literature, in the present study, we provide new insight into mechanistic links among IRF8, IRF4 and IL-21 producing  $T_{FH}$ . We demonstrate that IRF8 binds to the promoter region of IRF4 gene and that



**Fig. 7. Regulation of IRF4 expression by IRF8.** (A) Purified CD4<sup>+</sup> T cells from WT or *Irf8*<sup>-/-</sup> mice were stimulated with or without anti-CD3 and anti-CD28 for 48 h in the presence of IL-21. Microarray analysis of the gene expression by WT and *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells under T<sub>FH</sub> condition. (B) Quantitative real-time RT-PCR analysis of *irf4* gene level in WT and *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells under T<sub>FH</sub> condition for time course (n = 5). The ubiquitin gene (*Ubi*) was used as an internal control. (C) IRF8 ChIP-seq signals at the IRF4 gene locus. Blue line denotes the range of the IRF4 gene locus, and the peak patterns in red represent the ChIP-seq signals of IRF8 binding. Signals represent DNA fragments that were captured by the IRF8 antibody through ChIP. (D) Purified CD4<sup>+</sup> T cells from WT or *Irf8*<sup>-/-</sup> mice were stimulated with anti-CD3 and anti-CD28 for 48 h in the presence of IL-21, followed by ChIP assay. Three micrograms of an anti-IRF4 antibody or isotype-matched IgG as control antibody were used in the immunoprecipitation step. PCR was used to quantify the amount of precipitated DNA with primers flanking the *irf4*-binding site of the IL-21 promoter region. Each bar represents mean ± S.D. from three independent experiments, unpaired Student's t-test, \*P < 0.05, versus WT cells. (E) Western blot analysis of IRF4, IRF8, Bcl-6 and CXCR5 protein levels in WT and *Irf8*<sup>-/-</sup> CD4<sup>+</sup> T cells under T<sub>FH</sub> condition for time course. The β-actin was used as an internal control. (F) Purified CD4<sup>+</sup> T cells from WT or *Irf8*<sup>-/-</sup> mice were stimulated with anti-CD3 and anti-CD28 antibody for 48 h. CD40L expression was analyzed by flow cytometry gated on CD4<sup>+</sup> T cells and used isotype control. (G) *Cd40l*g RNA level was analyzed by real-time RT-PCR (n = 5). (H) Soluble CD40L protein level in culture media was measured by ELISA. (I) The experiment was repeated by using purified CD4<sup>+</sup> T cells of Lck-Cre<sup>+</sup> *Irf8*<sup>wt/wt</sup> (n = 6) or Lck-Cre<sup>+</sup> *Irf8*<sup>fl/fl</sup> (n = 6) mice. (J) ChIP assay of CD4<sup>+</sup> T cells from WT and *Irf8*<sup>-/-</sup> mice was analyzed for binding of IRF8 to the *Cd40l*g promoter (n = 4). Results shown are representative of three independent experiments. Data are given as means ± SEM. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

in the absence of IRF8, T cells express more IRF4. In addition, IRF8 deficiency significantly enhanced the IRF4 binding to the promoter region of IL-21 gene, together supporting the conclusion that IRF8 suppresses T<sub>FH</sub> differentiation by inhibiting transcription and transactivation of the TF IRF4.

Taken together, our murine and human data has uncovered potentially important pathogenic role for T<sub>FH</sub> cells in inflammatory diseases, and suggests that the suppressive function of IRF8 in T<sub>FH</sub> differentiation and the function may be potentially targeted to treat these diseases.

**Conflicts of interest**

The authors declare that they have no competing interests.

**Acknowledgments**

We are grateful for Dr. Feng Hong for technical support, Mr. Anthony Bonito and Mr. Zihang Zheng for reading the manuscript. Huabao Xiong was supported by the NIH grant R01AI104688. Peter. S Heeger was supported by R01AI132405 and R01AI071185. This work was supported in part by the Intramural Research Program of NICHD and NIAID.

**Appendix A. Supplementary data**

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jaut.2018.09.001>.

## References

- [1] J.E. Craft, Follicular helper T cells in immunity and systemic autoimmunity, *Nat. Rev. Rheumatol.* 8 (2012) 337–347.
- [2] C. King, S.G. Tangye, C.R.T. Mackay, Follicular helper (TFH) cells in normal and dysregulated immune responses, *Annu. Rev. Immunol.* 26 (2008) 741–766.
- [3] S.G. Tangye, C.S. Ma, R. Brink, E.K. Deenick, The good, the bad and the ugly - TFH cells in human health and disease, *Nat. Rev. Immunol.* 13 (2013) 412–426.
- [4] T. Chtanova, S.G. Tangye, R. Newton, N. Frank, M.R. Hodge, M.S. Rolph, C.R. Mackay, T follicular helper cells express a distinctive transcriptional profile, reflecting their role as non Th1/Th2 effector cells that provide help for B cells, *J. Immunol.* 173 (2004) 68–78.
- [5] J.J. O'Shea, W.E. Paul, Mechanisms underlying lineage commitment and plasticity of helper CD4<sup>+</sup> T cells, *Science* 327 (2010) 1098–1102.
- [6] D. Breitfeld, L. Ohl, E. Kremmer, J. Ellwart, F. Sallusto, M. Lipp, R. Förster, Follicular B helper T cells express CXC chemokine receptor 5, localize to B cell follicles, and support immunoglobulin production, *J. Exp. Med.* 192 (2000) 1545–1552.
- [7] C.H. Kim, L.S. Rott, I. Clark-Lewis, D.J. Campbell, L. Wu, E.C. Butcher, Subspecialization of CXCR5<sup>+</sup> T cells: B helper activity is focused in a germinal center-localized subset of CXCR5<sup>+</sup> T cells, *J. Exp. Med.* 193 (2001) 1373–1381.
- [8] R. Morita, N. Schmitt, S.E. Bentebibel, R. Ranganathan, L. Bourdery, G. Zurawski, E. Foucat, M. Dullaers, S. Oh, N. Sabzghabaei, Human blood CXCR5<sup>+</sup>CD4<sup>+</sup> T cells are counterparts of T follicular cells and contain specific subsets that differentially support antibody secretion, *Immunity* 34 (2011) 108–121.
- [9] M. Kitano, S. Moriyama, Y. Ando, M. Hikida, Y. Mori, T. Kurosaki, T. Okada, Bcl6 protein expression shapes pre-germinal center B cell dynamics and follicular helper T cell heterogeneity, *Immunity* 34 (2011) 961–972.
- [10] R.I. Nurieva, Y. Chung, G.J. Martinez, X.O. Yang, S. Tanaka, T.D. Matskevitch, Y.H. Wang, C. Dong, Bcl6 mediates the development of T follicular helper cells, *Science* 325 (2009) 1001–1005.
- [11] R.H. Zhang, K. Chen, L. Peng, H.B. Xiong, Regulation of T helper cell differentiation by interferon regulatory factor family members, *Immunol. Res.* 54 (1) (2012) 169–176.
- [12] R.J. Johnston, Y.S. Choi, J.A. Diamond, J.A. Yang, S. Crotty, STAT5 is a potent negative regulator of TFH cell differentiation, *J. Exp. Med.* 209 (2012) 243–250.
- [13] M.A. Kroenke, D. Eto, M. Locci, M. Cho, T. Davidson, E.K. Haddad, S. Crotty, Bcl6 and maf cooperate to instruct human follicular helper CD4 T cell differentiation, *J. Immunol.* 188 (2012) 3734–3744.
- [14] Nadine Bollig, Anne Brüstle, Kerstin Kellner, Waltraud Ackermann, Elfadil Abass, Raifer Hartmann, Bärbel Camara, Cornelia Brendel, Gavin Giel, Evita Bothur, Magdalena Huber, Christoph Paul, Alexandra Elli, Richard A. Kroccek, Roza Nuriev, Chen Dong, Ralf Jacob, Tak W. Mak, Michael Lohoff, Transcription factor IRF4 determines germinal center formation through follicular T-helper cell differentiation, *Proc. Natl. Acad. Sci.* 109 (22) (2012) 8664–8669.
- [15] Xindong Liu, Xin Chen, Bo Zhong, Aibo Wang, Xiaohu Wang, Fuliang Chu, Roza I. Nurieva, Xiaowei Yan, Ping Chen, Laurens G. van der Flier, Hiroko Nakatsukasa, Sattva S. Neelapu, Wanjun Chen, Hans Clevers, Qiang Tian, Hai Qi, Lai Wei, Chen Dong, Transcription factor achaete-scute homologue 2 initiates follicular T-helper-cell development, *Nature* 507 (27) (2014) 513–519.
- [16] B. Schiemann, J.L. Gommerman, K. Vora, T.G. Cachero, S. Shulga-Morskaya, M. Dobles, An essential role for BAFF in the normal development of B cells through a BCMA-independent pathway, *Science* 293 (2001) 2111–2114.
- [17] L. Bossaller, J. Burger, R. Draeger, B. Grimbacher, R. Knoth, A. Plebani, A. Durandy, U. Baumann, M. Schlesier, A.A. Welcher, ICOS deficiency is associated with a severe reduction of CXCR5<sup>+</sup>CD4<sup>+</sup> germinal center Th cells, *J. Immunol.* 177 (2006) 4927–4932.
- [18] H. Hu, X. Wu, W. Jin, M. Chang, X. Cheng, S.C. Sun, Noncanonical NF- $\kappa$ B regulates inducible costimulator (ICOS) ligand expression and T follicular helper cell development, *Proc. Natl. Acad. Sci.* 108 (2011) 12827–12832.
- [19] K. Lüthje, A. Kallies, Y. Shimohakamada, G.T. Belz, A. Light, D.M. Tarlinton, S.L. Nutt, The development and fate of follicular helper T cells defined by an IL-21 reporter mouse, *Nat. Immunol.* 13 (2012) 491–498.
- [20] D.T. Avery, E.K. Deenick, C.S. Ma, S. Suryani, N. Simpson, G.Y. Chew, T.D. Chan, U. Palendira, J. Bustamante, S. Boisson-Dupuis, B cell-intrinsic signaling through IL-21 receptor and STAT3 is required for establishing long-lived antibody responses in humans, *J. Exp. Med.* 207 (2010) 155–171.
- [21] J. Parrish-Novak, S.R. Dillon, A. Nelson, A. Hammond, C. Sprecher, J.A. Gross, J. Johnston, K. Madden, W. Xu, J. West, Interleukin-21 and its receptor are involved in NK cell expansion and regulation of lymphocyte function, *Nature* 408 (2000) 57–63.
- [22] J. Pène, J.F. Gauchat, S. Lécart, E. Drouet, P. Guglielmi, V. Boulay, A. Delwail, D. Foster, J.C. Lecron, H. Yssel, Cutting edge: IL-21 is a switch factor for the production of IgG1 and IgG3 by human B cells, *J. Immunol.* 172 (2004) 5154–5157.
- [23] I.S. Grewal, R.A. Flavell, CD40 and CD154 in cell-mediated immunity, *Annu. Rev. Immunol.* 16 (1998) 111–135.
- [24] R.J. Armitage, B.M. Macduff, M.K. Spriggs, W.C. Fanslow, Human B cell proliferation and Ig secretion induced by recombinant CD40 ligand are modulated by soluble cytokines, *J. Immunol.* 150 (1993) 3671–3680.
- [25] N.S. Butler, J. Moebius, L.L. Pewe, B. Traore, O.K. Doumbo, L.T. Tygrett, T.J. Waldschmidt, P.D. Crompton, J.T. Harty, Therapeutic blockade of PD-L1 and LAG-3 rapidly clears established blood-stage Plasmodium infection, *Nat. Immunol.* 13 (2012) 188–195.
- [26] K.L. Good-Jacobson, C.G. Szumilas, L. Chen, A.H. Sharpe, M.M. Tomayko, M.J. Shlomchik, PD-1 regulates germinal center B cell survival and the formation and affinity of long-lived plasma cells, *Nat. Immunol.* 11 (2010) 535–542.
- [27] C.S. Ma, E.K. Deenick, M. Batten, S.G. Tangye, The origins, function, and regulation of T follicular helper cells, *J. Exp. Med.* 209 (7) (2012) 1241–1253.
- [28] S. Crotty, Follicular helper CD4 T cells (TFH), *Annu. Rev. Immunol.* 29 (2011) 621–663.
- [29] J.L. Cannons, K.T. Lu, P.L. Schwartzberg, T follicular helper cell diversity and plasticity, *Trends Immunol.* 34 (5) (2013) 200–207.
- [30] R.J. Johnston, A.C. Poholek, D. DiToro, I. Yusuf, D. Eto, B. Barnett, A.L. Dent, J. Craft, S. Crotty, Bcl6 and Blimp-1 are reciprocal and antagonistic regulators of T follicular helper cell differentiation, *Science* 325 (2009) 1006–1010.
- [31] Y.M. Kerdiles, E.L. Stone, D.R. Beisner, M.A. McGargill, I.L. Ch'en, C. Stockmann, C.D. Katayama, S.M. Hedrick, Foxo transcription factors control regulatory T cell development and function, *Immunity* 33 (2010) 890–904.
- [32] H. Wang, M. Yan, J. Sun, S. Jain, R. Yoshimi, M. Abolfath, K. Ozato, G. Coleman Jr., P. Ng, D. Metcalf, et al., A reporter mouse reveals lineage-specific and heterogeneous expression of IRF8 during lymphoid and myeloid cell differentiation, *Immunol.* 193 (4) (2014) 1766–1777 15.
- [33] Kenya Honda, Tadatsugu Taniguchi, IRFs: master regulators of signalling by Toll-like receptors and cytosolic pattern-recognition receptors, *Nat. Rev. Immunol.* 6 (2006) 644–658.
- [34] Michael Lohoff, W. Tak, Roles of interferon-regulatory factors in T-helper-cell differentiation, *Nat. Rev. Immunol.* 5 (2005) 125–135.
- [35] H.S. Wang, C.H. Lee, C.F. Qi, P. Taylor, J. Feng, S. Abbasi, T. Atsumi, H.C. Morse 3rd, IRF8 regulates B-cell lineage specification, commitment, and differentiation, *Blood* 112 (10) (2008) 4028–4038.
- [36] A. Takaoka, T. Tamura, T. Taniguchi, Interferon regulatory factor family of transcription factors and regulation of oncogenesis, *Cancer Sci.* 99 (3) (2008) 467–478.
- [37] Y. Yoshida, R. Yoshimi, H. Yoshii, D. Kim, A. Dey, H. Xiong, J. Munasinghe, I. Yazawa, M.J. O'Donovan, O.A. Maximova, et al., The transcription factor IRF8 activates integrin-mediated TGF- $\beta$  signaling and promotes neuroinflammation, *Immunity* 40 (20) (2014) 187–198.
- [38] J.X. Feng, H.S. Wang, D.M. Shin, M. Masiuk, C.F. Qi, H.C. Morse 3rd, IFN regulatory factor 8 restricts the size of the marginal zone and follicular B cell pools, *J. Immunol.* 186 (2011) 1458–1466.
- [39] S. Hambleton, S. Salem, J. Bustamante, V. Bigley, S. Boisson-Dupuis, J. Azevedo, A. Fortin, M. Haniffa, L. Ceron-Gutierrez, C.M. Bacon, IRF8 mutations and human dendritic-cell immunodeficiency, *N. Engl. J. Med.* 365 (2) (2011) 127–138 10.
- [40] W. Huang, E. Horvath, E.A. Eklund, P.U.1, interferon regulatory factor (IRF) 2, and the interferon consensus sequence-binding protein (ICSBP/IRF8) cooperate to activate NF1 transcription in differentiating myeloid cells, *J. Biol. Chem.* 282 (9) (2007) 6629–6643.
- [41] Elke Glasmacher, Smita Agrawal, Abraham B. Chang, Theresa L. Murphy, Wenwen Zeng, Bryan Vander Lugt, Aly A. Khan, Maria Ciofani, Chauncey J. Spooner, Sascha Rutz, Jason Hackney, Roza Nurieva, Carlos R. Escalante, Wenjun Ouyang, Dan R. Littman, Kenneth M. Murphy, Harinder Singh, A genomic regulatory element that directs assembly and function of immune-specific AP-1-IRF complexes, *Science* 338 (16) (2012) 975–981.
- [42] X. Ouyang, R. Zhang, J. Yang, Q. Li, L. Qin, C. Zhu, J. Liu, H. Ning, M.S. Shin, M. Gupta, C.F. Qi, J.C. He, S.A. Lira, H.C. Morse 3rd, K. Ozato, L. Mayer, H. Xiong, Transcription factor IRF8 directs a silencing programme for TH17 cell differentiation, *Nat. Commun.* 2 (2011) 314, <https://doi.org/10.1038/ncomms1311>.
- [43] Stacey N. Harboure, Craig L. Maynarda, Carlene L. Zindla, Trenton R. Schoeb, Casey T. Weaver, Th17 cells give rise to Th1 cells that are required for the pathogenesis of colitis, *Proc. Natl. Acad. Sci.* 112 (22) (2015) 7061–7066.
- [44] N. Simpson, Expansion of circulating T cells resembling follicular helper T cells is a fixed phenotype that identifies a subset of severe systemic lupus erythematosus, *Arthritis Rheum.* 62 (2010) 234–244.
- [45] J.Y. Choi, J.H. Ho, S.G. Pasoto, V. Bunin, S.T. Kim, S. Carrasco, E.F. Borba, C.R. Gonçalves, P.R. Costa, E.G. Kallas, E. Bonfa, J. Craft, Circulating follicular helper-like T cells in systemic lupus erythematosus: association with disease activity, *Arthritis Rheum.* 67 (4) (2015) 988–999.
- [46] V.M. Liarski, N. Kaverina, A. Chang, D. Brandt, D. Yanez, L. Talasnik, G. Carlesso, R. Herbst, T.O. Utset, C. Labno, Y. Peng, Y. Jiang, M.L. Giger, M.R. Clark, Cell distance mapping identifies functional T follicular helper cells in inflamed human renal tissue, *Sci. Transl. Med.* 6 (230) (2014) 230–246 2.
- [47] N. Schmitt, Role of T Follicular helper cells in multiple sclerosis, *J. Nat. Sci.* 1 (7) (2015) 139–148.
- [48] R. Kenefack, C.J. Wang, T. Kapadi, L. Wardzinski, K. Attridge, L.E. Clough, F. Heuts, A. Kogmitzis, S. Patel, M. Rosenthal, Follicular helper T cell signature in type 1 diabetes, *J. Clin. Invest.* 125 (1) (2015) 292–303.
- [49] M. Sarra, I. Monteleone, C. Stolfi, M.C. Fantini, P. Sileri, G. Sica, R. Tersigni, T.T. Macdonald, F. Pallone, G. Monteleone, Interferon-gamma-expressing cells are a major source of interleukin-21 in inflammatory bowel diseases, *Inflamm. Bowel Dis.* 16 (8) (2010) 1332–1339.
- [50] V. Gateva, A large-scale replication study identifies TNIP1, PRDM1, JAZF1, UHRF1BP1 and IL10 as risk loci for systemic lupus erythematosus, *Nat. Genet.* 41 (2009) 1228–1233.
- [51] P.L. De Jager, Meta-analysis of genome scans and replication identify CD6, IRF8 and TNFRSF1A as new multiple sclerosis susceptibility loci, *Nat. Genet.* 11 (4) (2011) 776–782.
- [52] H. Elding, W. Lau, D.M. Swallow, N. Maniatis, Dissecting the genetics of complex inheritance: linkage disequilibrium mapping provides insight into Crohn disease, *Am. J. Hum. Genet.* 89 (2011) 798–805.