

Corrigendum

Corrigendum to “Global burden of molar incisor hypomineralization”
[J. Dent. 68C (2018) 10–18]



Falk Schwendicke^{a,*}, Karim Elhennawy^a, Seif Reda^a, Katrin Bekes^b, David J. Manton^c,
Joachim Krois^a

^a Operative and Preventive Dentistry, Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Germany

^b Department of Paediatric Dentistry, School of Dentistry, Medical University of Vienna, Vienna, Austria

^c Paediatric Dentistry, Melbourne Dental School, University of Melbourne, Australia

The authors regret:


- A first author's surname was misspelled in Fig. 3 and Table S1; the author's name is Wuollet
- The data point Wuollet 2016 was incorrectly extracted, the sample size was 287, not 87. This affects the prevalence estimates, as follows:
- Mean (95% CI) prevalence changed for Finland (16.2, 12.7/20.3), the super-region “High Income” (14.5, 12.3/17.0), and globally (12.9, 11.7/14.3).
- This also affects the case number to a minor degree, except for Finland, where now 849,278 prevalent cases and 9498 incident cases are calculated.
- Corrected Figures 3 and 4 and Table 1 are attached. Note that a minor mistake in color coding of Fig. 4a was also corrected. Also note that further, albeit minimal changes in prevalent and incident case estimates in other, non-affected countries may have occurred by model parameters being minimally deviant from the original model given the altered input data.

DOI of original article: <https://doi.org/10.1016/j.jdent.2017.12.002>

* Corresponding author at: Charité Centre for Dental Medicine, Department for Operative and Preventive Dentistry, Aßmannshäuser Str. 4–6, 14197, Berlin, Germany.

E-mail address: falk.schwendicke@charite.de (F. Schwendicke).

<https://doi.org/10.1016/j.jdent.2018.11.006>






Table 1: Prevalent cases in 2015 and incident cases in 2016 on super-regional, regional and regional levels. Details to each country can be found in the appendix Table S4.

GBD superregion	Prevalent Cases	Incident Cases	GBD region	Prevalent Cases	Incident Cases
Central Europe, Eastern Europe, and Central Asia	51.266.325	701.562	Central Asia	9.524.526	220.931
			Central Europe	14.994.426	162.588
High-income	191.396.872	2.354.845	Eastern Europe	26.747.373	318.043
			Australasia	5.693.232	76.052
			High-income Asia Pacific	20.802.458	186.487
			High-income North America	97.115.983	1.294.819
			Southern Latin America	8.129.550	137.760
			Western Europe	59.655.649	659.727
Latin America and Caribbean	78.735.493	1.482.305	Andean Latin America	7.239.745	162.323
			Caribbean	5.705.559	107.058
			Central Latin America	35.247.762	721.087
			Tropical Latin America	30.542.427	491.837
			North Africa and Middle East	61.918.666	1.533.740
North Africa and Middle East	61.918.666	1.533.740	South Asia	151.411.432	3.459.825
South Asia	151.411.432	3.459.825	East Asia	98.674.522	1.235.827
Southeast Asia, East Asia, and Oceania	180.723.796	2.866.638	Oceania	1.131.557	32.889
Sub-Saharan Africa	95.673.601	3.634.098	Southeast Asia	80.917.717	1.597.922
			Central Sub-Saharan Africa	11.292.692	480.396
			Eastern Sub-Saharan Africa	40.036.460	1.526.477
			Southern Sub-Saharan Africa	8.875.507	224.277
			Western Sub-Saharan Africa	35.468.942	1.402.948

The authors would like to apologise for any inconvenience caused.