



Mortality of acute-on-chronic liver failure: What is the role of obesity?

To the Editors:

I read with great interest the study by Hernaez *et al.*, entitled “Prevalence and short-term mortality of acute-on-chronic liver failure: A national cohort study from the USA.”¹ In their paper, the authors illustrate several notable findings regarding risk factors for mortality in patients with acute-on-chronic liver failure (ACLF) via analysis of the Veterans Affairs’ database.

In our previous paper, we demonstrated that class III obesity is a risk factor for the development of ACLF in patients with decompensated cirrhosis by analyzing the United network for Organ Sharing (UNOS) database and the Nationwide Inpatient Sample.² Unfortunately due to biases and limitations in both databases, we could not evaluate whether obesity was a risk factor for mortality in patients who had developed the ACLF. In the current study, the authors demonstrate that patients with ACLF have a greater prevalence of obesity based on body mass index (47% vs. 40%), compared to those without ACLF. Furthermore, among each grade of ACLF, the majority of patients had a BMI above 30 (41%–49%). These findings do suggest, at least, that obesity is more prevalent among patients with ACLF.

I am interested to know if the authors have performed or can perform an analysis regarding whether BMI, either as a continuous or categorical variable, was associated with mortality among patients who developed ACLF, particularly with regards to class III obesity. The results, whether positive or negative, would be clinically useful for the risk stratification of our patients with ACLF.

Conflict of interest

The author declares no conflicts of interest that pertain to this work.

Please refer to the accompanying ICMJE disclosure forms for further details.

Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jhep.2019.01.018>.

References

- [1] Hernaez R, Kramer JR, Liu Y, Tansel A, Natarajan Y, Hussain KB, et al. Prevalence and short-term mortality of acute-on-chronic liver failure: a national cohort study from the USA. *J Hepatol* 2019;70:639–647.
- [2] Sundaram V, Jalan R, Ahn JC, Charlton MR, Goldberg DS, Karvellas CJ, et al. Class III obesity is a risk factor for the development of acute-on-chronic liver failure in patients with decompensated cirrhosis. *J Hepatol* 2018;69:617–625.

Vinay Sundaram*

Division of Gastroenterology and Comprehensive Transplant Center,
Cedars-Sinai Medical Center, Los Angeles, CA, United States

*Address: 8900 Beverly Blvd, Los Angeles, CA 90048, United States.
Tel.: +1 310 423 6000, fax: +1 310 423 2356.
E-mail address: Vinay.sundaram@cshs.org



Reply to: “Mortality of acute-on-chronic liver failure: What is the role of obesity?”

To the Editors:

We appreciate the interest and comments by Dr. Sundaram. Out of 19,082 patients with acute-on-chronic liver failure (ACLF) in our study, 17.4% had missing data on body mass index (BMI).¹ Amongst those with non-missing data, obesity class III was present in 10.3%, obesity I-II in 36.4% and 35.9% were non-obese (BMI <30 kg/m²). The 28-day mortality rates were 20.5% in patients with class III obesity, 22.9% in those with class II obesity and 28.2% in non-obese patients. We further examined the independent association between mortality and BMI (as continuous and categorical variables). BMI was inversely associated with 28-day mortality (odds ratio [OR] 0.98; 95% CI 0.97–0.98) after adjustment of other demographic and clinical factors. We found a similar association when we examined obesity as a categorical variable. Using obesity classes, and compared to non-obese patients, both obesity class I-II and obesity class III were inversely associated with mortality risk (OR for class I-II 0.73; 95% CI 0.67–0.79; OR for class III 0.62; 95% CI 0.55–0.71).

In summary, despite the high prevalence of obesity in patients with ACLF,² these additional analyses show that BMI was inversely associated with mortality in our cohort. The higher risk of mortality observed in normal (and underweight) ACLF patients could at least partly be caused by residual confounding from malnutrition and sarcopenia – both markers of poor prognosis in decompensated cirrhosis. Future cohort studies that prospectively collect data on BMI, anthropometrics, nutrition, and sarcopenia will be important to examine the association between BMI and mortality in ACLF.

Financial support

The work is also supported in part by the Center for Gastrointestinal Development, Infection and Injury (NIDDK P30 DK 56338). Drs. Hernaez, Kramer and Kanwal are investigators at the Center for Innovations in Quality, Effectiveness and Safety (CIN 13-413), Michael E. DeBakey VA Medical Center, Houston, TX. This material is based upon work supported (or supported