

Conflict of interest

The authors declare no conflicts of interest that pertain to this work.

Please refer to the accompanying ICMJE disclosure forms for further details.

Authors' contributions

AS, PM & PD wrote the manuscript and revised the text.

Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jhep.2019.03.019>.

References

Author names in bold designate shared co-first authorship

- [1] Hessheimer A, Coll E, Torres F, Ruiz P, Gastaca M, Rivas J, et al. Normothermic regional perfusion versus super rapid recovery in controlled donation after circulatory death liver transplantation. *J Hepatol* 2019;70:658–665. <https://doi.org/10.1016/j.jhep.2018.12.013>.
- [2] Schlegel A, Kalisvaart M, Scalera I, Laing R, Mergental H, Mirza D, et al. The UK DCD Risk Score: a new proposal to define futility in donation-after-circulatory-death liver transplantation. *J Hepatol* 2018;456–464. <https://doi.org/10.1016/j.jhep.2017.10.034>.
- [3] DeOliveira ML, Jassem W, Valente R, Khorsandi SE, Santori G, Prachalias A, et al. Biliary complications after liver transplantation using grafts from donors after cardiac death: results from a matched control study in a single large volume center. *Ann Surg* 2011;254:716–722. <https://doi.org/10.1097/SLA.0b013e318235c572>, discussion 722–3.
- [4] Laing RW, Scalera I, Isaac J, Mergental H, Mirza DF, Hodson J, et al. Liver transplantation using grafts from donors after circulatory death: A propensity-matched study from a single centre. *Am J Transplant* 2016. <https://doi.org/10.1111/ajt.13699>.

- [5] Davila D, Ciria R, Jassem W, Briceño J, Littlejohn W, Vilca-Meléndez H, et al. Prediction models of donor arrest and graft utilization in liver transplantation from maastricht-3 donors after circulatory death. *Am J Transplant* 2012. <https://doi.org/10.1111/j.1600-6143.2012.04242.x>.
- [6] Muller X, Schlegel A, Würdinger M, Wendt M, Kron P, Eshmunov D, et al. Can hypothermic oxygenated perfusion (HOPE) rescue futile DCD liver grafts?. *HPB (Oxford)* 2019. <https://doi.org/10.1016/j.hpb.2019.01.004>, pii: S1365-182X(19)30058-9.
- [7] Mergental H, Laing R, Kirkham A, Perera T, Boteon Y, Attard J, et al. Transplantation of discarded livers following viability testing with normothermic machine perfusion: the VITAL (Viability Testing and Transplantation of mArginal Livers) trial outcomes. *AASLD 2018;2018. Abstract*.
- [8] Dutkowski P, Guarrera J, De Jonge J, Martins P, Porte R, Clavien P. Evolving trends in machine perfusion for liver transplantation. *Gastroenterology* 2019. <https://doi.org/10.1053/j.gastro.2018.12.037>.
- [9] Watson C, Kosmoliaptis V, Pley C, Randle L, Fear C, Crick K, et al. Observations on the ex situ perfusion of livers for transplantation. *Am J Transplant* 2018. <https://doi.org/10.1111/ajt.14687>.
- [10] Chouchani ET, Pell VR, James AM, Work LM, Saeb-Parsy K, Frezza C, et al. A unifying mechanism for mitochondrial superoxide production during ischemia-reperfusion injury. *CellMetab* 2016;23:254–263. <https://doi.org/10.1016/j.cmet.2015.12.009>.

Andrea Schlegel^{1,2}

Paolo Muiesan¹

Philipp Dutkowski^{3,*}

¹Liver Unit, Queen Elizabeth Hospital Birmingham, University Hospitals Birmingham National Health Service Foundation Trust, United Kingdom

²National Institute for Health Research Birmingham, Liver Biomedical Research Centre, College of Medical and Dental Sciences, University of Birmingham, United Kingdom

³Department of Surgery and Transplantation, Swiss HPB Centre, University Hospital Zurich, Switzerland

*Corresponding author. Address: Department of Surgery and Transplantation, Swiss HPB Centre, University Hospital Zurich, Raemistrasse 100, CH-8091 Zurich, Switzerland; Tel.: +41 44 255 4236; fax: +41 44 255 4999. E-mail address: philipp.dutkowski@usz.ch



Reply to: “Normothermic regional perfusion – What is the benefit?”

To the Editor:

We thank Drs. Schlegel, Muiesan, and Dutkowski for their interest in our manuscript¹ and are delighted to provide clearer and updated information regarding the use of normothermic regional perfusion (NRP) in controlled donation after circulatory death (cDCD) liver transplantation.

Our manuscript describes the Spanish experience with cDCD liver transplantation from national application in 2012 through 2016, comparing outcomes of transplants performed with NRP versus those performed with super rapid recovery (SRR).² Transplants were included from 20 centers, only 3 (15%) with previous experience performing uncontrolled donation after circulatory death. The results that we present can be achieved by not just perfusion and DCD “experts” but by virtually any liver transplant team. That said, the results we achieved with SRR are the same as those described by experienced centers using cDCD livers of a comparable profile,^{3,4} and improvements

achieved with NRP are strikingly similar to those recently reported by 2 UK centers describing 43 cDCD liver transplants performed with NRP.⁵

The Letter's authors claim the “risk” of the cDCD grafts we used was low based on a scoring system they developed.⁶ The UK DCD Risk Score has not been independently validated but has found to be ineffective at predicting 1-year cDCD liver survival (its aim) in our recipients⁷ and other British cohorts.⁸ The authors do correctly describe our median functional donor warm ischemia times: 12 and 15 minutes when NRP and SRR were used, respectively. Femoral cannulae were placed prior to withdrawal of ventilatory support in 87% of cases using NRP, and warm ischemia times were shorter when NRP was employed. That said, the implication that the authors consistently experience longer warm ischemia times than even those for livers recovered with SRR is surprising. Only 11 of 342 cDCD donors considered for liver donation during the study period




Fig. 1. Liver transplant approaches in Spain. (A) Year-to-year evolution of cDCD liver transplant activity in Spain, divided according to organ recovery method: NRP, super rapid cold perfusion and SRR. (B) In the past 2 years, NRP has become the recovery method of choice in cDCD liver transplantation in Spain. In 2018, it was applied 217 times at 62 different hospitals. In the majority of these cases (62%), NRP was performed at non-liver transplant centers using either in-hospital or transported equipment. Source: Organización Nacional de Trasplantes (www.ont.es). cDCD, controlled donation after circulatory death; NRP, normothermic regional perfusion; SRR, super rapid recovery.

were turned down due to prolonged warm ischemia (3%). The facts that i) indication for proceeding with cDCD in Spain is strict and predicated on likelihood of arrest within 60 minutes of withdrawal of care and ii) cDCD donors in Spain have been in intensive care 7–10 days prior to withdrawal (vs. 2–3 days in the UK)^{3,5} may explain the consistently shorter warm ischemia times we experienced.

While we did not argue in our manuscript that our discard rates are similar to those in other countries, the authors are correct in pointing this out. Between 2012 and 2016, 38% of cDCD livers recovered with SRR and 34% recovered with NRP in Spain were ultimately discarded, similar to the 33% of retrieved cDCD livers declined in 2017/2018 in the UK.⁹ Our figures are national averages, and individual centers have lower discard rates. Obviously, the “rate” depends on the denominator, and comparing national averages with rates described in smaller pilot studies, where pre-selection has been performed on grafts included, is misleading.¹

As in donation after brain death (DBD) liver transplantation, cDCD liver evaluation is not exact, and visual assessment by the surgical team is still the ultimate measure of viability when *ex situ* perfusion is not employed. The majority of cDCD livers turned down in our study were observed to be moderately-to-severely steatotic, poorly perfused, fibrotic, or cirrhotic.² While hepatic aminotransferases in the perfusate were assessed as an indication of hepatic injury, they rose very little in most cases (only 4 livers were turned down due to rising aminotransferases), and liver viability assessment in the strictest sense was not performed. This does not mean that true viability assessment may not be performed during NRP, as bile is produced, and evaluation of bile production and biochemistry may serve as useful means to assess significant biliary injury.

When NRP is employed, the cost of the cDCD process increases €2,500–5,000 with respect to standard DBD or cDCD with SRR. This cost is lower than that required to perfuse a liver *ex situ*, where disposable components of the machine perfusion circuit alone are at the higher end of the aforementioned range. Considering costs, one also has to keep in mind that NRP is simultaneously used to recover cDCD kidneys, pancreata, and hearts, as well, and benefits of NRP in terms of post-transplantation outcomes extend to these organs.¹⁰

In their final comments, the Letter’s authors call for a moratorium on widespread application of any perfusion technology in human liver transplantation pending results of randomized trials. This recommendation defies reality in countries such as Spain, France, and Italy. In Spain, the use of NRP in cDCD has risen exponentially since 2012 (Fig. 1A). In 2018, 189 cDCD livers were transplanted: 151 with NRP and 38 with SRR. NRP was applied in >200 cDCD donors at 62 hospitals, in a third of cases at non-transplant hospitals by local teams (Fig. 1B). The application of NRP is widely disseminated here and has allowed for implementation of cDCD at all levels. Given excellent post-transplant results,² it seems improbable if not unethical that centers currently using NRP would abandon it and risk increased biliary complications and graft loss pending a level 1 clinical trial.

While NRP increases upfront costs associated with cDCD transplantation compared with SRR, it may be used by both transplant and perfusion experts and less experienced professionals to treat and potentially assess the quality of multiple organs, not just the liver. At a time when healthcare systems are concerned with achieving the greatest benefit at the lowest cost, NRP appears to be the DCD perfusion strategy that best meets this need.

Conflict of interest

The authors declare no conflicts of interest that pertain to this work.

Please refer to the accompanying ICMJE disclosure forms for further details.

Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jhep.2019.04.005>.

References

Author names in bold designate shared co-first authorship

- [1] Schlegel A, Muiesan P, Dutkowski P. Normothermic regional perfusion – what is the benefit? *J Hepatol* 2019;71:441–443.

- [2] Hessheimer AJ, Coll E, Torres F, Ruiz P, Gastaca M, Rivas JI, et al. Normothermic regional perfusion vs. super-rapid recovery in controlled donation after circulatory death liver transplantation. *J Hepatol* 2019;70 (4):658–665.
- [3] Schlegel A, Scalera I, Perera MTPR, Kalisvaart M, Mergental H, Mirza DF, et al. Impact of donor age in donation after circulatory death liver transplantation: Is the cutoff “60” still of relevance? *Liver Transpl* 2018;24:352–362.
- [4] Croome KP, Mathur AK, Lee DD, Moss AA, Rosen CB, Heimbach JK, et al. Outcomes of donation after cardiac death liver grafts from donors $\geq$ 50 years of age: a multi-center analysis. *Transplantation* 2018.
- [5] Watson C, Hunt F, Messer S, Currie I, Large S, Sutherland A, et al. In situ normothermic perfusion of livers in controlled circulatory death donation may prevent ischemic cholangiopathy and improve graft survival. *Am J Transplant* 2019.
- [6] Schlegel A, Kalisvaart M, Scalera I, Laing RW, Mergental H, Mirza DF, et al. The UK DCD risk score: a new proposal to define futility in donation-after-circulatory-death liver transplantation. *J Hepatol* 2018;68:456–464.
- [7] Hessheimer AJ, Coll E, Ruiz P, Gastaca M, Rivas JI, Gomez M, et al. The UK DCD risk score: still no consensus on futility in DCD liver transplantation. *J Hepatol* 2019;16.
- [8] Oniscu GC, Watson CJE, Wigmore SJ. Redefining futility in DCD liver transplantation in the era of novel perfusion technologies. *J Hepatol* 2018;68:1327–1328.
- [9] UK NHS Blood and Transplant. 2017/18 Organ Donation and Transplantation Activity Report. British Transplantation Society; 2018 Apr 1.
- [10] Demiselle J, Augusto JF, Videcoq M, Legeard E, Dube L, Templier F, et al. Transplantation of kidneys from uncontrolled donation after circulatory determination of death: comparison with brain death donors with or without extended criteria and impact of normothermic regional perfusion. *Transpl Int* 2016;29:432–442.

Amelia J. Hessheimer¹
 Elisabeth Coll²
 Patricia Ruíz³
 Mikel Gastaca³
 José Ignacio Rivas⁴
 Manuel Gómez⁴
 Belinda Sánchez⁵
 Julio Santoyo⁵
 Pablo Ramírez⁶
 Pascual Parrilla⁶
 Luis Miguel Marín⁷
 Miguel Ángel Gómez-Bravo⁷
 Juan Carlos García-Valdecasas¹
 Javier López-Monclús⁸
 Andrea Bosca⁹
 Rafael López-Andújar⁹
 Yiliam Fundora-Suárez¹⁰
 Jesús Villar¹⁰
 Álvaro García-Sesma¹¹

Carlos Jiménez¹¹
 Gonzalo Rodríguez-Laíz¹²
 Laura Lladó¹³
 Juan Carlos Rodríguez¹⁴
 Manuel Barrera¹⁵
 Ramón Charco¹⁶
 Jose Ángel López-Baena¹⁷
 Javier Briceño¹⁸
 Fernando Pardo¹⁹
 Gerardo Blanco²⁰
 David Pacheco²¹
 Beatriz Domínguez-Gil²
 Víctor Sánchez Turrión⁸
 Constantino Fondevila^{1,*}

¹Department of General & Digestive Surgery, Institut de Malalties Digestives i Metabòliques (IMDiM), Hospital Clínic, CIBERehd, IDIBAPS, University of Barcelona, Spain

²Organización Nacional de Trasplantes, Madrid, Spain

³Hospital Universitario Cruces, Bilbao, Spain

⁴Complejo Hospitalario Universitario La Coruña, Spain

⁵Hospital Regional Universitario de Málaga, Spain

⁶Hospital Clínico Universitario Virgen de la Arrixaca (IMIB), Murcia, Spain

⁷Hospital Universitario Virgen del Rocío, Seville, Spain

⁸Hospital Universitario Puerta de Hierro, Majadahonda, Spain

⁹Hospital Universitario y Politécnico La Fe, Valencia, Spain

¹⁰Hospital Universitario Virgen de las Nieves, Granada, Spain

¹¹Hospital Universitario 12 de Octubre, Madrid, Spain

¹²Department of General & Digestive Surgery, ISABIAL, Hospital General Universitario de Alicante, Spain

¹³Hospital Universitario de Bellvitge, Hospitalet de Llobregat, Spain

¹⁴Hospital Universitario Marqués de Valdecilla, Santander, Spain

¹⁵Hospital Universitario Nuestra Señora de Candelaria, Santa Cruz de Tenerife, Spain

¹⁶Hospital Universitario Vall d'Hebrón, Barcelona, Spain

¹⁷Hospital General Universitario Gregorio Marañón, Madrid, Spain

¹⁸Hospital Universitario Reina Sofía, Córdoba, Spain

¹⁹Clínica Universitaria de Navarra, Pamplona, Spain

²⁰Hospital Universitario Infanta Cristina, Badajoz, Spain

²¹Hospital Universitario Río Hortega, Valladolid, Spain

*Corresponding author. Address: Chief of the Department of General & Digestive Surgery, Hospital Clínic, University of Barcelona, C/ Villarreal 170, 08036 Barcelona, Spain. Tel.: +34 93 227 57 18, fax +34 93 227 55 89.

E-mail address: cfonde@clinic.cat