



ELSEVIER

Available online at www.sciencedirect.com

Journal of Hospital Infection

journal homepage: www.elsevier.com/locate/jhin



Preoperative mechanical and oral antibiotic bowel preparation to reduce infectious complications of colorectal surgery – the need for updated guidelines

C.L.F. Battersby^{a,b,*}, N.J. Battersby^c, D.A.J. Slade^b, M. Soop^b, C.J. Walsh^d

^a Wrexham Maelor Hospital, Wrexham, UK

^b Salford Royal Foundation Trust, Salford, UK

^c Wessex School of Surgery, Winchester, Hampshire, UK

^d Wirral University Teaching Hospital NHS Foundation Trust, Birkenhead, Wirral, UK

ARTICLE INFO

Article history:

Received 3 November 2018

Accepted 14 December 2018

Available online 21 December 2018

Keywords:

Bowel preparation

Complications

Surgical site infection



SUMMARY

Background: Increasing evidence indicates that combined mechanical and oral antibiotic bowel preparation reduces the infectious complications of colorectal surgery. Anecdotal evidence suggests the combination is rarely used in the UK and Europe.

Aim: To establish colorectal surgeons' current use and awareness of the benefits of such bowel preparation, and to identify decision-making influences surrounding preoperative bowel preparation.

Method: An electronic survey was emailed to all members of the Association of Coloproctology of Great Britain and Ireland, and promoted via Twitter.

Findings: A total of 495 respondents completed the survey: 413 (83.2%) UK, 39 (7.9%) other European, 43 (8.7%) non-European. Respondents used oral antibiotics for 12–20% of cases. Mechanical bowel preparation (MBP), phosphate enema, and no preparation, respectively, ranged between 9 and 80%. Combined MBP and oral antibiotic bowel preparation ranged between 5.5 and 18.6%.

Fifty-three percent (260/495) agreed that combined mechanical and oral antibiotic bowel preparation reduces surgical site infection; 32% (157/495) agreed that the combination reduces risk of anastomotic leak.

Kappa statistics between 0.06 and 0.27 indicate considerable incongruity between surgeons' awareness of the literature, and day-to-day practice.

Twenty-four percent (96/495) believed MBP to be incompatible with enhanced recovery after surgery (ERAS); 41% (204/495) believed that MBP delays return to normal intestinal function.

Conclusions: Few UK and European colorectal surgeons use mechanical and oral antibiotic bowel preparation, despite evidence of its efficacy in reducing infectious complications. The influence of ERAS pathways and UK and European guidelines may explain this. In contradiction to the UK and Europe, North American guidelines recommend incorporating combined mechanical and oral antibiotic bowel preparation into ERAS programmes. This

* Corresponding author. Address: Department of General Surgery, Wrexham Maelor Hospital, Croesnewydd Road, Wrexham LL13 7TD, UK. Tel.: +44 7855 946766.

E-mail address: clfbattersby@doctors.org.uk (C.L.F. Battersby).

study suggests that future UK and European guidelines incorporate combined mechanical and oral antibiotic bowel preparation into the ERAS pathway.

© 2018 The Healthcare Infection Society. Published by Elsevier Ltd. All rights reserved.

Introduction

Infectious complications of colorectal surgery continue to present a significant burden to both patients and healthcare providers. Complications such as surgical site infection (SSI) and anastomotic leak (AL) cause considerable morbidity and mortality, cost, prolonged length of stay (LOS) and impaired quality of life (QOL) [1,2]. Colorectal surgery accounts for the greatest number, and most costly SSIs within NHS hospitals in England [3]. Despite the scale of the problem, recording and reporting of SSIs is currently very poor; to reduce the incidence of this problem requires recognition of the nature of the problem, and clinical leadership to drive change [3,4]. It is well recognized that rates of SSI can be significantly reduced using infection-control bundles. Combined mechanical bowel preparation and preoperative oral antibiotic preparation (MBP/OABP) are common in such bundles [5–7].

Establishing an optimal bowel preparation regime to reduce the incidence and burden of infectious complications in colorectal surgery has been the subject of debate for over a century [8]. The use of mechanical bowel preparation (MBP), oral antibiotic bowel preparation (OABP), or the combination of both, has fallen in and out of favour over the decades [8].

The use of combined MBP/OABP in elective colorectal surgery remains the subject of considerable debate, reflected in the discrepancy observed between guidelines issued by various authorities around the world. NICE Guidelines (Clinical Guideline 74) recommend against the routine use of MBP, citing the 2011 Cochrane Review; these are currently the only published guidelines pertinent to UK practice [9]. European Society for Coloproctology (ESCP) guidelines also advise against the use of MBP [10].

There has been an increasing volume of recent evidence to suggest that the preoperative use of combined MBP/OABP is associated with significant reduction in the incidence of infectious complications of elective colorectal surgery [8,11–16]. The use of combined MBP/OABP is supported by randomized controlled trial data, especially for reducing SSI, however, much of the support for the use of combined MBP/OABP is from large North American cohort studies, including American College of Surgery National Surgical Quality Improvement Program (ACS NSQIP) data [13].

In contrast to UK and European guidelines, North American guidelines recommend routine use of combined MBP/OABP for elective colorectal surgery [17]. One reason for this discrepancy is that some European authors are critical of the use of NSQIP data as studies are based on retrospective analysis of prospectively collected data, rather than data collected within a randomized trial setting and all of the studies are based on the same database. Due to the nature of the data collection, it is not always possible to establish the MBP/OABP regime employed. It is also the case that there is insufficient data comparing oral antibiotic preparation alone, with combined MBP and OAB [10].

The benefits of combined MBP/OABP is further supported by the recent ESCP Left Sided Colorectal Resection audit [18]. This was a prospectively designed study, with AL as a primary endpoint, and detailed data regarding nature of preparation, and other risk factors for AL were collected. Of 3676 patients from 52 countries, 618 (16.8%) received MBP & OABP, 1945 MBP (52.9%) and 1099 patients were in the NBP group (received no MBP and no OABP) (29.9%). Patients undergoing MBP & OABP had the lowest overall rate of AL (6.1%, 9.2%, 8.7%, respectively) and by mixed-effects multivariable regression MBP & OABP was associated with a two-fold reduction in the AL (odds ratio (OR) 0.52, 0.30–0.92, $P = 0.02$) but MBP alone did not reduce AL risk (OR 0.92, 0.63–1.36, $P = 0.69$) compared to NBP [18]. A randomized controlled trial, COLONPREP, designed to compare MBP/OABP with OABP alone is due to commence recruitment in the near future. The trial has SSI as a primary endpoint, with AL as a secondary endpoint. The outcomes may be informative with regard to preventing SSI, but it is unlikely that the trial will be adequately powered to evaluate the effect of preparation regime in reducing risk of AL.

Prior to the introduction of enhanced recovery after surgery (ERAS) pathways, MBP was regarded as standard practice. As ERAS has become increasingly popular, use of MBP has reduced, on the basis that the physiological response to the depletion of fluids and electrolytes prolongs postoperative recovery, MBP is unpleasant for the patients, and the lack of data to demonstrate benefits from its use [9,14]. The evidence, however, suggests that combined MBP/OABP does not delay time to discharge, and may even reduce LOS [13]. Oral antibiotic use alone has been investigated as an alternative to MBP, with some data to show reduced SSI and AL rates [11,19].

Despite the evidence to support the use of MBP/OABP in reducing infectious complications of elective colorectal surgery, anecdotal evidence suggests that the use of MBP and OABP regimes amongst colorectal surgeons is variable. This study aimed to survey current practice within the UK, and to extend the reach of the survey to the wider colorectal surgical community using social media. In addition to describing current practice, this work also sought to investigate factors that influence decision-making regarding use of MBP and OABP, especially within the setting of ERAS, which has been widely adopted by colorectal surgeons throughout the world.

Methods

An electronic survey was emailed to all members of the Association of Coloproctology of Great Britain and Ireland (ACPGBI). The survey was in English and the data collection period lasted for one month (19 March 2018 to 18 April 2018).

Promotional links were sent via Twitter in the week prior to launching the survey, on the day of the launch and regularly during the survey, with the #colorectalsurgery link, to disseminate the survey to a wider audience. Regular reminders were also sent via e-mail, the ACPGBI newsletter and Twitter.

Questions assessed use of MBP and OAB for various colorectal procedures, nature of respondent's practice (consultant or trainee), use of ERAS pathways, and opinions on the influence of MBP and OAB on the incidence of SSI and AL.

The strength of agreement between reported current practice and opinions regarding SSI and AL in relation to the use of MBP and OABP was tested using Kappa statistic. A Kappa value of ≤ 0.20 was interpreted as 'Poor', 0.21–0.40 as 'Fair', 0.41–0.60 as 'Moderate', 0.61–0.80 as 'Good', and 0.81–1.00 as 'Very good' [20]. Data analysis was with carried out with SPSS (v21.0; IBM Corp, Armonk, NY).

Results

A total of 495 respondents completed the survey, with 33 countries represented. Four-hundred and thirteen (83.2%) were from the UK, 39 (7.9%) from other European countries and 43 (8.7%) from non-European countries. Four-hundred and forty-three (89%) respondents were consultant colorectal surgeons; 52 (11%) were trainees with an interest in colorectal surgery. One-hundred and twenty-six (26%) have been practising ERAS for less than five years, 247 (50%) for five to 10 years, and 122 (24%) for over 10 years.

Oral antibiotics and MBP

Use of MBP, phosphate enemas and oral antibiotics for various colorectal procedures is detailed in Table I. Use of MBP was predominantly seen in left-sided resections, especially those with a planned defunctioning stoma, but remained low for other procedures. Between 30 and 47% of respondents used phosphate enemas as a substitute for MBP in left-sided procedures. The use of oral antibiotics alone was generally low for all procedures; consequently, the use of combined MBP/OABP was also low for all procedures.

Fifty-three percent (260/495) agreed that combined MBP/OAB reduces SSI; 32% (157/495) agreed that combined MBP and OAB reduces risk of AL.

Opinion regarding effect of MBP and OAB on SSI and AL

Kappa statistics are given for respondents' agreements with the statements regarding effect of OABP (Table II) and MBP (Table III) on the reduction of SSI and AL, and what those

Table II

Comparison between the strength of agreement of surgeons' reported current oral antibiotic bowel preparation practice and their view on the role of combined mechanical and oral antibiotic bowel preparation for preventing surgical site infection and anastomotic leak

MBP with oral antibiotics reduces risk of surgical site infection?				
	Current practice	Agrees with question	Disagrees with question	Kappa agreement
Right hemicolectomy	Use OABP	10.7%	1.4%	0.17
	Do not use OABP	41.8%	46.1%	
Left-sided resection without stoma	Use OABP	15.2%	2.6%	0.23
	Do not use OABP	37.4%	44.8%	
Left-sided resection with stoma	Use OABP	16.0%	4.4%	0.20
	Do not use OABP	36.6%	43.0%	
Panproctocolectomy	Use OABP	11.9%	2.4%	0.17
	Do not use OABP	40.6%	45.1%	
Abdominoperineal excision	Use OABP	12.3%	2.4%	0.18
	Do not use OABP	40.2%	45.1%	
MBP with oral antibiotics reduces risk of anastomotic leak?				
		Agrees with question	Disagrees with question	Kappa agreement
Right hemicolectomy	Use OABP	7.1%	5.1%	0.18
	Do not use OABP	24.6%	63.2%	
Left-sided resection without stoma	Use OABP	9.3%	8.5%	0.19
	Do not use OABP	22.4%	59.8%	
Left-sided resection with stoma	Use OABP	9.7%	10.7%	0.16
	Do not use OABP	22.0%	57.6%	
Panproctocolectomy	Use OABP	7.5%	6.9%	0.16
	Do not use OABP	24.2%	61.4%	
Abdominoperineal excision	Use OABP	7.7%	7.1%	0.16
	Do not use OABP	24.0%	61.2%	

These are listed according to operation type. Green box: current practice is consistent with the response to the question. Blue box: agrees with question but not in practice. Red box, surgeon uses OABP but not for this reason. MBP, mechanical bowel preparation; OABP, oral antibiotic bowel preparation.

respondents did in their own practice. The Kappa values are below 0.21 for OABP and MBP practice for all procedures except left-sided resection without a stoma. For left-sided procedures without a stoma the Kappa values between 0.23 and 0.27 indicate a fair agreement between the surgeon's opinion and practice.

MBP and ERAS

Twenty-four percent (96/495) believed MBP to be incompatible with ERAS; 41% of respondents (204/495) believed that MBP delays return to normal intestinal function.

Table I

Summary of reported bowel preparation practice for elective colorectal resections according to the survey responses

Operation type % (N/495)	MBP		Phosphate enema		No preparation		OABP		Combined MBP/OABP	
Right hemicolectomy	10%	[48/495]	9%	[47/495]	81%	[400/495]	12%	[60/495]	5.4%	[27/495]
Left-sided resection ^a without a planned defunctioning stoma	43%	[211/495]	47%	[235/495]	10%	[49/495]	18%	[88/495]	13.3%	[66/495]
Left sided resection ^a with a planned defunctioning stoma	80%	[395/495]	16%	[81/495]	4%	[19/495]	20%	[101/495]	18.6%	[92/495]
Panproctocolectomy	18%	[90/495]	30%	[149/495]	52%	[256/495]	14%	[71/495]	7.1%	[35/495]
Abdominoperineal excision resection	18%	[90/495]	43%	[211/495]	39%	[194/495]	15%	[73/495]	7.9%	[39/495]

MBP, mechanical bowel preparation; OABP, oral antibiotic preparation. MBP, mechanical bowel preparation.

^a Includes (low) anterior resections.

Table III

Comparison between the strength of agreement of surgeons' reported current mechanical bowel preparation practice and their view on the role of combined mechanical and oral antibiotic bowel preparation for preventing surgical site infection and anastomotic leak

MBP with oral antibiotics reduces risk of SSI?				
	Current practice	Agrees with question	Disagrees with question	Kappa agreement
Right hemicolectomy	Use MBP	7.5%	2.2%	0.09
	Do not use MBP	45.1%	45.3%	
Left-sided resection without stoma	Use MBP	29.3%	13.3%	0.27
	Do not use MBP	23.2%	34.1%	
Left-sided resection with stoma	Use MBP	44.2%	35.6%	0.10
	Do not use MBP	8.3%	11.9%	
Panproctocolectomy	Use MBP	13.1%	5.1%	0.14
	Do not use MBP	39.4%	42.4%	
Abdominoperineal excision	Use MBP	12.5%	5.7%	0.12
	Do not use MBP	40.0%	41.8%	
MBP with oral antibiotics reduces risk of anastomotic leak?				
		Agrees with question	Disagrees with question	Kappa agreement
Right hemicolectomy	Use MBP	5.7%	4.0%	0.15
	Do not use MBP	26.1%	64.2%	
Left-sided resection without stoma	Use MBP	19.0%	23.6%	0.23
	Do not use MBP	12.7%	44.6%	
Left-sided resection with stoma	Use MBP	27.5%	52.3%	0.07
	Do not use MBP	4.2%	16.0%	
Panproctocolectomy	Use MBP	7.7%	10.5%	0.10
	Do not use MBP	24.0%	57.8%	
Abdominoperineal excision	Use MBP	6.9%	11.3%	0.06
	Do not use MBP	24.8%	57.0%	

These are listed according to operation type. Green box, current practice is consistent with the response to the question. Blue box, agree with question but not in practice. Red box, surgeon uses MBP but not for this reason. MBP, mechanical bowel preparation; SSI, surgical site infection.

Discussion

This study demonstrated a considerable variation in the use of MBP and oral antibiotic preparation, both individually and in combination, amongst practising colorectal surgeons. Despite European and UK guidelines advising against routine use of MBP, surgeons continue to use MBP, especially for left-sided resections with a planned stoma. Oral antibiotic preparation was used in between 12% and 20% of cases, with very few surgeons using the combination of MBP/OABP, suggesting that some surgeons use oral antibiotics as an alternative to MBP to reduce infectious complications of colorectal surgery. A wide variation in practice was also demonstrated depending upon the operation being carried out. It is particularly interesting to note the data from this study showing the low rate of both MBP and OABP for patients undergoing panproctocolectomy and abdominoperineal excision resection (APER); deep organ space infection, and perineal wound infections carry a major burden for this population [21]. The findings suggest that decision-making with regard to bowel preparation is driven by reducing the risk of AL from low anterior resection, rather than preventing SSIs, or AL from other resections. Recent American College of Surgeons National Surgical Quality Improvement Program (ACS NSQIP) data supports the use of combined MBP/OAB for all colectomies [14].

Although few surgeons prescribe MBP/OAB, 53% of respondents believe that combined MBP/OABP reduces SSI; 32% believe that the combination reduces AL, indicating a discrepancy between awareness of the benefit of MBP/OABP and day-to-day practice. Based on data from previous surveys, it was

anticipated that surgeons prescribing combined MBP/OABP would be in the minority [22], this study therefore sought to investigate whether ERAS protocols might influence decision-making. It was postulated that surgeons who omit MBP do so on the basis that it is not compatible with ERAS. Forty-one percent of respondents believe that MBP delays return to normal intestinal function; however, only 24% of respondents believe that MBP is incompatible with ERAS. It is possible that despite awareness of the benefit of MBP/OAB, most surgeons have not yet modified their ERAS pathways. Whilst most ERAS pathways do not include MBP, avoidance of MBP was not one of the central tenets described by Kelhlet [23], and it has been shown that MBP/OABP can be incorporated into ERAS pathways [2, 17]. A further barrier to change in practice may be the lack of clarity regarding the precise MBP/OABP preparation. Various combinations of kanamycin, metronidazole, neomycin and erythromycin have been used with variable dosing [13], however, detailed advice to support decision making remains scarce.

This study also explored the relationship between surgeon's views on the effects of MBP and OABP on SSI and AL, and their own practice (Tables II and III). As there were two options for each criterion (agree/disagree; prescribe/do not prescribe), responses fell into one of four groups (e.g., agree/prescribe; agree/do not prescribe). Scrutiny of the outcomes indicated considerable incongruity between individual surgeons' agreement with the statements, and their actual practice. The relationships were quantified with kappa agreements, which showed either a poor or fair agreement for all outcomes. These weak agreements indicate that a large proportion of surgeons agreed with the questions, and believe that combined MBP/OABP reduces SSI and AL, but do not currently prescribe such a regime. Data from this study also indicates that a considerable number of respondents were either unaware of current literature or did not agree with it, e.g., respondents who did not believe that MBP/OABP reduces SSI and AL.

Although many surgeons recognise the benefit of MBP/OAB, very few UK surgeons currently employ the regime as a routine element of pre-operative management. The recent influence of ERAS pathways may have contributed to the omission of MBP/OAB, especially as UK and European ERAS guidelines do not endorse routine use of MBP/OAB. In contradiction to the UK and Europe, North American guidelines do recommend the routine use of MBP/OAB, within the context of an ERAS pathway. Infectious complications of colorectal surgery are likely to counteract many of the advantages gained from ERAS. It is therefore reasonable to suggest that future iterations of UK and European guidelines for elective colorectal surgery should recognize the data that supports the use of MBP/OABP, and empower colorectal surgeons to incorporate combined MBP/OABP into ERAS pathways.

Conflict of interest statement

The authors have no conflicts of interest to declare.

Funding sources

None.

References

- [1] Artinyan A, Orcutt ST, Anaya DA, Richardson P, Chen GJ, Berger DH. Infectious postoperative complications decrease long-

- term survival in patients undergoing curative surgery for colorectal cancer: a study of 12,075 patients. *Ann Surg* 2015;261(3):497–505.
- [2] Holubar SD, Hedrick T, Gupta R, Kellum J, Hamilton M, Gan TJ, et al. American Society for Enhanced Recovery (ASER) and Perioperative Quality Initiative (POQI) joint consensus statement on prevention of postoperative infection within an enhanced recovery pathway for elective colorectal surgery. *Perioper Med (Lond)* 2017;6:4.
- [3] Troughton R, Birgand G, Johnson AP, Naylor N, Gharbi M, Aylin P, et al. Mapping national surveillance of surgical site infections (SSIs) to national needs and priorities: an assessment of England's surveillance landscape. *J Hosp Infect* 2018;100:378–85.
- [4] Abercrombie J. General surgery GIRFT programme national speciality report. 2017.
- [5] Gorgun E, Rencuzogullari A, Ozben V, Stocchi L, Fraser T, Benlice C, et al. An effective bundled approach reduces surgical site infections in a high-outlier colorectal unit. *Dis Colon Rectum* 2018;61(1):89–98.
- [6] Harris J. Success of a colorectal surgical site infection prevention bundle in a multihospital system. *AORN J* 2018;107(5):592–600.
- [7] Keenan JE, Speicher PJ, Nussbaum DP, Adam MA, Miller TE, Mantyh CR, et al. Improving outcomes in colorectal surgery by sequential implementation of multiple standardized care programs. *J Am Coll Surg* 2015;221(2):404–414.e1.
- [8] Klinger AL, Green H, Monlezun DJ, Beck D, Kann B, Vargas HD, et al. The role of bowel preparation in colorectal surgery: results of the 2012–2015 ACS-NSQIP Data. *Ann Surg* 2017. <https://doi.org/10.1097/SLA.0000000000002568>.
- [9] Guenaga KF, Matos D, Wille-Jorgensen P. Mechanical bowel preparation for elective colorectal surgery. *Cochrane Database Syst Rev* 2011;(9):Cd001544.
- [10] Slim K, Kartheuser A. Mechanical bowel preparation before colorectal surgery in enhanced recovery programs: discrepancy between the American and European Guidelines. *Dis Colon Rectum* 2018;61(2):e13–4.
- [11] Cannon JA, Altom LK, Deierhoi RJ, Morris M, Richman JS, Vick CC, et al. Preoperative oral antibiotics reduce surgical site infection following elective colorectal resections. *Dis Colon Rectum* 2012;55(11):1160–6.
- [12] Kiran RP, Murray AC, Chiuzan C, Estrada D, Forde K. Combined preoperative mechanical bowel preparation with oral antibiotics significantly reduces surgical site infection, anastomotic leak, and ileus after colorectal surgery. *Ann Surg* 2015;262(3):416–25. discussion 23–5.
- [13] McSorley ST, Steele CW, McMahon AJ. Meta-analysis of oral antibiotics, in combination with preoperative intravenous antibiotics and mechanical bowel preparation the day before surgery, compared with intravenous antibiotics and mechanical bowel preparation alone to reduce surgical-site infections in elective colorectal surgery. *BJs Open* 2018;2(4):185–94.
- [14] Midura EF, Jung AD, Hanseman DJ, Dhar V, Shah SA, Rafferty JF, et al. Combination oral and mechanical bowel preparations decreases complications in both right and left colectomy. *Surgery* 2018;163(3):528–34.
- [15] Scarborough JE, Mantyh CR, Sun Z, Migaly J. Combined mechanical and oral antibiotic bowel preparation reduces incisional surgical site infection and anastomotic leak rates after elective colorectal resection: an analysis of colectomy-targeted ACS NSQIP. *Ann Surg* 2015;262(2):331–7.
- [16] Vo E, Massarweh NN, Chai CY, Tran Cao HS, Zamani N, Abraham S, et al. Association of the addition of oral antibiotics to mechanical bowel preparation for left colon and rectal cancer resections with reduction of surgical site infections. *JAMA Surg* 2018;153(2):114–21.
- [17] Carmichael JC, Keller DS, Baldini G, Bordeianou L, Weiss E, Lee L, et al. Clinical Practice Guidelines for enhanced recovery after colon and rectal surgery from the American Society of Colon and Rectal Surgeons and Society of American Gastrointestinal and Endoscopic Surgeons. *Dis Colon Rectum* 2017;60(8):761–84.
- [18] The 2017 European Society of Coloproctology (ESCP) Collaborating Group. Association of mechanical bowel preparation with oral antibiotics and anastomotic leak following left sided colorectal resection: An international, multi-centre, prospective audit. *Colorectal Dis* 2018;20(Suppl 6):15–32.
- [19] Garfinkle R, Abou-Khalil J, Morin N, Ghitulescu G, Vasilevsky CA, Gordon P, et al. Is there a role for oral antibiotic preparation alone before colorectal surgery? ACS-NSQIP analysis by coarsened exact matching. *Dis Colon Rectum* 2017;60(7):729–37.
- [20] Altman DG. *Practical statistics for medical research*. London: Chapman & Hall/CRC.; 1991.
- [21] Li W, Stocchi L, Elagili F, Kiran RP, Strong SA. Healing of the perineal wound after proctectomy in Crohn's disease patients: only preoperative perineal sepsis predicts poor outcome. *Tech Coloproctol* 2017;21(9):715–20.
- [22] Devane LA, Proud D, O'Connell PR, Panis Y. A European survey of bowel preparation in colorectal surgery. *Colorectal Dis* 2017;19(11):O402–6.
- [23] Kehlet H. Enhanced Recovery Protocols (ERP): Need for action. *Ann Surg* 2018;268(6):e85.