



Bloodstream infections caused by carbapenem-resistant *Acinetobacter baumannii*: Clinical features, therapy and outcome from a multicenter study



Alessandro Russo^a, Matteo Bassetti^a, Giancarlo Ceccarelli^b, Novella Carannante^c, Angela Raffaella Losito^d, Michele Bartoletti^e, Silvia Corcione^f, Guido Granata^g, Antonella Santoro^h, Daniele Roberto Giacobbe^{i,j}, Maddalena Peghin^a, Antonio Vena^a, Francesco Amadori^k, Francesco Vladimiro Segala^f, Maddalena Giannella^e, Giovanni Di Caprio^c, Francesco Menichetti^k, Valerio Del Bono^l, Cristina Mussini^h, Nicola Petrosillo^g, Francesco Giuseppe De Rosa^f, Pierluigi Viale^e, Mario Tumbarello^d, Carlo Tascini^c, Claudio Viscoli^{i,j}, Mario Venditti^{b,*}, ISGRI-SITA (Italian Study Group on Resistant Infections of the Società Italiana Terapia Antinfettiva)

^a Infectious Diseases Clinic, Department of Medicine, University of Udine and Azienda Sanitaria Universitaria Integrita Presidio Ospedaliero Universitario Santa Maria della Misericordia, Udine, Italy

^b Department of Public Health and Infectious Diseases, Sapienza University of Rome, Policlinico Umberto I, Viale dell'Università 37, 00161 Rome, Italy

^c First division of Infectious Diseases, Cotugno Hospital, AORN dei Colli, Naples, Italy

^d Institute of Infectious Diseases, Università Cattolica del Sacro Cuore, Fondazione Policlinico Universitario Agostino Gemelli, Rome, Italy

^e Department of Medical and Surgical Sciences, University of Bologna, Bologna, Italy

^f Department of Medical Sciences, University of Turin, Infectious Diseases, City of Health and Sciences, Turin, Italy

^g Clinical and Research Department for Infectious Diseases, Unit Systemic and Immunodepression-Associated Infections, National Institute for Infectious Diseases L. Spallanzani, Rome, Italy

^h Clinic of Infectious Disease, University Hospital, Modena, Italy

ⁱ Infectious Diseases Unit, Ospedale Policlinico San Martino – IRCCS per l'Oncologia, Italy

^j Department of Health Sciences, University of Genoa, Genoa, Italy

^k Infectious Diseases Clinic, Nuovo Santa Chiara University Hospital, Azienda Ospedaliera Universitaria Pisana, Pisa, Italy

^l Infectious Diseases Unit, Azienda Ospedaliera S. Croce e Carle, Cuneo, Italy

ARTICLE INFO

Article history:

Accepted 24 May 2019

Available online 28 May 2019

Keywords:

Bacteremia

Septic shock

Acinetobacter

Multidrug-resistant

Colistin

SUMMARY

Objectives: bloodstream infections (BSI) due to multidrug-resistant (MDR) *Acinetobacter baumannii* (AB) have been increasingly observed among hospitalized patients.

Methods: prospective, observational study conducted among 12 large tertiary-care hospitals, across 7 Italian regions. From June 2017 to June 2018 all consecutive hospitalized patients with bacteremia due to MDR-AB were included and analyzed in the study.

Results: During the study period 281 episodes of BSI due to MDR-AB were observed: 98 (34.8%) episodes were classified as primary bacteremias, and 183 (65.2%) as secondary bacteremias; 177 (62.9%) of them were associated with septic shock. Overall, 14-day mortality was observed in 172 (61.2%) patients, while 30-day mortality in 207 (73.6%) patients. On multivariate analysis, previous surgery, continuous renal replacement therapy, inadequate source control of infection, and pneumonia were independently associated with higher risk of septic shock. Instead, septic shock and Charlson Comorbidity Index >3 were associated with 14-day mortality, while adequate source control of infection and combination therapy with survival. Finally, septic shock, previous surgery, and aminoglycoside-containing regimen were associated with 30-day mortality, while colistin-containing regimen with survival.

Conclusions: BSI caused by MDR-AB represents a difficult challenge for physicians, considering the high rates of septic shock and mortality associated with this infection.

© 2019 The British Infection Association. Published by Elsevier Ltd. All rights reserved.

* Corresponding author.

E-mail address: mario.venditti@uniroma1.it (M. Venditti).

Introduction

In recent years, bloodstream infections (BSI) due to multidrug-resistant (MDR) gram-negative bacteria such as *Acinetobacter baumannii* (AB) have been increasingly observed among hospitalized patients admitted to the intensive care unit (ICU), surgical and medical wards.^{1–3} MDR-AB has been defined one of the top priority pathogens by the World Health Organization^{4,5}; specifically, in Italy an increased incidence of MDR-AB was observed in the last years.⁶

Acinetobacter baumannii bacteria are usually resistant to carbapenems and to β -lactams, aminoglycosides, rifampin, and fluoroquinolones.^{7,8} There are limited therapeutic options resulting often in inappropriate therapy and subsequent negative impact on outcome. New agents with microbiological activity against MDR-AB strains have been recently developed and they will be available in the near future.^{9–11} Mortality rate over 60% has been reported for MDR-AB infections,¹² particularly in patients with septic shock.¹³

Early diagnosis and adequate administration of antimicrobials are the milestone for the management of critically ill patients with MDR-AB,^{14–16} and recent data were reported in literature comparing monotherapy with combination therapy.¹⁷ On these basis, physicians should recognize peculiar clinical characteristics and treat MDR-AB infections appropriately in hospitalized patients,¹⁸ so the aim of the present study was to analyse clinical features, antimicrobial treatment and outcome of patients with BSI due to MDR-AB.

Patients and method

Study design and patient selection

Prospective, observational study conducted among 12 large tertiary-care hospitals across 7 Italian regions. From June 2017 to June 2018 all consecutive hospitalized patients with bacteremia caused by MDR-AB were included in the study. Inclusion criteria included: (1) age \geq 18 years; (2) blood culture positive for MDR-AB; (3) clinical signs consistent with infection. Polymicrobial etiology was excluded; only one episode of MDR-AB infection for each patient was reported in the study period. The study was conducted according to the principles stated in the Declaration of Helsinki and has been approved by local Ethical Committees (No 4547-2017); informed consent was waived by Ethical Committees. The present study was promoted by ISGRI-SITA (Italian Study Group on Resistant Infections of the Società Italiana Terapia Antinfettiva).

Patient data were collected from medical charts and from hospital computerized databases or clinical charts according to a pre-established questionnaire. The following information were reviewed: demographics; clinical and laboratory findings; comorbid conditions; microbiological data; duration of ICU and hospital stay; any MDR infection during hospitalization; treatment and procedures (e.g. non-invasive ventilation, mechanical ventilation, continuous renal replacement therapy [CRRT]) carried out during hospitalization and/or in the 30 days prior to infection; class of antibiotics received on admission and/or during admission before a positive culture of a biological sample was obtained; the simplified acute physiology score (SAPS II); sequential organ failure assessment (SOFA) and quick (q)-SOFA at time of infection; anamnestic MDR-AB colonization or during hospitalization; source of infection and its adequate control; antibiotic regimens used for MDR-AB infection; development of septic shock; all-cause 14- and 30-day mortality.

Definitions and MDR-AB identification

Infections were defined according to the standard definitions of the European Centers for Disease Control and Prevention (eCDC).¹⁹

Infection was defined as the presence of at least 1 positive blood culture for MDR-AB in individuals with signs and symptoms consistent with infection;²⁰ concomitant isolation of MDR-AB in other sites like urine, skin swabs or biopsies, lung or abdomen were also recorded. Infection onset was defined as the date of collection of the first positive blood culture for MDR-AB. Infection was considered of unknown origin when the source of infection was not established after routine clinical, microbiological, and radiological examination. Primary BSI was defined as BSI occurring in patients without a recognized source of infection. The central venous catheter (CVC) was considered the source of infection if one of the following condition were present: (1) a positive result of semi-quantitative (>15 colony-forming units per catheter segment) catheter culture, whereby the same species was isolated from the catheter segment and a peripheral blood culture; (2) growth in a blood culture obtained through a catheter hub is detected by an automated system at least 2 h earlier than a peripheral blood culture of equal volume simultaneously collected, providing that the same species was isolated.²¹

Septic shock was defined according to international definitions.²² The severity of clinical conditions was determined by using SAPS II, SOFA and qSOFA scores calculated at the time of infection onset. Length of hospital and ICU stay were calculated as the number of days from the date of admission to the date of discharge or death.

Identification of MDR-AB strains was based accordingly with local laboratory techniques. The Vitek 2 automated system (bioMérieux, Marcy l'Etoile, France) was used for isolate identification and antimicrobial susceptibility testing. Minimum inhibitory concentrations (MICs) were established according to the European Committee on Antimicrobial Susceptibility Testing (EUCAST) breakpoints.²³ Isolated strains were classified as multidrug-resistant (MDR), extensively drug-resistant (XDR) and pandrug-resistant (PDR) according to Magiorakos et al.²⁴




Fig. 1. Wards of hospitalization at time of infection.

Table 1
Univariate analysis comparing survivors and non-survivors at 30 days from infection onset.

Variables	Survivors n = 74 (%)	Non-survivors n = 207 (%)	HR (CI 95%)	P
Age, mean ± SD (years)	53.9 ± 17.4	63.8 ± 15.9	2.2 (1.9–4.1)	<0.001
Male sex	60 (81.1)	134 (64.7)	0.42 (0.2–0.7)	0.009
Comorbidities				
Chronic liver disease	6 (8.1)	16 (7.7)	0.94 (0.8–1.3)	1.0
Neoplasm	11 (14.9)	33 (15.9)	1.08 (0.8–1.2)	1.0
Diabetes	19 (25.7)	57 (27.5)	1.1 (0.8–1.4)	0.87
Heart failure	6 (8.1)	49 (23.7)	3.51 (2.4–5.3)	0.003
Coronary artery disease	26 (35.1)	103 (49.8)	1.82 (1.4–3.4)	0.04
Chronic renal disease	10 (13.5)	41 (19.8)	1.58 (0.7–2.1)	0.29
COPD	23 (31.1)	59 (28.5)	0.88 (0.6–1.2)	0.76
Neurological disease	7 (9.4)	10 (4.8)	1.01 (0.8–1.2)	1.0
SOT/HSCT	2 (2.7)	9 (4.3)	2.98 (0.9–3.5)	0.2
>2 comorbidities	27 (36.4)	48 (23.1)	0.91 (0.5–1.3)	0.59
Charlson comorbidity index, mean ± SD	3.8 ± 3.5	6.4 ± 3.9	3.1 (2.4–4.8)	0.004
Previous hospitalization (90 days)	26 (35.1)	69 (33.3)	0.92 (0.6–1.5)	0.77
Previous ICU admission (90 days)	7 (9.5)	29 (14)	1.55 (0.7–1.9)	0.41
Previous surgery (30 days)	11 (14.8)	64 (30.9)	2.77 (2.1–5.2)	0.004
Previous antibiotic therapy (30 days)	38 (51.4)	115 (55.6)	1.18 (0.8–1.7)	0.58
Previous <i>Acinetobacter</i> spp colonization/infection	4 (5.4)	8 (3.8)	0.85 (0.6–1.2)	1.0
<i>Acinetobacter</i> colonization prior infection	11 (14.9)	20 (9.7)	0.61 (0.5–1.3)	0.27
Source of infection				
Primary bacteremia	32 (43.2)	66 (31.8)	0.58 (0.3–1.1)	0.06
CVC-related bacteremia	1 (1.4)	10 (4.8)	3.37 (0.8–3.9)	0.29
Pneumonia	38 (51.4)	114 (55.1)	1.16 (0.8–1.4)	0.59
Catheter-related urinary tract	0	6 (2.9)	1.23 (0.8–1.3)	0.34
SSTI	2 (2.7)	0	0.25 (0.1–1.1)	0.06
Intra-abdominal	1 (1.4)	11 (5.3)	4.09 (0.8–4.3)	0.19
Fever	24 (32.4)	31 (14.9)	0.8 (0.5–1.8)	0.67
NIV	2 (2.7)	7 (3.3)	2.25 (0.8–2.8)	0.48
Mechanical ventilation	33 (44.5)	61 (29.4)	0.65 (0.4–1.7)	0.24
CRRT	8 (10.8)	22 (10.6)	1.01 (0.8–1.1)	0.2
ECMO	2 (2.7)	2 (0.9)	0.6 (0.4–1.8)	0.63
Inadequate source control of infection	22 (29.7)	82 (39.7)	1.51 (0.8–1.9)	0.16
Previous MDR infections during hospital stay	30 (40.5)	62 (30)	0.62 (0.4–1.3)	0.11
PCT at time of infection onset, mean ± SD	18.2 ± 49.8	33.9 ± 68.1	1.7 (0.8–1.9)	0.26
CRP at time of infection onset, mean ± SD	29.9 ± 51.3	51.1 ± 83.9	1.3 (0.7–2)	0.25
Steroid therapy	28 (37.8)	88 (42.5)	1.21 (0.8–1.8)	0.49
Transfer in ICU	7 (9.5)	35 (16.9)	7.5 (2.2–9.3)	0.001
Septic shock	31 (41.9)	146 (70.5)	3.32 (2.5–8.9)	<0.001
Length of hospitalization, mean ± SD (days)	54.2 ± 35.6	32.5 ± 27.3	0.6 (0.2–0.8)	<0.001
Length of ICU stay, mean ± SD (days)	42.9 ± 30.4	24.1 ± 22.7	0.5 (0.3–0.7)	<0.001
Length of definitive antibiotic therapy, mean ± SD (days)	15.1 ± 8.2	8.4 ± 8.3	0.63 (0.4–0.9)	<0.001
SAPS II at time of infection onset, mean ± SD	35.7 ± 14.1	46.6 ± 15.3	2.1 (1.8–4.2)	<0.001
qSOFA at time of infection onset, mean ± SD	1.5 ± 0.9	2.1 ± 1	1.2 (1.1–2.3)	0.003
SOFA at time of infection onset, mean ± SD	6.2 ± 3	7.2 ± 3.1	1.1 (0.8–1.4)	0.11

OR: odds ratio; SD: standard deviation; COPD: chronic obstructive pulmonary disease; SOT/HSCT: solid organ transplant/hematopoietic stem cell transplant; ICU: intensive care unit; CVC: central venous catheter; SSTI: skin and soft-tissue infection; NIV: non-invasive ventilation; CRRT: continuous renal replacement therapy; ECMO: extracorporeal membrane oxygenation; MDR: multidrug-resistant; PCT: procalcitonin; CRP: c-reactive protein; SAPS: simplified acute physiology score; SOFA: sequential organ failure assessment.

Table 2
Cox regression analysis about risk factors associated with 14-day and 30-day mortality.

14-day mortality				30-day mortality			
Variables	HR	CI 95%	p	Variables	HR	CI 95%	p
Septic shock	10.79	1.12–141	0.04	Septic shock	1.54	1.04–2.27	0.03
Adequate source control of infection	0.22	0.01–0.42	0.01	Colistin-containing regimen	0.41	0.27–0.63	<0.001
Charlson comorbidity index > 3	1.44	1.04–2	0.02	Aminoglycoside-containing regimen	2.57	1.33–4.94	0.005
Combination therapy	0.36	0.01–0.89	0.03	Previous surgery (30 days)	1.63	1.16–2.29	0.005

HR: hazard ratio; CI: confidence interval.

Antimicrobial treatment evaluation and adequate source control of infection

Depending on the number of drugs used (1 or >1), treatment regimens were classified either as monotherapy or combination therapy. Initial antibiotic therapy, defined as antimicrobial chemotherapy implemented within 24 h after the onset of infection, was assessed along with definitive antibiotic therapy, defined as antimicrobial treatment based on *in vitro* MDR-AB isolate sus-

ceptibilities. Drugs in definitive therapy must have been administered for at least 50% of the total duration of therapy (except for patients who died while on definitive therapy, who were included if they received at least 1 complete day of therapy). Time to initial definitive therapy was the period between the infection onset and initial definitive therapy.

Adequate control of source of infection was defined as the removal of any preexisting contaminated CVC as well as the drainage of intra-abdominal abscesses or other fluid collections have been

Table 3
Univariate analysis comparing patients developing or not septic shock.

Variables	No septic shock n = 104 (%)	Septic shock n = 177 (%)	OR	P
Age, mean ± SD (years)	58.6 ± 18.1	62.7 ± 15.8	2.07 (0.8–2.4)	0.46
Male sex	74 (71.2)	120 (67.8)	0.7 (0.5–1.4)	0.59
Comorbidities				
Chronic liver disease	12 (11.5)	10 (5.6)	0.45 (0.3–1.9)	0.1
Neoplasm	20 (19.2)	24 (13.6)	0.65 (0.3–1.4)	0.23
Diabetes	27 (26)	49 (27.7)	1.09 (0.6–1.5)	0.78
Heart failure	15 (14.4)	40 (22.6)	1.72 (0.8–2.2)	0.11
Coronary artery disease	33 (31.7)	96 (54.2)	2.55 (1.8–4.5)	<0.001
Chronic renal disease	16 (15.4)	35 (19.8)	1.35 (0.7–1.7)	0.42
COPD	32 (30.8)	50 (28.2)	0.88 (0.5–1.3)	0.68
Neurological disease	11 (10.6)	6 (3.3)	0.8 (0.5–1.5)	0.78
SOT/HSCT	5 (4.8)	6 (3.3)	0.64 (0.4–1.5)	0.34
>2 comorbidities	44 (42.3)	31 (17.5)	0.8 (0.5–1.4)	0.72
Charlson Comorbidity Index, mean SD	5.1 ± 4	5.3 ± 3.9	1.26 (0.8–1.4)	0.85
Previous hospitalization (90 days)	40 (38.5)	55 (31.1)	0.72 (0.5–1.3)	0.24
Previous ICU admission (90 days)	12 (11.5)	24 (13.6)	1.2 (0.6–1.9)	0.71
Previous surgery (30 days)	16 (15.4)	59 (33.3)	2.76 (1.5–4.6)	0.001
Previous antibiotic therapy (30 days)	59 (56.7)	94 (53.1)	0.86 (0.3–1.3)	0.62
Previous <i>Acinetobacter</i> spp colonization/infection	4 (3.8)	8 (4.5)	3.42 (0.9–3.9)	0.06
<i>Acinetobacter</i> colonization prior infection	11 (10.6)	20 (11.3)	1.07 (0.9–1.2)	1.0
Source of infection				
Primary bacteremia	52 (50)	46 (25.9)	0.33 (0.1–0.6)	<0.001
CVC-related bacteremia	4 (3.8)	7 (4)	1.02 (0.9–1.1)	1.0
Pneumonia	44 (42.3)	108 (61)	2.13 (1.8–2.9)	0.003
Catheter-related urinary tract	4 (3.8)	2 (1.1)	0.28 (0.2–1.3)	0.19
SSTI	0	2 (1.1)	1.2 (0.8–1.4)	0.53
Intra-abdominal	0	12 (6.8)	2.6 (2.1–4.4)	0.005
Fever	37 (35.5)	18 (10.1)	0.51 (0.4–1.2)	0.13
NIV	7 (8.6)	2 (3.8)	0.41 (0.2–1.3)	0.48
Mechanical ventilation	48 (59.3)	46 (88.5)	5.27 (3.1–7.2)	<0.001
CRRT	1 (0.9)	29 (16.4)	3.66 (2.8–5.8)	0.003
ECMO	1 (0.9)	3 (1.6)	4.8 (0.8–4.8)	0.3
Inadequate source control of infection	20 (19.2)	84 (47.5)	3.79 (3.1–6.4)	<0.001
Previous MDR infections during hospital stay	43 (41.3)	49 (27.7)	0.54 (0.2–0.7)	0.02
PCT at time of infection onset, mean ± SD	19.1 ± 45.2	37 ± 75.8	1.7 (0.8–2.2)	0.198
CRP at time of infection onset, mean ± SD	38.6 ± 55.9	51.2 ± 95.8	1.4 (0.8–1.7)	0.45
Steroid therapy	31 (29.8)	85 (48)	2.17 (2–5.1)	0.004
Transfer in ICU	23 (22.1)	19 (10.7)	0.33 (0.1–0.6)	0.009
Length of hospitalization, mean ± SD (days)	47.6 ± 37.8	32.7 ± 25	0.87 (0.4–0.9)	<0.001
Length of ICU stay, mean ± SD (days)	31.3 ± 33.2	26.2 ± 21.6	0.5 (0.3–1.3)	0.13
Length of definitive antibiotic therapy, mean ± SD (days)	11 ± 7.7	9.7 ± 9.3	0.98 (0.5–1.3)	0.2
SAPS II at time of infection onset, mean ± SD	37.4 ± 16.4	47.4 ± 14	1.9 (1.2–2.6)	<0.001
qSOFA at time of infection onset, mean ± SD	1.5 ± 0.9	2.45 ± 0.9	2.4 (1.9–3.2)	<0.001
SOFA at time of infection onset, mean ± SD	6.1 ± 2.6	8.1 ± 3.5	2.3 (1.8–3.8)	0.004
30-day mortality	61 (58.7)	146 (82.5)	3.32 (2.4–5.3)	<0.001

OR: odds ratio; SD: standard deviation; COPD: chronic obstructive pulmonary disease; SOT/HSCT: solid organ transplant/hematopoietic stem cell transplant; ICU: intensive care unit; CVC: central venous catheter; SSTI: skin and soft-tissue infection; NIV: non-invasive ventilation; CRRT: continuous renal replacement therapy; ECMO: extracorporeal membrane oxygenation; MDR: multidrug-resistant; PCT: procalcitonin; CRP: c-reactive protein; SAPS: simplified acute physiology score; SOFA: sequential organ failure assessment.

performed within 24 h after the onset of infection. Timing of CVC removal was based on medical record review and was confirmed by review of patient radiographs.

Primary endpoint and statistical analysis

The primary endpoint of the study was to evaluate risk factors associated with survival or death at 30 days from BSI onset. Secondary endpoints were risk factors associated with 14-day mortality and septic shock.

To detect significant differences between groups, we used Chi-square test or Fisher exact test for categorical variables, and the 2-tailed *t*-test or Mann-Whitney test for continuous variables, when appropriate. All pre-treatment variables identified at univariate analysis were tested at logistic regression analysis to identify risk factors associated with development of septic shock. In a multivariate analysis of survival, the Cox regression model was tested using a proportional hazards model analysis with backward stepwise selection and $p < 0.05$ for all variables, to determine the effects of all anamnestic, clinical and therapeutic variables

on 14- and 30-day survival. Kaplan–Meier curves were used to determine survival at 14- and 30-day in patients treated either with monotherapy or combination therapy. Survival curves for time-to-event variables, constructed with the use of Kaplan–Meier estimates, were based on all available data and were compared with the use of the log-rank test. Wald confidence intervals and tests for hazard- and odds-ratios were computed based on the estimated standard errors. Possible confounding factors and interactions were weighted during analyses. Statistical significance was established at ≤ 0.05 . All reported *P* values are 2-tailed. The results obtained were analyzed using a commercially available statistical software packages (SPSS, version 20.0; SPSS Inc, Chicago, Illinois).

Results

During the study period, 281 BSI caused by MDR-AB strains were observed. Out of these, 98 episodes (34.8%) were classified as primary bacteremia, and 183 (65.2%) cases as secondary bacteremia; 177 (62.9%) episodes were associated with development of septic shock. Resistance rates were the following: colistin 1.4%,

Table 4

Logistic regression analysis about risk factors associated with development of septic shock.

Variables	OR	CI 95%	p
Previous surgery (30 days)	33.9	4.86–236.82	<0.001
CRRT	15.4	3.77–63.09	<0.001
Inadequate source control of infection	4.21	1.06–16.7	0.04
Pneumonia	5.29	1.35–20.59	0.016

OR: odds ratio; CI: confidence interval; CRRT: continuous renal replacement therapy; SOFA: sequential organ failure assessment.

gentamicin 87.3%, amikacin 89.1%, and meropenem 100%. On these basis, 98.6% of AB strains were considered XDR and 1.4% PDR. Finally, 14-day mortality was observed in 172 (61.2%) patients, while 30-day mortality in 207 (73.6%) patients.

Fig. 1 shows hospital wards at time of infection onset: 83% of patients were hospitalized in ICU, 9.7% in medical wards, 3.6% in surgical wards, and 1.7% in emergency department.

Table 1 shows univariate analysis comparing survivors and non-survivors at 30 days from infection onset. Differences between survivors and non-survivors were reported for age (53.9 ± 17.4 Vs 63.8 ± 15.9 years, $p < 0.001$), male sex (81.1% Vs 64.7, $p = 0.009$), heart failure (8.1% Vs 23.7%, $p = 0.003$), Charlson Comorbidity Index (3.8 ± 3.5 Vs 6.4 ± 3.9 points, $p = 0.004$), patients transferred to ICU (9.5% Vs 16.9%, $p = 0.001$), septic shock (41.9% Vs 70.5%, $p < 0.001$), SAPS II at time of infection onset (35.7 ± 14.1 Vs 46.6 ± 15.3 points, $p < 0.001$), and qSOFA at time of infection onset (1.5 ± 0.9 Vs 2.1 ± 1 points, $p = 0.003$).

As reported in Table 2, Cox regression analysis of factors associated with 14-day mortality in MDR-AB BSI showed that septic shock (HR 10.79, CI95% 1.12–141, $p = 0.04$), and Charlson Comorbidity Index >3 (HR 1.44, CI95% 1.04–2, $p = 0.02$) were associated with death, while adequate source control of infection (HR 0.22, CI95% 0.01–0.42, $p = 0.01$) and combination therapy (HR 0.36, CI95% 0.01–0.89, $p = 0.03$) were associated with survival. Conversely, Cox regression analysis of factors associated with 30-day mortality showed that septic shock (HR 1.54, CI95% 1.04–2.27, $p = 0.03$), previous surgery (HR 1.63, CI95% 1.16–2.29, $p = 0.005$), and aminoglycoside-containing regimen (HR 2.57, CI95% 1.33–4.94, $p = 0.005$) were associated with death, while colistin-containing regimen (HR 0.41, CI95% 0.27–0.63, $p < 0.001$) is associated with 30-day survival.

Univariate analysis comparing patients with septic shock or not secondary to MDR-AB infection is reported in Table 3. In patients developing septic shock were recorded more frequently previous surgery (33.3% Vs 15.4%, $p = 0.001$), pneumonia (61% Vs 42.3%, $p = 0.003$), intra-abdominal source of infection (6.8% Vs 0, $p = 0.005$), mechanical ventilation (88.5% Vs 59.3%, $p < 0.001$), CRRT (16.4% Vs 0.9%, $p = 0.003$), adequate source control of infection (47.5% Vs 19.2%, $p < 0.001$), steroid therapy (48% Vs 29.8%, $p = 0.004$), if compared with patients without septic shock. Primary bacteremia was observed more frequently in patients without septic shock (50% Vs 25.9%, $p < 0.001$). Finally, 30-day mortality was 82.5% in patients with septic shock ($p < 0.001$).

A logistic regression analysis about risk factors associated with development of septic shock is reported in Table 4: previous surgery (OR 33.9, CI95% 4.86–236.82, $p < 0.001$), CRRT (OR 15.4, CI95% 3.77–63.09, $p < 0.001$), no source control of infection (OR 4.21, CI95% 1.06–16.7, $p = 0.04$), and pneumonia (OR 5.29 CI95% 1.35–20.59, $p = 0.016$) were independently associated with higher risk of septic shock associated with BSI due to MDR-AB strain.

Characteristics of antibiotic regimens in definitive therapy used in overall population are reported in Table 5. Monotherapy was the choice for 46 (16.3%) patients; a combination of two antibiotics was used in 133 (47.3%) patients; the use of a colistin-

Table 5

Antibiotic regimens as definitive therapy among overall population.

Antibiotic therapy ^a	Overall population n = 281 (%)
Use of only 1 antibiotic as definitive therapy	46 (16.3)
Use of 2 antibiotics in combination as definitive therapy	133 (47.3)
Use of 3 antibiotics in combination as definitive therapy	74 (26.3)
Use of 4 antibiotics in combination as definitive therapy	16 (5.6)
Use of 5 antibiotics in combination as definitive therapy	1 (0.4)
Colistin-containing regimen as definitive therapy	231 (82.2)
Tigecycline-containing regimen as definitive therapy	69 (24.5)
Aminoglycoside-containing regimen as definitive therapy	16 (5.7)
Rifampin-containing regimen as definitive therapy	71 (25.2)
Ampicillin/sulbactam-containing regimen as definitive therapy	2 (0.7)
Fosfomycin-containing regimen as definitive therapy	4 (1.4)
Trimethoprim/sulfamethoxazole-containing regimen as definitive therapy	5 (1.7)
Vancomycin-containing regimen as definitive therapy	14 (4.9)
Carbapenem-containing regimen as definitive therapy	194 (69)
Use of colistin aerosol inhalation therapy	28 (9.9)
Time to initial definitive therapy, mean \pm SD (days)	4.4 \pm 1.2

SD: standard deviation.

^a During the study period, the usual antimicrobial dosages, adopted for the most used antibiotics, were the following: for colistin, a loading dose of 9 million IU followed by 4.5 million IU every 12 h; for tigecycline, a loading dose of 150–200 mg followed by 100 mg every 12 h; for gentamicin, a dosage of 5 mg/kg every 24 h; for rifampin, a dosage of 10 mg/kg/day; for meropenem, a dosage of 2 g every 8 h or 1.5 g every 6 h; for fosfomycin 12–24 g/day divided every 6–8 h; for ampicillin/sulbactam 3 g every 6 h; for trimethoprim/sulfamethoxazole 15–20 mg/kg/day divided every 6 h; for vancomycin 40 mg/kg/day divided every 12 h.

containing regimen was used for 231 (82.2%) patients, and a carbapenem-containing regimen in 194 (69%) patients. Colistin was used as a monotherapy regimen in 40 cases; in combination therapy mainly with a carbapenem (73.5% of cases), and tigecycline (34.5% of cases). No differences were reported in 30-day mortality about monotherapy or combination therapy with colistin (15 [37.5%] Vs 131 [68.5%] patients, $p = 0.06$). The colistin aerosol inhalation therapy was used in 28 (9.9%) patients; the time to initial definitive therapy was 4.4 ± 1.2 days. Comparison between characteristics of patients treated with monotherapy and combination therapy is reported in Table 6. Older age (67.7 ± 15.2 Vs 59.9 ± 16.8 years, $p = 0.004$), and >2 comorbidities (54.3% Vs 21.2%, $p = 0.001$) were reported more frequently in patients treated with monotherapy; conversely, adequate source control of infection (42.1% Vs 10.9%, $p < 0.001$), transfer in ICU (33% Vs 19.5%, $p = 0.004$), and higher SAPS II at time of infection (44.7 ± 15.1 Vs 39 ± 17.8 points, $p = 0.02$) were more frequent in patients treated with combination therapy. Finally, statistically significant differences were reported for 14-day mortality ($p < 0.001$), but not for 30-day mortality. Rates of 30-day mortality in patients treated with monotherapy (71.7%), combination therapy (74.04%), two-drug combination (70.6%), three-drug combination (78.3%), four drug combination (68.7%), colistin-containing (70.9%), and carbapenem-containing regimens (73.1%) are reported in Fig. 2.

Finally, Kaplan–Meier analysis of 14- and 30-day survival of patients treated with monotherapy or combination therapy is reported in Fig. 3.

Discussion

In the present study we observed very high rates of 14-day (61.2%) and 30-day (73.6%) mortality; most of patients developed septic shock (62.9%) with a very high mortality (82.5%). The high mortality related to *Acinetobacter baumannii* infections was firstly reported in patients with hematologic malignancies;²⁵ afterwards, Freire et al. evaluated mortality in cancer patients with BSI caused

Table 6
Comparison between patients treated with monotherapy or combination therapy.

Variables	Monotherapy n = 46 (%)	Combination therapy n = 235 (%)	P
Age, mean \pm SD (years)	67.7 \pm 15.2	59.9 \pm 16.8	0.004
Male sex	30 (65.2)	164 (69.8)	0.6
Comorbidities			
Chronic liver disease	8 (17.4)	14 (6)	0.01
Neoplasm	10 (21.7)	34 (14.5)	0.26
Diabetes	20 (43.5)	56 (23.8)	0.01
Heart failure	13 (28.3)	42 (17.9)	0.1
Coronary artery disease	26 (56.5)	103 (43.8)	0.14
Chronic renal disease	10 (21.7)	41 (17.4)	0.53
COPD	22 (47.8)	60 (25.5)	0.004
Neurological disease	5 (10.9)	12 (5.1)	1.0
SOT/HSCCT	1 (2.1)	10 (4.2)	0.45
>2 comorbidities	25 (54.3)	50 (21.2)	0.001
Charlson Comorbidity Index, mean SD	5.5 \pm 3.1	5 \pm 4.3	0.64
Previous hospitalization (90 days)	22 (47.8)	73 (31.1)	0.04
Previous ICU admission (90 days)	5 (10.9)	31 (13.2)	0.81
Previous surgery (30 days)	10 (21.7)	65 (27.6)	0.7
Previous antibiotic therapy (30 days)	28 (60.9)	125 (53.2)	0.41
Previous <i>Acinetobacter</i> spp colonization/infection	1 (3.3)	10 (9.6)	0.29
<i>Acinetobacter</i> colonization prior infection	5 (10.9)	26 (11.1)	1.0
Source of infection			
Primary bacteremia	17 (37)	81 (34.4)	0.73
CVC-related bacteremia	0	11 (4.7)	0.22
Pneumonia	23 (50)	129 (54.9)	0.62
Catheter-related urinary tract	2 (4.3)	4 (1.7)	0.25
SSTI	0	2 (2.9)	1.0
Intra-abdominal	4 (8.7)	8 (3.4)	0.11
Fever	21 (45.6)	34 (14.5)	0.17
NIV	2 (4.3)	7 (2.9)	1.0
Mechanical ventilation	15 (32.6)	79 (33.6)	0.07
CRRT	7 (15.2)	23 (9.7)	1.0
ECMO	1 (2.1)	3 (1.2)	1.0
Source control of infection	5 (10.9)	99 (42.1)	<0.001
Previous MDR infections during hospital stay	21 (45.7)	71 (30.2)	0.05
PCT at time of infection onset, mean \pm SD	34.2 \pm 96.7	25.4 \pm 51.6	0.63
CRP at time of infection onset, mean \pm SD	69.4 \pm 114.2	36.1 \pm 57.1	0.09
Steroid therapy	18 (39.1)	98 (41.7)	0.87
Transfer in ICU	9 (19.5)	33 (14)	0.004
Septic shock	26 (56.5)	151 (64.3)	0.32
Length of hospitalization, mean \pm SD (days)	32.6 \pm 30.3	39.3 \pm 31.2	0.18
Length of ICU stay, mean \pm SD (days)	13.8 \pm 11	30.1 \pm 27.2	<0.001
Length of definitive antibiotic therapy, mean \pm SD (days)	8.3 \pm 2	10.5 \pm 9.3	0.14
SAPS II at time of infection onset, mean \pm SD	39 \pm 17.8	44.7 \pm 15.1	0.02
qSOFA at time of infection onset, mean \pm SD	1.9 \pm 1	1.8 \pm 1	0.75
SOFA at time of infection onset, mean \pm SD	6.5 \pm 2.6	7 \pm 3.2	0.51
14-day mortality	32 (69.5)	140 (59.5)	<0.001
30-day mortality	33 (71.7)	174 (74)	0.71

SD: standard deviation; COPD: chronic obstructive pulmonary disease; SOT/HSCCT: solid organ transplant/hematopoietic stem cell transplant; ICU: intensive care unit; CVC: central venous catheter; SSTI: skin and soft-tissue infection; NIV: non-invasive ventilation; CRRT: continuous renal replacement therapy; ECMO: extracorporeal membrane oxygenation; MDR: multidrug-resistant; PCT: procalcitonin; CRP: c-reactive protein; SAPS: simplified acute physiology score; SOFA: sequential organ failure assessment.

by MDR-AB, with 7-day and 30-day mortality rates of 71.7% and 83.7%, respectively.²⁶ Recent studies confirmed these data with mortality over 90% in patients with septic shock.^{12,13} Moreover, our data show that MDR-AB BSI remains a peculiar ICU-acquired infection, although is now more frequently observed even in medical and surgical wards.

Risk factors affecting 14- and 30-day outcome have been explored in our population. As previously reported in literature, these data confirm the weight of comorbidities and septic shock in determining higher 30-day mortality.^{3,13,16} Interestingly, previous surgery was associated with 30-day mortality while adequate source control with 14-day survival. These observations emphasize the role of early source control, that remains particularly complicated in abdominal surgical patient.

Different antibiotic combinations have been studied for treatment of severe infections sustained by MDR-AB;^{27,28} in this context, the role of aerosolized colistin as adjunctive treatment for treatment of severe pneumonia was also investigated.^{29–31} In a randomized clinical trial,³² in patients with MDR-AB infections

mortality was not reduced by addition of rifampicin to colistin; further *in vitro* studies explored the synergism of some drug combinations, especially colistin plus carbapenem for treatment of MDR-AB infections,^{33,34} suggesting the advantage of this combination based on high *in vitro* synergy rates. Combination of a carbapenem plus colistin seems to be the first option for treatment of MDR-AB infections,³⁵ even in our population. Recently, Paul and coworkers reported data from a randomised controlled trial comparing colistin alone versus colistin plus meropenem for treatment of severe infections caused by carbapenem-resistant Gram-negative bacteria.¹⁷ Authors concluded that combination therapy was not more efficient than monotherapy, and that adding meropenem to colistin did not improve clinical failure in severe MDR-AB infections. Of importance, Dickstein and coworkers performed a subgroup analysis on patients with *Acinetobacter* infections and reported that colistin monotherapy was associated with a better outcome compared to colistin-meropenem combination therapy.³⁶ It is important to underline that studies comparing efficacy of monotherapy (mainly colistin) with combination regimens for *Acinetobacter baumannii*





Fig. 2. 30-day mortality and antibiotic regimens in definitive therapy among overall population.



* $p=0.003$

** $p=0.82$

Fig. 3. Kaplan–Meier curves about 14-day (A) and 30-day (B) survival of patients treated with monotherapy (green line*) or combination therapy (blue line**).

infections included a spectrum of different severe infections, like ventilator associated pneumonia, but not always associated with bacteremia. On this basis, our study confirms that comparative studies on MDR-AB therapy should include bacteremic patients. In our population, combination therapy was associated with higher survival at 14 days, but no differences were reported at 30 days if compared to monotherapy (see Fig. 3); however, all patients (except in 1 case) treated with monotherapy died in the first 14 days from the diagnosis of BSI (see Table 4). In our interpretation, sepsis and septic shock are associated with a lethal cascade of events that is unlikely to be interrupted even by an appropriate initial

antimicrobial treatment. In addition, most of our patients were severely ill and would probably have been unable to survive their infections independently of the administration of an adequate initial antimicrobial treatment.

Despite the relevant association between colistin use and 30-day survival, recent EUCAST recommendations,³⁷ based on recent observations³⁸ advertised about potential false susceptibility to colistin in approximately 50% of *Acinetobacter baumannii* strains with automated systems or Etest. Therefore, the very high rates of mortality observed in our population and in published studies^{13,26} might be also attributed to a reported false susceptibility to colistin

in patients for whom physicians were confident in the prescribing a colistin-based regimen. In strains with a real susceptibility to colistin the use of this drug was associated with long term survival in severe infections.

Finally, interesting data observed in our population show the association between aminoglycoside-containing regimen and 30-day mortality. A possible explanation should be that in a significant proportion of patients (54%), especially septic ones (61%), pneumonia was reported as primary source of infection. As reported above, severe pneumonia remains a difficult-to-treat infection and, considering the poor lung penetration of aminoglycosides, these drugs should be probably avoided for MDR AB pneumonia.^{39,40}

Our study reveals some limitations that should be acknowledged. Firstly, the observational nature of the study brings about an intrinsic limitation in the analysis. Secondly, overall incidence of BSI caused by MDR-AB strains was very inhomogeneous in the involved centres. Finally, the underlying mechanisms of resistance in these strains were not routinely assessed and *in vitro* synergistic combinations were not performed, except for few cases. Further analysis about *in vitro* activity of antibiotic regimens have been avoided, considering that 91 (32.3%) of patients were treated for a previous MDR infections with similar antibiotic regimens. Moreover, antibiotic regimens with two or more drugs displaying *in vitro* activity against MDR-AB have been observed in <10% of strains. However, this real-life clinical experience from prestigious hospitals in Italy provides useful suggestions to clinicians about management of this difficult-to-treat infection.

In conclusion, BSI caused by MDR-AB strains represents a challenge for physicians, considering the high rates of septic shock and mortality associated with this infection. Our data showed peculiar clinical features and use of different antibiotic regimens in this setting of infection, with a predominant role for colistin. The lack of scientific data might explain the very high mortality rate observed in this population of patients.^{41,42}

Acknowledgments

This work was made with the contribution of SITA (Società Italiana Terapia Antinfettiva).

Funding

No funding.

Conflict of interest

None to declare.

References

- Levy MM, Artigas A, Phillips GS, Rhodes A, Beale R, Osborn T, et al. Outcomes of the Surviving Sepsis Campaign in intensive care units in the USA and Europe: a prospective cohort study. *Lancet Infect Dis* 2012;**12**:919–24.
- Perez F, Endimiani A, Ray AJ, Decker BK, Wallace CJ, Hujer KM, et al. Carbapenem-resistant *Acinetobacter baumannii* and *Klebsiella pneumoniae* across a hospital system: impact of post-acute care facilities on dissemination. *J Antimicrob Chemother* 2010;**65**:1807–18.
- Bassetti M, Righi E, Vena A, Graziano E, Russo A, Peghin M. Risk stratification and treatment of ICU-acquired pneumonia caused by multidrug-resistant/extensively drug-resistant/pandrug-resistant bacteria. *Curr Opin Crit Care* 2018;**24**:385–93.
- http://www.who.int/drugresistance/AMR_Importance/en. Accessed on 6th November 2018.
- Isler B, Doi Y, Bonomo RA, Paterson DL. New Treatment Options against Carbapenem-Resistant *Acinetobacter baumannii* Infections. *Antimicrob Agents Chemother* 2018;**63**(1) doi:10.1128/AAC.01110-18.
- <https://atlas.ecdc.europa.eu/public/index.aspx>. Accessed on 26th April 2019.
- Perez F, Hujer AM, Hujer KM, Decker BK, Rather PN, Bonomo RA. Global challenge of multidrug-resistant *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2007;**51**:3471–84.
- Opal SM, Calandra T. Antibiotic usage and resistance: gaining or losing ground on infection in critically ill patients? *JAMA* 2009;**302**:2367–8.
- Vázquez-Ucha JC, Maneiro M, Martínez-Gutián M, Buynak J, Bethel CR, Bonomo RA, et al. Activity of the β -Lactamase inhibitor LN-1-255 against carbapenem-hydrolyzing class D β -lactamases from *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2017;**61**(11).
- Hackel MA, Tsuji M, Yamano Y, Echols R, Karlowsky JA, Sahn DF. In Vitro activity of the siderophore cephalosporin, cefiderocol, against a recent collection of clinically relevant gram-negative bacilli from north america and europe, including carbapenem-nonsusceptible isolates (SIDERO-WT-2014 Study). *Antimicrob Agents Chemother* 2017;**61**(9).
- Durand-Réville TF, Guler S, Comita-Prevoir J, Chen B, Bifulco N, Huynh H, et al. ETX2514 is a broad-spectrum β -lactamase inhibitor for the treatment of drug-resistant Gram-negative bacteria including *Acinetobacter baumannii*. *Nat Microbiol* 2017;**2**:17104.
- Busani S, Serafini G, Mantovani E, Venturelli C, Giannella M, Viale P, et al. Mortality in patients with septic shock by multidrug resistant bacteria. *J Intensive Care Med* 2017. doi:10.1177/0885066616688165.
- Russo A, Giuliano S, Ceccarelli G, Alessandri F, Giordano A, Brunetti G, et al. Comparison of septic shock due to multidrug-resistant *Acinetobacter baumannii* or *Klebsiella pneumoniae* carbapenemase-producing *K. pneumoniae* in intensive care unit patients. *Antimicrob Agents Chemother* 2018;**62**(6).
- Dickinson JD, Kollef MH. Early and adequate antibiotic therapy in the treatment of severe sepsis and septic shock. *Curr Infect Dis Rep* 2011;**13**:399–405.
- Rossolini GM, Mantengoli E. Antimicrobial resistance in Europe and its potential impact on empirical therapy. *Clin Microbiol Infect* 2008;**14**:2–8.
- Bassetti M, Carnelutti A, Peghin M. Patient specific risk stratification for antimicrobial resistance and possible treatment strategies in gram-negative bacterial infections. *Expert Rev Anti Infect Ther* 2017;**15**:55–65.
- Paul M, Daikos GL, Durante-Mangoni E, Yahav D, Carmeli Y, Benatar YD, et al. Colistin alone versus colistin plus meropenem for treatment of severe infections caused by carbapenem-resistant Gram-negative bacteria: an open-label, randomised controlled trial. *Lancet Infect Dis* 2018;**18**:391–400.
- Garnacho-Montero J, Dimopoulos G, Poulakou G, Akova M, Cisneros JM, De Waele J, et al. European society of intensive care medicine. task force on management and prevention of *Acinetobacter baumannii* infections in the ICU. *Intensive Care Med*. 2015;**41**:2057–75.
- European Centre for Disease Prevention and Control. *Annual epidemiological report on communicable diseases in Europe*. Stockholm, Sweden: European Centre for Disease Prevention and Control; 2014 <https://ecdc.europa.eu/sites/portal/files/media/en/publications/Publications/AER-VPD-IBD-2014.pdf>.
- Russell JA. Management of sepsis. *N Engl J Med* 2006;**355**:1699–713.
- O'Grady NP, Alexander M, Burns LA, Dellinger EP, Garland J, Heard SO, et al. Healthcare infection control practices advisory committee. Guidelines for the prevention of intravascular catheter-related infections. *Am J Infect Control* 2011;**39**:S1–34.
- Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, et al. The third international consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA* 2016;**315**:801–10.
- European Committee on Antimicrobial Susceptibility Testing (EUCAST). *Breakpoint tables for interpretation of MICs and zone diameters*, Växjö, Sweden: European Committee on Antimicrobial Susceptibility Testing; 2013. version 3.0 http://www.eucast.org/clinical_breakpoints/.
- Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 2012;**18**:268–81.
- Turkoglu M, Mirza E, Tunçcan OG, Erdem GU, Dizbay M, Yağcı M, et al. *Acinetobacter baumannii* infection in patients with hematologic malignancies in intensive care unit: risk factors and impact on mortality. *J Crit Care* 2011;**26**:460–7.
- Freire MP, de Oliveira Garcia D, Garcia CP, Campagnari Bueno MF, Camargo CH, Kono Magri ASG, et al. Bloodstream infection caused by extensively drug-resistant *Acinetobacter baumannii* in cancer patients: high mortality associated with delayed treatment rather than with the degree of neutropenia. *Clin Microbiol Infect* 2016;**22**:352–8.
- Bassetti M, Repetto E, Righi E, Boni S, Diverio M, Molinari MP, et al. Colistin and rifampicin in the treatment of multidrug-resistant *Acinetobacter baumannii* infections. *J Antimicrob Chemother* 2008;**61**:417–20.
- Poulakou G, Bassetti M, Tsiodras S. "Excess mortality" and colistin-tigecycline for extensively drug-resistant *Acinetobacter baumannii* bacteremia. *Crit Care Med* 2015;**43**:e470–1.
- Tumbarello M, De Pascale G, Treccarichi EM, De Martino S, Bello G, Maviglia R, et al. Effect of aerosolized colistin as adjunctive treatment on the outcomes of microbiologically documented ventilator-associated pneumonia caused by colistin-only susceptible gram-negative bacteria. *Chest* 2013;**144**:1768–75.
- Hsieh TC, Chen FL, Ou TY, Jean SS, Lee WS. Role of aerosolized colistin methanesulfonate therapy for extensively-drug-resistant *Acinetobacter baumannii* complex pneumonia and airway colonization. *J Microbiol Immunol Infect* 2016;**49**:523–30.
- Kim YK, Lee JH, Lee HK, Chung BC, Yu SJ, Lee HY, et al. Efficacy of nebulized colistin-based therapy without concurrent intravenous colistin for ventilator-associated pneumonia caused by carbapenem-resistant *Acinetobacter baumannii*. *J Thorac Dis* 2017;**9**:555–67.
- Durante-Mangoni E, Signoriello G, Andini R, Mattei A, De Cristoforo M, Murino P, et al. Colistin and rifampicin compared with colistin alone for

- the treatment of serious infections due to extensively drug-resistant *Acinetobacter baumannii*: a multicenter, randomized clinical trial. *Clin Infect Dis* 2013;**57**:349–58.
33. Ni W, Shao X, Di X, Cui J, Wang R, Liu Y. In vitro synergy of polymyxins with other antibiotics for *Acinetobacter baumannii*: a systematic review and meta-analysis. *Int J Antimicrob Agents* 2015;**45**:8–18.
 34. Zusman O, Avni T, Leibovici L, Adler A, Friberg L, Stergiopoulou T, et al. Systematic review and meta-analysis of in vitro synergy of polymyxins and carbapenems. *Antimicrob Agents Chemother* 2013;**57**:5104–11.
 35. Cheng A, Chuang YC, Sun HY, Sheng WH, Yang CJ, Liao CH, et al. Excess mortality associated with colistin-tigecycline compared with colistin-carbapenem combination therapy for extensively drug-resistant *acinetobacter baumannii* bacteremia: a multicenter prospective observational study. *Crit Care Med* 2015;**43**:1194–204.
 36. Dickstein Y, Lellouche J, Dalak Amar MB, Schwartz D, Nutman A, Daitch V, et al. Treatment outcomes of colistin and carbapenem-resistant *Acinetobacter baumannii* infections: an exploratory subgroup analysis of a randomized clinical trial. *Clin Infect Dis* 2018 [Epub ahead of print]. doi:10.1093/cid/ciy988.
 37. www.eucast.org/ast_of_bacteria/warnings. Accessed 13 of October 2018.
 38. Matuschek E, Ahman J, Webster C, Kahlmeter G. Antimicrobial susceptibility testing of colistin - evaluation of seven commercial MIC products against standard broth microdilution for *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, and *Acinetobacter* spp.. *Clin Microbiol Infect* 2018;**24**:865–70.
 39. Panidis D, Markantonis SL, Boutzouka E, Karatzas S, Baltopoulos G. Penetration of gentamicin into the alveolar lining fluid of critically ill patients with ventilator-associated pneumonia. *Chest* 2005;**128**:545–52.
 40. Boselli E, Breilh D, Djabarouti S, Guillaume C, Rimmelé T, Gordien JB, et al. Reliability of mini- bronchoalveolar lavage for the measurement of epithelial lining fluid concentrations of tobramycin in critically ill patients. *Intensive Care Med* 2007;**33**:1519–23.
 41. Paterson DL, Rogers BA. How soon is now? The urgent need for randomized, controlled trials evaluating treatment of multidrug-resistant bacterial infection. *Clin Infect Dis* 2010;**51**:1245–7.
 42. Wong D, Nielsen TB, Bonomo RA, Pantapalangkoor P, Luna B, Spellberg B. Clinical and pathophysiological overview of *Acinetobacter* Infections: a century of challenges. *Clin Microbiol Rev* 2017;**30**:409–47.