



Correction to: A cluster of disseminated small cortical lesions in MELAS: its distinctive clinical and neuroimaging features

Yu Hongo¹ · Juntaro Kaneko¹ · Hiroki Suga¹ · Daisuke Ishima¹ · Eiji Kitamura¹ · Tsugio Akutsu¹ · Yuya Onozawa² · Naomi Kanazawa¹ · Tomohide Goto³ · Kazutoshi Nishiyama¹ · Takahiro Iizuka¹ 

Published online: 9 April 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Correction to: Journal of Neurology
<https://doi.org/10.1007/s00415-019-09283-3>

The author would like to correct the errors in the publication of the original article. The corrected details are given below for your reading.

In “Clinical features and neuroimaging patterns of SLEs” section, sixth sentence of the fifth paragraph should as:

In patient 11, brain MRIs obtained 16 weeks after the symptoms onset showed extensive gadolinium enhancement along the gyri of the affected cortexes, indicating not only focal disruption of the BBB, but also same subacute stage of all disseminated lesions (Fig. 2g–i).

In “Discussion” section, ninth sentence of the third paragraph should as:

We also reported a patient with m.617G>A who developed recurrent embolic strokes associated with carotid artery stenosis as a phenotype of macroangiopathy [19].

The original article can be found online at <https://doi.org/10.1007/s00415-019-09283-3>.

✉ Takahiro Iizuka
takahiro@med.kitasato-u.ac.jp

- ¹ Department of Neurology, Kitasato University School of Medicine, 1-15-1 Kitasato, Minami-ku, Sagamihara, Kanagawa 252-0374, Japan
- ² Department of Clinical Laboratory, Kitasato University Hospital, Sagamihara, Japan
- ³ Department of Neurology, Kanagawa Children’s Medical Center, Yokohama, Japan