



The Swiss Narcolepsy Scale (SNS) and its short form (sSNS) for the discrimination of narcolepsy in patients with hypersomnolence: a cohort study based on the Bern Sleep–Wake Database

Panagiotis Bargiotas¹ · Anelia Dietmann¹ · Alan G. Haynes² · Ulf Kallweit^{1,3} · Marta Garcia Calle¹ · Markus Schmidt¹ · Johannes Mathis¹ · Claudio L. Bassetti¹

Received: 22 February 2019 / Revised: 5 May 2019 / Accepted: 7 May 2019 / Published online: 24 May 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

Previous studies reported high sensitivity and specificity of the Swiss Narcolepsy Scale (SNS) for the diagnosis of narcolepsy type 1. We used data from the Bern Sleep–Wake Database to investigate the discriminating capacity of both the SNS and the Epworth Sleepiness Scale (ESS) to identify narcolepsy type 1 and type 2 in patients with central disorders of hypersomnolence (CDH) or sleepy patients with obstructive sleep apnea (OSA). In addition, we aimed to develop a simplified version of the SNS. We created the two-item short-form SNS (sSNS), based on the discriminative capability of the models including all possible combinations of the five questions of the SNS. Using the previously published co-efficiencies, we confirmed the high capacity of the SNS in identifying narcolepsy type 1. The updated SNS (based on new co-efficiencies and cutoff) and the sSNS showed high capacity and were both superior to ESS in identifying narcolepsy type 1. The sSNS correlated significantly with the SNS ($r = -0.897$, $p < 0.001$). No scale showed sufficient discrimination for narcolepsy type 2. This is the largest cohort study that confirms the discriminating power of SNS for narcolepsy type 1 in patients with hypersomnolence and the first study to assess its discriminative power for narcolepsy type 2. The easy-to-use and easy-to-calculate short-form scale has a high discriminating power for narcolepsy type 1 and may be used as screening tool, especially among general practitioners, to identify patients and accelerate their referral to a center of expertise.

Keywords Hypersomnolence · Hypersomnia · Screening tool · Excessive daytime sleepiness · Validation

Panagiotis Bargiotas and Anelia Dietmann have equally contributed to this work.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00415-019-09365-2>) contains supplementary material, which is available to authorized users.

✉ Panagiotis Bargiotas
panagiotis.bargiotas@insel.ch

¹ Sleep Wake Epilepsy Center, Department of Neurology, Inselspital University Hospital Bern, University of Bern, Bern, Switzerland

² CTU Bern, and Institute of Social and Preventive Medicine (ISPM), University of Bern, Bern, Switzerland

³ Clinical Sleep and Neuroimmunology, Institute of Immunology, University of Witten/Herdecke, Witten, Germany

Introduction

Narcolepsy belongs to the group of central disorder of hypersomnolence (CDH) and is clinically characterized by excessive daytime sleepiness (EDS), cataplexy (in narcolepsy type I, NT1), hypnagogic or hypnopompic hallucinations, sleep paralysis and sleep fragmentation [1].

Narcolepsy has an estimated prevalence of 0.05% and symptom onset typically peaks during the second decade of life [2, 3].

There is evidence for a delayed referral of patients with narcolepsy to a specialized center [4] and delayed diagnosis of narcolepsy [5, 6] often due to lack of recognition of signs and symptoms and lack of knowledge about CDH in a broad medical community [4].

The correct diagnosis of narcolepsy is based on clinical features and objective measures including multiple sleep

latency test (MSLT), polysomnography, and/or measurement of decreased or absent hypocretin in cerebrospinal fluid supported by HLA DQB1*0602 testing. Questionnaires such as the Ullanlinna Narcolepsy Scale (UNS) [7], Narcolepsy Severity Scale [8] and the Epworth Sleepiness Scale (ESS) [9, 10] are frequently used in screening of EDS as well as evaluation of treatment effects in narcolepsy, but have a limited discriminative capability especially among patients with hypersomnolence [9–12]. ESS allows calculating a score that quantifies daytime sleepiness, respectively how likely participants fall asleep in different passive situations. It consists of eight items with a four-step rating scale. Total score ranges from 0 to 24. Excessive daytime sleepiness is defined by an ESS value ≥ 10 [13, 14].

In 2004, we introduced a new scale, the Swiss Narcolepsy Scale (SNS) [12] aiming to develop a simple, short and specific screening questionnaire for identifying patients with narcolepsy. The SNS is a self-administered narcolepsy screening instrument which contains five questions and assesses the following parameters: (1) the inability to fall asleep; (2) feeling unrefreshed in the morning; (3) taking a nap at noon; (4) weak knees/buckling of the knees during emotions such as laughing, happiness, or anger; and (5) sagging of the jaw during emotions such as laughing, happiness, or anger. A calculated value (with defined multipliers) of < 0 is indicating the presence of narcolepsy [12].

This initial study compared the SNS with the UNS in 57 patients with NT1, 56 with non-narcoleptic hypersomnolence, and 40 healthy controls, and reported high sensitivity (96%) and specificity (98%) for NT1 compared to a similar sensitivity (98%) but lower specificity (56%) of the UNS [12]. This initial study and a recent validation study [15] focused mainly on the diagnostic accuracy of SNS in detecting NT1 against other types of hypersomnolence. Studies comparing the discriminating power of SNS and ESS for NT2 among patients with hypersomnolence are lacking.

In the current study, we aim (1) to assess the capacity of the SNS and the ESS in discriminating NT1 and NT2 in a larger cohort of new patients with disorders of hypersomnolence, (2) to provide an update version of the SNS based on new scoring coefficients and optimal cutoff point and (3) to develop a simplified short form of the SNS to increase practicability in a daily practice of a broad community of physicians. In an effort to increase applicability of the SNS and mainly of the sSNS in a general practitioner's usual practice, we additionally assessed the discriminative power of the scales in an expanded cohort that included also patients suffering from obstructive sleep apnea, one of the most common causes of excessive sleepiness among patients who visit a general practitioner.

Methods

This is a retrospective cohort study based on data from the Bern Sleep–Wake Database (Dietmann et al. submitted). The protocol for the establishment of the database for clinical and research purposes was approved by the local ethics committee (Kantonale Ethikkommission Bern, 2016-00409).

Patients and clinical assessment

For this study, datasets collected between 2001 and 2016 for clinical purposes were used. Patients have been admitted to the Sleep–Wake-Epilepsy Center, Department of Neurology, Inselspital, University Hospital Bern, Bern, Switzerland for evaluation of suspected disorder of hypersomnolence. Patients underwent clinical routine workup including clinical consultations and electrophysiological examinations (polysomnography, multiple sleep latency test, maintenance of wakefulness test, psychovigilance test and actigraphy), all patients filled in a set of questionnaires related to sleep–wake-disorders. Final diagnosis was reviewed for this study by two independent sleep specialists (A.D. and M.G.C.) according to the International Classification of Sleep Disorders (ICSD-3) [16] using medical history from hospital records, laboratory data, electrophysiological workup of subjective complaints (including PSG, MSLT, MWT and actigraphy) and a battery of sleep–wake questionnaires. Patients included in this study were diagnosed with narcolepsy (type 1 and 2) or other disorder of hypersomnolence, including idiopathic hypersomnia, hypersomnolence due to a medical disorder, hypersomnolence associated with a psychiatric disorder, insufficient sleep syndrome, long sleepers, EDS of unknown origin and sleepy (ESS > 10) patients who have completed the SNS scale and were diagnosed with obstructive sleep apnea (Apnea–Hypopnea Index, AHI $> 5/h$).

The assessment of the clinical and epidemiological sleep–wake profile of the patients was based on the “Bern Sleep Questionnaire”. The questionnaire contains 110 questions about demographics, reasons for referral to sleep laboratory, general information about sleep–wake habits and sleep problems, breathing and circulation, parasomnias and potential trigger factors, dreaming, waking-up, tiredness and sleepiness, cataplexy, hallucinations, stress, well-being, drugs, medication, neurological, psychiatric and other comorbidities, as well as information on family history. Furthermore, all questions included in the ESS, SNS and UNS are included in the Bern Sleep Questionnaire.

Statistical model

To assess the diagnostic accuracy of the SNS, sensitivity and specificity were calculated for the cutoff value of 0. Additionally, logistic regression was used to model the effect of the SNS on the probability of NT1 and NT2 separately.

Table 1 Demographic and clinical characteristics of the cohort

Demographic and clinical characteristics	<i>n</i> (%) or median (LQ, UQ)
Age	32.9 (22.8, 47.7)
Gender	
Male	177 (59%)
Female	122 (41%)
ESS mean score (\pm SD)	11.6 (\pm 6.8)
Disorders of hypersomnolence	299 (100%)
NT1	69
NT2	16
Idiopathic hypersomnia	35
Hypersomnia due to a medical disorder	12
Hypersomnia due to a medication or substance	2
Hypersomnia associated with a psychiatric disorder	59
Insufficient sleep syndrome	91
Long sleeper	10
EDS unclear etiology	35

LQ lower quartile, UQ upper quartiles, NT1 narcolepsy type 1, NT2 narcolepsy type 2, EDS excessive daytime sleepiness

Model coefficients of the individual questions in the SNS were also re-assessed using logistic regression. The Hosmer–Lemeshow goodness-of-fit test and the Brier score were subsequently calculated to test for calibration and agreement between diagnosis and prediction, respectively. Complete cases analyses were performed.

To derive the new short-form SNS (sSNS), logistic regression and AUC for all combinations of questions was used to rank the discriminative ability of the combinations of questions. Although models with four and five questions typically showed higher discriminative abilities, a two-question model (including at least one question on cataplexy and one question on awakening) was desirable. The best two-question model was chosen for further analyses. The cutoff for predicting narcolepsy was selected by summing the sensitivity and specificity for each possible predicted value from the model and selecting the cutoff with the largest sum. Cutoffs reported are on the linear predictor scale. Validation of the sSNS for NT1 and the re-parameterized SNS to predict NT1 was assessed via internal bootstrap validation to estimate the optimism in the AUC, Brier score and calibration plot slope and intercept. Briefly, a training sample was drawn and the model was refit in that sample, with Briers score and a test of calibration plot intercept and slope calculated for the training sample. The statistics from that training model were then compared to the original sample for an estimate of the test performance. Optimism was then calculated based on the difference between training and test performance. This procedure was performed 2000 times and the average optimism subtracted from the statistics from the original model to derive a corrected performance.

As an exploratory analysis, the best cutoff for ESS to determine NT1 and NT2 was assessed by calculating sensitivity and specificity at each possible cutoff (each value between the minimum and maximum), summing the sensitivity and specificity and determining the cutoff with the highest sum. AUC was also calculated.

Analyses were performed in Stata 15.1 and R 3.4.2.

Results

Patients

In our dataset, we identified 299 individuals with a disorder of hypersomnolence who have completed the SNS scale. The final cohort consisted of patients with NT1 (30%), NT2 (7%), idiopathic hypersomnia (15%), hypersomnolence due to a medical condition or a substance (5%), hypersomnolence associated with a psychiatric disorder (26%) and hypersomnolence of unknown origin (17%). Mean age was 33 years (range 23–48) and the male/female ratio was 1.45. For NT1 and NT2 mean age was 31 (range 23–42) and 25 years (range 19–34), and the male/female ratio was 2.0 and 2.2, respectively. Table 1 presents the demographic and clinical characteristics of the patients with CDH.

Expanding the cohort of patients who completed the SNS and including not only patients with CDH but also sleepy (ESS > 10) patients with obstructive sleep apnea (AHI > 5/h), we could identify 473 individuals.

Table 2 Sensitivities and specificities of SNS and its short form (sSNS) in discriminating NT1 and NT2 in a cohort of individuals with hypersomnolence

	SNS*		Updated SNS**		sSNS	
	NT1	NT2	NT1	NT2	NT1	NT2
Cutoff	0	0	− 1.83	− 2.75	− 0.68	− 2.26
Sensitivity (95% CI)	0.86 (0.75–0.92)	0.25 (0.10–0.49)	0.93 (0.84–0.97)	0.63 (0.39–0.82)	0.80 (0.69–0.88)	0.44 (0.23–0.67)
Specificity (95% CI)	0.88 (0.83–0.91)	0.71 (0.65–0.76)	0.82 (0.76–0.86)	0.70 (0.64–0.75)	0.92 (0.87–0.95)	0.83 (0.78–0.87)
Hosmer–Lemeshow goodness-of-fit	0.70	0.02	0.39	0.28	0.26	0.00
Brier score	0.87	0.87	0.07	0.05	0.08	0.22
AUC	0.95	0.54	0.96	0.68	0.92	0.82

NT1 narcolepsy type I, NT2 narcolepsy type II, CI confidence interval, AUC area under the estimated curve, SNS Swiss Narcolepsy Scale, sSNS short-form SNS

*SNS in the current cohort based on the previously published scoring coefficients (Sturzenegger and Bassetti 2004)

**SNS in the current cohort based on the updated scoring coefficients




Fig. 1 Distribution of the SNS for NT1. Black points indicate correct predictions, gray points indicate incorrect (false negative) predictions. Sensitivity and specificity values are shown. NT1, narcolepsy type 1; SNS, Swiss Narcolepsy Scale

The discriminating power of the SNS for NT1 and NT2

In our cohort, the SNS showed a sensitivity of 86% and a specificity of 88% in detecting NT1 against other types of hypersomnolence using the previously published scoring coefficients and the published cutoff of zero [12]. A Brier score of 0.87 indicated relatively poor agreement between observed and predicted outcome (NT1), although the Hosmer–Lemeshow test ($p = 0.70$) was not statistically significant suggesting a normal calibration (Table 2). For the detection of NT2 against other types of hypersomnolence the SNS showed a sensitivity of 25% and a specificity of 71% (Table 2). The distribution of the SNS values in patients with NT1 is shown in Fig. 1.

The discriminating power of the updated SNS for NT1 and NT2

Based on this new cohort, we recalculated the published model coefficients for scoring SNS and determined a new

cutoff point of $- 1.83$. The parameterization of the model to predict NT1 (rather than not-NT1, as in the original parameterization) resulted in the following formula:

$$Q1 \times (-0.47) + Q2 \times (-0.83) + Q3 \times 0.58 + Q4 \times 0.56 + Q5 \times 1.45 - 2.75 \geq -1.83.$$

The sensitivity of the updated SNS for detecting NT1 against other types of hypersomnolence was 91% and the specificity 82%. Brier score was 0.07, indicating good agreement between observed and predicted outcomes (NT1). The Hosmer–Lemeshow test was also non-significant ($p = 0.39$) suggesting a normal calibration (Table 2).

The updated SNS showed a sensitivity of 63% and a specificity of 70% in detecting NT2 against other types of hypersomnolence, using the revised scoring coefficients and the revised cutoff of $- 2.75$ (Table 2).

The discriminating power of the simplified form SNS (sSNS) for NT1 and NT2

The two-item simplified form from the SNS (sSNS) was created based on the discriminative capability of the models including all possible combinations of the five SNS questions. Among them, the combination of SNS question 2 (feeling of being unrefreshed in the morning) and question 5 (reports of episodes with muscle weakness in the face/neck related to emotions) showed the highest discriminative capability for NT1 (supplementary table 1 and supplementary table 2). The accuracy for the sSNS in detecting NT1 was comparable to that of the SNS reaching 80% sensitivity and 92% specificity (Table 2). Brier score was 0.08, indicating good agreement, and the Hosmer–Lemeshow test was non-significant ($p = 0.26$), suggesting a good calibration. The sSNS score correlate significantly with the updated SNS ($r = - 0.89$, $p < 0.001$).




Fig. 2 Distribution of the sSNS score for NT1 patients. Black points indicate correct predictions, gray points indicate incorrect (false negative) predictions. Sensitivity and specificity values are shown. NT1, narcolepsy type 1; sSNS, short-form Swiss Narcolepsy Scale




Fig. 3 Grid to predict NT1. Black regions suggest the possibility of NT1. Values in the cell represent probability of NT1 against other types of hypersomnolence. P: probability; NT1, narcolepsy type 1

Based on the sensitivity and specificity of the models to predict NT1, we determined a cutoff point of -0.68 for the sSNS. The parameterization of the model to predict NT1 resulted in the following formula:

$$Q2 \times (-0.82) + Q5 \times 1.70 - 0.74 \geq -0.68.$$

The distribution of the sSNS values in patients with NT1 are shown in Fig. 2. Model parameters are presented in Table 2.

We used the above formula to create an easy-to-use grid, based on the possible answers in sSNS (Q2 and Q5 of the SNS), to assess probability of NT1 diagnosis against other disorders of hypersomnolence (Fig. 3).

None of the combinations among SNS questions showed sufficient capacity in discriminating NT2 within the cohort of patients with hypersomnolence (data not shown).

Table 3 Sensitivities and specificities of ESS in discriminating NT1 and NT2 in the cohort of patients with hypersomnolence

	ESS	
	NT1	NT2
Sensitivity (95% CI)	0.56 (0.46–0.65)	0.96 (0.82–0.99)
Specificity (95% CI)	0.68 (0.61–0.75)	0.07 (0.05–0.11)
Hosmer–Lemeshow goodness-of-fit	0.288	0.225
Brier score	0.210	0.093
AUC	0.681	0.501

ESS Epworth Sleepiness Scale, NT1 narcolepsy type I, NT2 narcolepsy type II, CI confidence interval, AUC area under the curve

The discriminating power of the original SNS, the updated SNS and the sSNS for NT1 among patients with CDH or OSA

We then assessed the discriminating capability of SNS, updated SNS and sSNS to discriminate NT1 among patients with CDH or sleepy patients with obstructive sleep apnea (OSA).

The original SNS, using the previously published scoring coefficients and the published cutoff of zero [12], showed a sensitivity of 83.3% (95% CI 73.1–90.2%) and a specificity of 90.6% (95% CI 86.9–93.2%) in detecting NT1 among patients with CDH or OSA.

The sensitivity of the reparametrized SNS for detecting NT1 among patients with CDH or OSA was 93% (95% CI 84.8–97.0%) and the specificity 82.3% (95% CI 77.9–86.0%).

The sensitivity of the sSNS for detecting NT1 among patients with CDH or OSA was 83% (95% CI 72.9–89.7%), and the specificity 82.6% (95% CI 78.6–86.0%).

The discriminating power of the ESS for NT1 and NT2

In our cohort, the sensitivity of ESS score, using the typical cutoff of sleepiness ($ESS \geq 10$) was 68% and specificity was 56% for the identification of NT1 against other disorders of hypersomnolence (Table 3). Using various ESS cutoffs, finally applying the one ($ESS \geq 18$) with the best AUC, the sensitivity (51%) and specificity (78%) for identifying NT1 remained low.

Similarly, using various ESS cutoffs, applying the one ($ESS \geq 10$) with the best AUC, the sensitivity for identifying NT2 against other disorders of hypersomnolence was 93% but specificity was very low (17%).

Discussion

This is the largest known retrospective cohort study to assess the discriminative capacity of the SNS for narcolepsy. Our data confirmed the previously reported high capacity of the SNS in identifying NT1 against other types of hypersomnolence [12]. In addition, we updated the SNS by recalculating the model coefficients for scoring SNS and defining new cutoff scores for the scale. The capacity of the updated SNS for discriminating NT1 against other CDH was comparable to the capacity of the original SNS.

There is ample evidence for a delayed diagnosis of narcolepsy often due to the absence or lack of recognition of common narcolepsy symptoms, especially cataplexy. Therefore, the implementation of simple, easy-to-use, and reliable questionnaires on the symptom-based suspicion of narcolepsy and its subtypes in primary care may significantly increase the referral accuracy and improve resource utilization by narrowing the differential diagnosis upon referral. In an effort to increase its practicability, we aimed to simplify the SNS and introduced a simplified form (sSNS) which contains only two questions.

The sSNS correlated with the full SNS and demonstrated comparable validity with the SNS in detecting NT1 against other types of hypersomnolence. We created an easy-to-use grid, based on the sSNS formula, to simplify further the prediction of NT1, by selecting the relevant cell.

We then applied the original SNS, the updated SNS and the sSNS in a larger cohort of patients including sleepy patients with obstructive sleep apnea (OSA). The discriminative power of all scales, including the two-question sSNS, remained satisfactory. This increases the applicability of the SNS and mainly of the sSNS in a general practitioner's usual practice, as OSA is one of the most common causes of pathological level of sleepiness among patients who visit a general practitioner.

Our data from the sSNS suggest that the combined reports of episodes with muscle weakness in the face/neck related to emotions and the subjective feeling of restorative night time sleep can better predict the NT1 among patients with hypersomnolence. Indeed, the cataplectic attacks and the restorative nature of sleep are considered, among sleep specialists, typical symptoms for NT (and specifically for NT1), in contrast to the absence of cataplexy in NT2 or to the non-restorative nature of sleep in other CDH such as idiopathic hypersomnia. However, often these two important sleepiness features are not implemented in the standard first-line screening of a sleepy patient.

Furthermore, we compared the capability of the SNS, the sSNS and the ESS to discriminate NT1 against other

disorders of hypersomnolence. Our data further support the superiority of SNS and sSNS against ESS in identifying NT1, even if these higher cutoff ESS scores were used.

Finally, this is the first study to report the poor discriminative capacity of SNS for NT2. Both SNS and the updated SNS showed low capacity in identifying NT2 against other disorders of hypersomnolence. No combination of the five SNS questions showed a satisfactory discriminative ability. Similarly, ESS showed low validity in identifying NT2 against other disorders of hypersomnolence even if the higher cutoff ESS scores were used. These findings are consistent with previous data showing that the distinction between narcolepsy without cataplexy (NT2) and other disorders of hypersomnolence (mainly the idiopathic hypersomnia) remains ambiguous, not seldom due to a diagnosis change over the time [17].

The retrospective approach and the small N number of patients with NT2 consist of two important limitations of our study. Due of the retrospective design of the study, not all patients with CDH have completed the SNS, and, therefore, had to be excluded from the analysis. In addition, the small sample size of NT2 patients may have limited our ability to detect the discriminative power of the scales for NT2 in this cohort. Prospective cohort studies would overcome these limitations.

Conclusions

The SNS is a useful and valid complementary tool for the diagnosis of NT1 against other types of hypersomnolence. In this study, we introduce its short form (sSNS), a two-question, simple, easy-to-use, easier-to-calculate and reliable questionnaire in particular to be used in primary care as a screening tool for narcolepsy in patients with hypersomnolence. This could decrease the delay and increase the accuracy of referral of patients with hypersomnolence to a specialized sleep center for narcolepsy-specific diagnostic. Finally, our data suggest that SNS and ESS are not the ideal tools for the discrimination of NT2 against other disorders of hypersomnolence.

Although further confirmatory studies and most importantly prospective studies on clinical biomarkers for disorders of hypersomnolence are needed, our data could be used for the improvement of diagnostic processes in these conditions and the development of more specific screening scales in the future.

Acknowledgements We thank Corinne Roth, Sandra Röthlisberger and Tanja Gerber for their assistance with the data transfer to the Bern Sleep–Wake Database. We also thank Christian Sturzenegger for the consultation.

Compliance with ethical standards

Conflicts of interest The authors thank Jazz Pharmaceuticals for funding of this study. Jazz Pharmaceuticals also reviewed the manuscript and had the opportunity to provide suggestions to the authors for their consideration. Although Jazz Pharmaceuticals reviewed the content of this manuscript, the ultimate interpretation, and the decision to submit it for publication was made by the authors independently. “Jazz pharmaceuticals holds a royalty-bearing non-exclusive license to the SNS from the University of Bern”. *Panagiotis Bargiotas* has no specific conflict of interest with respect of present work. Dr. Bargiotas has received congress fees and travel reimbursements from Lundbeck Foundation. *Anelia Dietmann, Marta Garcia Calle, Markus Schmidt, and Johannes Mathis* have no specific conflict of interest with respect of present work and have nothing to declare. *Alan G. Haynes* is affiliated with CTU Bern, University of Bern, which has a staff policy of not accepting honoraria or consultancy fees. However, CTU Bern is involved in design, conduct, or analysis of clinical studies funded by not-for-profit and for-profit organizations. In particular, pharmaceutical and medical device companies provide direct funding to some of these studies. For an up-to-date list of CTU, Bern’s conflicts of interest see “https://www.ctu.unibe.ch/research/declaration_of_interest/index_eng.html”. *Ulf Kallweit* has received Honoraria for consultancy/advisory board and/or speaking engagements from: AOP Orphan Pharma; Bioprojet Pharma; Harmony Biosciences; Jazz Pharma; UCB Pharma. *Claudio Bassetti* has no specific conflict of interest with respect of present work. Prof. Bassetti has received honoraria for consultancy, lectures, and board memberships from the following institutions/companies: Jazz, Servier, UCB, Zambon, Cephalon Lundbeck, Pfizer Boehringer Ingelheim. His research is currently supported by grants of the following institutions/companies: Swiss National Science Foundation (SNF), ResMed, Respiroics, Vifor Pharma, UCB Pharma, Schweizerische Herzstiftung, Tropos Stiftung, Parkinson Schweiz.

References

- Scammell TE (2015) Narcolepsy. *N Engl J Med* 373(27):2654–2662. <https://doi.org/10.1056/NEJMra1500587>
- Dauvilliers Y, Montplaisir J, Molinari N, Carlander B, Ondze B, Besset A, Billiard M (2001) Age at onset of narcolepsy in two large populations of patients in France and Quebec. *Neurology* 57(11):2029–2033
- Ohayon MM, Ferini-Strambi L, Plazzi G, Smirne S, Castronovo V (2005) How age influences the expression of narcolepsy. *J Psychosom Res* 59(6):399–405. <https://doi.org/10.1016/j.jpsychores.2005.06.065>
- Morrish E, King MA, Smith IE, Shneerson JM (2004) Factors associated with a delay in the diagnosis of narcolepsy. *Sleep Med* 5(1):37–41
- Thorpy MJ (1992) The clinical use of the multiple sleep latency test. The standards of practice committee of the American Sleep Disorders Association. *Sleep* 15(3):268–276
- Luca G, Haba-Rubio J, Dauvilliers Y, Lammers GJ, Overeem S, Donjacour CE, Mayer G, Javidi S, Iranzo A, Santamaria J, Peraita-Adrados R, Hor H, Kutalik Z, Plazzi G, Poli F, Pizza F, Arnulf I, Lecendreux M, Bassetti C, Mathis J, Heinzer R, Jennum P, Knudsen S, Geisler P, Wierzbicka A, Feketeova E, Pfister C, Khatami R, Baumann C, Tafti M, European Narcolepsy N (2013) Clinical, polysomnographic and genome-wide association analyses of narcolepsy with cataplexy: a European Narcolepsy Network study. *J Sleep Res* 22(5):482–495. <https://doi.org/10.1111/jsr.12044>
- Hublin C, Kaprio J, Partinen M, Koskenvuo M, Heikkila K (1994) The Ullanlinna Narcolepsy Scale: validation of a measure of symptoms in the narcoleptic syndrome. *J Sleep Res* 3(1):52–59
- Dauvilliers Y, Beziat S, Pesenti C, Lopez R, Barateau L, Carlander B, Luca G, Tafti M, Morin CM, Billiard M, Jaussent I (2017) Measurement of narcolepsy symptoms: the Narcolepsy Severity Scale. *Neurology* 88(14):1358–1365. <https://doi.org/10.1212/WNL.0000000000003787>
- Hublin C, Kaprio J, Partinen M, Koskenvuo M, Heikkila K, Koskimies S, Guilleminault C (1994) The prevalence of narcolepsy: an epidemiological study of the Finnish Twin Cohort. *Ann Neurol* 35(6):709–716. <https://doi.org/10.1002/ana.410350612>
- Johns MW (2000) Sensitivity and specificity of the multiple sleep latency test (MSLT), the maintenance of wakefulness test and the epworth sleepiness scale: failure of the MSLT as a gold standard. *J Sleep Res* 9(1):5–11
- Wing YK, Li RH, Ho CK, Fong SY, Chow LY, Leung T (2000) A validity study of Ullanlinna Narcolepsy Scale in Hong Kong Chinese. *J Psychosom Res* 49(5):355–361
- Sturzenegger C, Bassetti CL (2004) The clinical spectrum of narcolepsy with cataplexy: a reappraisal. *J Sleep Res* 13(4):395–406. <https://doi.org/10.1111/j.1365-2869.2004.00422.x>
- Bloch KE, Schoch OD, Zhang JN, Russi EW (1999) German version of the Epworth Sleepiness Scale. *Respiration* 66(5):440–447. <https://doi.org/10.1159/000029408>
- Vignatelli L, Plazzi G, Barbato A, Ferini-Strambi L, Manni R, Pompei F, D’Alessandro R (2003) Italian version of the Epworth Sleepiness Scale: external validity. *Neurol Sci* 23(6):295–300. <https://doi.org/10.1007/s100720300004>
- Sturzenegger C, Baumann CR, Lammers GJ, Kallweit U, van der Zande WL, Bassetti CL (2018) Swiss Narcolepsy Scale: a simple screening tool for hypocretin-deficient narcolepsy with cataplexy. *Clin Transl Neurosci* 2(2):2514183X18794175
- AASM (2007) Manual for the scoring of sleep and associated events: rules TaTS, 1st edn. American Academy of Sleep Medicine, Westchester
- Trotti LM, Staab BA, Rye DB (2013) Test–retest reliability of the multiple sleep latency test in narcolepsy without cataplexy and idiopathic hypersomnia. *JCSM* 9(8):789–795. <https://doi.org/10.5664/jcsm.2922>