



Correction to: Female sexual dysfunction in Egyptian women with anxiety: prevalence and patterns

A. E. Arafa¹ · S. A. Senosy¹

Published online: 18 July 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Correction to: J Public Health: From Theory to Practice
<https://doi.org/10.1007/s10389-018-0896-6>

In table 4: column 3, row 1 - $n = 78$ (%) should be 35.0 ± 7.4 .
The correct table is shown below.

Table 4 Socio-demographic, obstetric and clinical characteristics associated with female sexual dysfunction in the participating women with anxiety

Characteristic	No dysfunction $n = 41$ (%)	Dysfunction $n = 78$ (%)	p value	
Age (mean \pm sd) in years	30.1 ± 5.6	35.0 ± 7.4	$< 0.001^*$	
Parity			0.269	
	≤ 2	20(48.8)	32(41.0)	
	> 2	21 (51.2)	46 (59.0)	
Education	Literate	34 (82.9)	57 (73.1)	0.165
	Illiterate	7 (17.1)	21 (26.9)	
Menstrual cycle	Regular	35 (85.4)	43 (55.1)	0.001*
	Irregular	6 (14.6)	35 (44.9)	
Menarche age (mean \pm sd) in years	12.7 ± 1.5	12.9 ± 1.8	0.447	
Marriage duration (mean \pm sd) in years	8.6 ± 4.9	12.3 ± 6.4	0.002*	

*p value significant

The original article has been corrected.

The online version of the original article can be found at <https://doi.org/10.1007/s10389-018-0896-6>

✉ S. A. Senosy
shoshoahmed80@yahoo.com; shaymaa.sayed@med.bsu.edu.eg

¹ Department of Public Health and Community Medicine, Faculty of Medicine, Beni-Suef University, Beni-Suef, Mukbel 62511, Egypt