



Anti-Tumour Treatment

Management of adverse events associated with tyrosine kinase inhibitors: Improving outcomes for patients with hepatocellular carcinoma

Lorenza Rimassa^a, Romano Danesi^b, Tiziana Pressiani^a, Philippe Merle^{c,*}^a Medical Oncology and Hematology Unit, Humanitas Cancer Center, Humanitas Clinical and Research Center – IRCCS, Via Manzoni 56, 20089 Rozzano, Milan, Italy^b Clinical Pharmacology and Pharmacogenetics Unit, Department of Clinical and Experimental Medicine, University of Pisa, Via Roma 67, Pisa, Italy^c Department of Hepatology, Hôpital de la Croix-Rousse, Hospices Civils de Lyon, Université Lyon 1, 103 Grande rue de la Croix Rousse, Lyon, France

ARTICLE INFO

Keywords:

Advanced hepatocellular carcinoma
Adverse event management
Cabozantinib
Lenvatinib
Regorafenib
Sorafenib

ABSTRACT

Hepatocellular carcinoma (HCC) is the most common form of primary liver cancer. Sorafenib, regorafenib, lenvatinib and cabozantinib are tyrosine kinase inhibitors (TKIs) that target, in part, vascular endothelial growth factor receptors, and are approved in various regions of the world for the treatment of advanced HCC. All these agents are associated with a range of adverse events (AEs) that can have a substantial impact on patients' health-related quality of life. Fatigue, diarrhoea, hand–foot skin reaction, nausea, vomiting, decreased appetite, hypertension and weight loss are among the most common AEs experienced with these four TKIs. In this review, we discuss strategies for the management of these AEs in patients with advanced HCC, with the aim of maximizing treatment benefits and minimizing the need for TKI treatment discontinuation. We also consider potential TKI–drug interactions and discuss the use of TKIs in patients with liver dysfunction or who have experienced tumour recurrence after liver transplantation. Use of appropriate AE management strategies and avoidance of contraindicated drugs should help patients with advanced HCC to achieve optimal outcomes with TKIs.

Introduction

Tyrosine kinase inhibitors (TKIs) that target pro-angiogenic molecules, such as vascular endothelial growth factor receptors (VEGFRs), in the tumour microenvironment induce apoptosis by inhibiting tumour cell proliferation and angiogenesis [1]. TKIs are effective in treating a range of tumour types, including advanced hepatocellular carcinoma (HCC) [2–5]. HCC accounts for the majority of cases of primary liver cancer, which is the sixth most common cancer and the fourth most common cause of cancer-related death worldwide [6]. This is in part due to the fact that many patients with HCC have advanced disease at the time of diagnosis [7,8]. In 2007 sorafenib was the first VEGFR TKI to be approved in advanced HCC; before this there were no approved systemic therapies for advanced HCC, and the prognosis was poor [9]. The first phase 3 trial of sorafenib versus placebo in advanced HCC demonstrated a significant, yet moderate improvement in median overall survival (OS; 10.7 vs. 7.9 months, respectively; hazard ratio [HR], 0.69; 95% confidence interval, 0.55–0.87; $p < 0.001$) [2]. A further phase 3 trial showed a similar benefit of sorafenib over placebo in Asian patients with advanced HCC [10].

In 2016, a phase 3 trial investigated the efficacy of regorafenib versus placebo in patients with advanced HCC who had progressed on

sorafenib. These data demonstrated that regorafenib moderately improved median OS (10.6 months vs. 7.8 months; HR, 0.63; 95% CI: 0.50–0.79; $p < 0.0001$) and progression-free survival (PFS; 3.1 months vs. 1.5 months; HR, 0.46; 95% CI: 0.37–0.56; $p < 0.0001$) compared with placebo [3]. Regorafenib was subsequently approved in 2017 for the treatment of advanced HCC following sorafenib failure.

When well tolerated, treatment with sorafenib followed by regorafenib (at radiologic progression on sorafenib) can result in a median OS of 26 months [11]. However, when this sequence is not possible due to lack of efficacy or intolerance to these drugs, the prognosis remains poor. This has led to the investigation of further treatment options. Two further anti-angiogenic TKIs, lenvatinib and cabozantinib, have recently shown benefit in advanced HCC. A phase 3 trial of lenvatinib demonstrated non-inferiority to sorafenib in the first-line treatment of advanced HCC [4], with a median OS of 13.6 months vs. 12.3 months, respectively (HR, 0.92; 95% CI: 0.79–1.06). Based on these data, lenvatinib was approved in Japan, Europe and the USA as a first-line treatment. Cabozantinib, which in addition to inhibiting VEGFR 1, 2 and 3 also inhibits c-MET and AXL, was evaluated in a phase 3 study versus placebo in patients who had received up to two prior systemic therapies, including sorafenib, and whose disease had progressed following at least one prior treatment. Cabozantinib demonstrated

* Corresponding author.

E-mail address: philippe.merle@inserm.fr (P. Merle).<https://doi.org/10.1016/j.ctrv.2019.05.004>

Received 22 January 2019; Received in revised form 10 May 2019; Accepted 13 May 2019

0305-7372/ © 2019 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

significant improvements versus placebo in median OS (10.2 months vs. 8.0 months, respectively; HR, 0.76; 95% CI: 0.63–0.92; $p = 0.005$), PFS (5.2 months vs. 1.9 months, respectively; HR, 0.44; 95% CI: 0.36–0.52; $p < 0.001$) and objective response rate (ORR; 4% vs. < 1%, respectively; $p = 0.009$) [5]. Based on these data, cabozantinib was approved in Europe and the USA for the second-line treatment of advanced HCC.

All TKIs such as sorafenib, regorafenib, lenvatinib and cabozantinib are associated with various adverse events (AEs) that can negatively impact a patient's health-related quality of life. Effective AE prevention and management are therefore important to maximize treatment benefits. Here, we review the tolerability profiles of TKIs used to treat advanced HCC, with a focus on AE management strategies. We also discuss the implications of comorbidities and drug–drug interactions for TKI tolerability in this patient population.

Methods

We identified AEs associated with sorafenib, regorafenib, lenvatinib or cabozantinib treatment of patients with advanced HCC from the initial phase 3 clinical trials for these agents. AE management strategies were identified by searching PubMed in May 2018 for English-language articles, with no date restrictions, using the following terms:

("adverse events"[title/abstract] OR "side effects"[title/abstract] OR safety[title/abstract] OR tolerability[title/abstract] OR toxicity[title/abstract]) AND (sorafenib OR regorafenib OR lenvatinib OR cabozantinib) AND (carcinoma[title/abstract] OR cancer[title/abstract]) AND (management[title/abstract] OR manage[title/abstract]).

Search results were screened manually to identify relevant articles. Owing to the wealth of sorafenib literature, only articles about sorafenib AE management in HCC were included. For regorafenib, lenvatinib and cabozantinib, articles about AE management in any solid tumours were included. Reference lists from these articles were reviewed for other potentially relevant publications. TKI–drug interaction information was identified from product labels and a literature search.

Search results

In total, 281 articles were identified and evaluated for relevance, and 30 were included in this review. An additional 16 articles that were cited in the search results were also included.

Summary of AEs associated with TKI treatment of advanced HCC

In the four seminal phase 3 trials of sorafenib, regorafenib, lenvatinib and cabozantinib, most patients experienced drug-related AEs (Table 1) [2–5]. The incidence of grade ≥ 3 AEs ranged from 45 to 75%. The most common AEs were generally similar for all four agents, and included fatigue, diarrhoea, hand–foot skin reaction (HFSR), nausea, vomiting, decreased appetite, hypertension and weight loss.

In each of these studies, a large proportion of patients experienced either an AE-related dose reduction or interruption. With sorafenib, AEs led to dose reductions in 26% of patients and dose interruptions in 44%, compared with 7% and 30% of placebo-treated patients, respectively [2]. AE-related dose reductions or interruptions were reported in 68% of patients treated with regorafenib (vs. 31% with placebo) and 62% of patients treated with cabozantinib (vs. 13% with placebo) [3,5]. The trial evaluating lenvatinib versus sorafenib as first-line treatment only reported dose reductions and interruptions owing to treatment-related AEs; these occurred in 37% and 40% of lenvatinib-treated patients, and 38% and 32% of sorafenib-treated patients, respectively [4].

The most common AEs leading to sorafenib dose reductions in the original phase 3 trial were diarrhoea (8%), HFSR (5%) and rash or desquamation (3%) [2]. HFSR (22%) and diarrhoea (10%) were also the most common AEs leading to cabozantinib dose reductions, along

Table 1

Common adverse events in phase 3 trials of tyrosine kinase inhibitors in advanced hepatocellular carcinoma.

Trial details	Adverse events occurring in $\geq 15\%$ of patients	Incidence of grade ≥ 3 adverse events ^{a,b} (%)
Sorafenib:	Any	45
NCT00105443	Diarrhoea	10–11
(N = 602) [2]	Fatigue	10
Sorafenib 400 mg twice daily	Abdominal pain	9
	Weight loss	2
	Decreased appetite	3
	Nausea	1
	Ascites	6–7
	HFSR	8
	Rash/desquamation	1
	Oedema, limb	3
	Vomiting	2
Regorafenib:	Any	66
NCT01774344	HFSR	13
(N = 573) [3]	Diarrhoea	3
Regorafenib 160 mg daily during weeks 1–3 of each 4-week cycle	Fatigue	9
	Hypertension	15
	Decreased appetite	3
	Increased blood bilirubin	11
	Abdominal pain	3
	Increased AST	11
	Nausea	1
	Constipation	< 1
	Ascites	4
	Anaemia	5
	Limb oedema	1
	Increased ALT	4
	Hypoalbuminaemia	2
Lenvatinib:	Any	75
NCT01761266	Hypertension	23
(N = 954) [4]	Diarrhoea	4
Lenvatinib 12 mg daily for bodyweight ≥ 60 kg or 8 mg daily for bodyweight < 60 kg	Decreased appetite	5
	Weight loss	8
	Fatigue	4
	HFSR	3
	Proteinuria	6
	Dysphonia	< 1
	Nausea	1
	Decreased platelet count	5
	Abdominal pain	2
	Vomiting	1
	Constipation	1
	Increased blood bilirubin	7
Cabozantinib:	Any	68
NCT01908426	Diarrhoea	10
(N = 707) [5]	Decreased appetite	6
Cabozantinib 60 mg daily	HFSR	17
	Fatigue	10
	Nausea	2
	Hypertension	16
	Vomiting	< 1
	Increased AST	12
	Asthenia	7
	Dysphonia	1
	Constipation	< 1
	Abdominal pain	2
	Weight loss	1
	Increased ALT	5

ALT: alanine aminotransferase; AST: aspartate transaminase; HFSR: hand–foot skin reaction.

^a Adverse event grading from different versions of the National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE) were used for sorafenib (version 3.0), regorafenib (version 4.03), lenvatinib (version 4.0) and cabozantinib (version 4.0).

^b Incidences of grade ≥ 3 adverse events for sorafenib, regorafenib and cabozantinib were calculated by summing grade 3 and grade 4 events.

Table 2

A general algorithm for dose interruptions and reductions according to the severity of tyrosine kinase inhibitor-associated adverse events.

Grade or severity of AE	Treatment	Modification of dose
Grade 0, 1 (mild) or 2 (moderate)	Unchanged	No modification
Grade 3 (severe)	Interruption until recovery to < grade 2	First episode: try to restart at full dose Second episode: reduce the dose until the AE has reduced in severity by 1 grade
Grade 3–4	As there is a risk of death, consider a permanent reduction or interruption	

AE: adverse event.

with fatigue (7%), hypertension (7%) and increased aspartate transaminase levels (6%) [5]. AEs leading to dose reductions and interruptions were not reported for regorafenib or lenvatinib [3,4]. In addition to AE-related dose reductions or interruptions, a smaller but substantial proportion of patients discontinued treatment and withdrew from each study due to AEs. Discontinuations owing to treatment-related AEs occurred in similar proportions of patients treated with sorafenib (11%), regorafenib (10%), lenvatinib (9%) and cabozantinib (16%) [2–5]. AE management and minimization may improve patients' quality of life and reduce treatment discontinuation.

To ensure that treatment efficacy is optimal, maintaining the dose intensity recommended on each TKI's label is of major interest. Dose reductions or treatment interruptions should be considered on a case-by-case basis according to the severity and nature of each AE. Given that such strategies are specific to each TKI, a general algorithm for TKI-associated AE management is provided in Table 2.

Association of TKI AEs with positive treatment outcomes

Some TKI-associated AEs correlate with improved patient outcomes; sorafenib-related dermatological AEs are associated with improved survival [12–17]. A prospective study in 147 sorafenib-treated patients with advanced HCC demonstrated that those who experienced dermatological AEs within the first 60 days of treatment had significantly increased time to progression (8.1 months vs. 3.9 months; $p = 0.016$) and OS (18.2 months vs. 10.1 months; $p = 0.009$), compared with those who did not [14]. A retrospective, bivariate analysis of patients enrolled in the phase 3 study of regorafenib showed that HFSR is also associated with better outcomes for regorafenib-treated patients [18].

TKI-related diarrhoea and hypertension have also been associated with positive outcomes in advanced HCC [12,13,19,20]. An observational study of patients who experienced diarrhoea or hypertension after sorafenib treatment showed significant improvements in OS compared with those who did not (median OS: diarrhoea, 14 months vs. 7 months; hypertension, 13 months vs. 8 months; for patients with and without each AE, respectively) [13]. Although the association between AEs and outcomes in advanced HCC has not been assessed for lenvatinib, median OS for patients with differentiated thyroid cancer treated with lenvatinib was significantly longer for patients who experienced grade ≥ 2 hypertension than for those who did not (not reached vs. 21.7 months) [21]. Currently, there are no data available on any associations between AEs and outcomes of cabozantinib treatment in any tumour type.

Occurrence of HFSR, diarrhoea and hypertension are all likely to be related to the mechanism of action of TKIs. Inhibition of platelet-derived growth factor receptors (PDGFRs) and c-Kit in the eccrine glands of the dermis and epidermis of the palms and soles may contribute to HFSR [22,23]. A decrease in peripheral blood flow following sorafenib treatment has also been linked to the emergence of this AE [24]. Hypertension may result from TKI inhibition of vascular endothelial growth factor (VEGF)-mediated nitric oxide synthase upregulation, thereby inhibiting vasomotor function and promoting degeneration of small blood vessels [25]. VEGF and VEGFRs are also expressed in intestinal endothelial cells, and c-Kit is important for the function of interstitial cells of Cajal in the gastrointestinal tract [26,27].

Consequently, it has been suggested that VEGFR and c-Kit inhibition may cause diarrhoea [28]. Management of HFSR, diarrhoea and hypertension is especially important; although they are associated with positive treatment outcomes, they are among the most frequently experienced AEs [2–5]. Furthermore, if TKI tolerability can be improved, patients are more likely to remain on treatment and therefore experience better outcomes [29,30].

Management of AEs associated with TKI therapy

Over the last decade, a wealth of strategies have been developed for the management of sorafenib-associated AEs. Most of these strategies can be used in the management of AEs associated with other TKIs. Moreover, many AEs associated with regorafenib, lenvatinib and cabozantinib in the treatment of advanced HCC are consistent with AEs observed in the treatment of other solid tumours [31–35]. Consequently, where information is lacking for managing AEs observed in the treatment of advanced HCC, AE management approaches that are established for other tumour types can be considered here. Management recommendations for the most common TKI-associated AEs are discussed below and summarized in Table 3.

Hand-foot skin reaction

TKI-associated HFSR (also known as palmar–plantar erythrodysesthesia or hand–foot syndrome) generally affects the palms of the hands and soles of the feet. Occasionally, symptoms may also occur on other areas such as the knees and elbows. HFSR of any grade was the most common AE associated with regorafenib in the phase 3 advanced HCC study, occurring in 53% of patients (Table 1) [3]. It was also among the most common AEs in the sorafenib, lenvatinib and cabozantinib phase 3 advanced HCC studies (occurring in 21%, 27% and 46% of patients, respectively) [2,4,5]. Grade 3 HFSR, which can substantially impair patients' ability to perform daily tasks and overall quality of life, occurred in 3–17% of patients in these studies (Table 1) [2–5].

Many recommendations for managing HFSR are based on clinical experience rather than scientific evidence. For patients with advanced HCC, most recommendations are based on experience with sorafenib; however, these are consistent with recommendations made for regorafenib, cabozantinib and lenvatinib in other tumour types (Table 3). Before starting TKI treatment, it is important to educate patients about the symptoms of HFSR, and to take steps to soften and remove any existing areas of hyperkeratosis or callused skin on patients' hands and feet [36–42]. Patients should be regularly monitored to proactively manage any occurrence of skin toxicities once TKI treatment has started, especially at the start of therapy (e.g. every few weeks for the first 2–4 months) [36,37,42–44]. To minimize the development of calluses and hyperkeratosis, patients should use emollients regularly [36,39,41,42,45–49]. In sorafenib-treated patients with advanced HCC, prophylactic emollients containing 10% urea, used three times a day, have been shown to reduce the incidence and delay the onset of HFSR [48]. Minimizing unnecessary friction can be achieved by wearing cotton socks and gloves, padded insoles and well-fitting shoes [37,39,41,42,46,47,50] and avoiding heavy lifting [46]. Other

Table 3
Management of common adverse events associated with tyrosine kinase inhibitors for advanced hepatocellular carcinoma.

Adverse event	Recommended management strategies
HFSR	Prophylactic management: - Existing areas of hyperkeratosis or callused skin should be softened and removed before TKI initiation - Emollients containing 10% urea should be used three times daily - Hot water should be avoided - Non-fragranced, non-foaming skin products should be used, hand sanitizers containing alcohol should be avoided; hands should be completely dried after washing - Sun exposure and unprotected cold exposure should be avoided - Friction may be minimized using cotton socks and gloves, padded insoles and well-fitting shoes, and by avoiding heavy lifting AE management: - Preventative measures should be continued - Use emollients containing 20–40% urea - Moisturizers containing salicylic acid, ammonium lactate or alpha hydroxyl acid may be used to soften and exfoliate hyperkeratotic and callused areas - Cooling hand and foot baths containing magnesium sulphate may reduce pain and soften calluses - For grade 2/3 HFSR, consider using topical treatments such as cortisone and 0.05% clobetasol - If topical treatments are ineffective, consider using oral analgesia such as pregabalin and opioids (use with caution) - To prevent infection, cracked skin may be soaked in equal parts vinegar and water for 10 min each day - In severe cases it may be necessary for a trained professional to drain blisters and remove areas of hyperkeratosis
Diarrhoea	Prophylactic management: - Using a stool diary to help identify foods that may trigger digestive problems may be useful AE management: - Caffeine, alcohol, spicy or fatty foods, dairy products and foods high in insoluble fibre should be avoided - Consumption of bananas, rice, potatoes, apple sauce, toast and probiotics may be helpful - Concomitant lactulose dose reduction may be necessary - Fluid intake should be increased, and electrolytes should be monitored and replaced when necessary - In cases that cannot be managed by dietary changes, loperamide (4 mg then 2 mg every 4 h, or after each loose stool until the desired effect is achieved) may be prescribed. For patients who frequently experience diarrhoea, loperamide may also be taken pre-emptively, 30 min before TKI treatment - Loperamide-refractory diarrhoea may be treated with atropine-diphenoxylate, codeine or tincture of opium, if appropriate
Fatigue	AE management: - Rest periods should be incorporated into the patient's daily schedule - For patients who are fit enough, daily exercise such as walking, or weight-bearing exercises may be useful - A nutritious diet and proper hydration should be encouraged - The presence of underlying conditions including hypothyroidism, low testosterone in men (via an endocrinology consultation), pain, sleep dysfunction, emotional distress, depression and anaemia should be evaluated and treated as required. However, care should be taken with the administration of testosterone, due to the potential hormone-sensitivity of HCC cells. - Treatment of other adverse events such as diarrhoea, vomiting, nausea and weight loss may reduce fatigue - Psychostimulants such as caffeine, or methylphenidate or modafinil for more severe cases, may be considered; however, care should be taken when prescribing modafinil owing to potential interactions with TKIs - TKI dosing in the evening rather than the morning may reduce daytime fatigue
Nausea and vomiting	AE management: - Chocolate, caffeine, alcohol and nicotine should be avoided - Pharmacological treatment with metoclopramide or levosulpiride may be considered - Ondansetron and granisetron should be used with caution owing to potential interactions with TKIs
Decreased appetite and weight loss	Prophylactic management: - Appetite and weight should be monitored in each treatment cycle - A nutritious diet should be encouraged AE management: - Appetite stimulants such as dronabinol or megestrol acetate should be considered - Any underlying nausea should be treated - A high-calorie diet and dietary supplements should be recommended
Hypertension	Prophylactic management: - Blood pressure should be controlled before initiating TKI treatment - Blood pressure should be monitored regularly for the first few months of treatment AE management: - Angiotensin-converting enzyme inhibitors, angiotensin receptor blockers or beta blockers should be used to treat hypertension - Calcium channel blockers may be considered, but careful selection is necessary to avoid interactions with TKIs - Caution should be taken when using thiazide diuretics owing to the risk of diarrhoea

AE: adverse event; HFSR: hand–foot skin reaction; NSAIDs: non-steroidal anti-inflammatory drugs; TKI: tyrosine kinase inhibitor.

preventative strategies include avoiding hot water [41–43,51] and hand sanitizers containing alcohol [42], using non-fragranced, non-foaming skin cleansers [42,46,47], ensuring that hands are dried completely after washing, and avoiding sun exposure and unprotected exposure to cold [51].

Once HFSR develops, preventative measures should be continued. Emollients containing 20–40% urea may be used, and additional topical treatments should be considered [50,52]. Moisturizers containing salicylic acid, ammonium lactate or alpha hydroxyl acid may be useful to

soften and exfoliate hyperkeratotic and callused areas [36,37,39–41,45,50]. Cooling hand and foot baths containing magnesium sulphate are also suggested to reduce pain and soften calluses, making them easier to remove [37,40,43,47,51]. Topical cortisone, corticosteroids such as 0.05% clobetasol, and topical analgesia that includes lidocaine are commonly recommended for symptomatic relief of grade 2/3 HFSR [38,39,42,43]. If these are not effective, oral analgesics including non-steroidal anti-inflammatory drugs (NSAIDs), pregabalin and opioids may be considered with caution [36,44,52,53].

To prevent infection of cracked and painful skin, patients may be advised to soak the affected area in a solution of equal parts vinegar and water for 10 min twice a day [41]. In cases where severe hyperkeratosis and blistering occur, a trained professional should drain the blisters and remove areas of hyperkeratosis using sterile instruments; antibiotics should be used when infection occurs [41,46]. When severe HFSR does not adequately respond to the methods described above, TKI dose interruptions or reductions should be considered [36,37,39–41,45]. Once severity decreases to grade 1, the full dose can be resumed; however, if this results in worsening of the HFSR, a permanent lower dose may be an option [45,50,52]. Discontinuation of TKI therapy should be considered only in the most severe or recurrent cases of HFSR [37,40,44,51].

Diarrhoea

Diarrhoea was the most common AE associated with sorafenib and cabozantinib, occurring in 55% and 54% of patients with advanced HCC, respectively (Table 1) [2,5]. It was the second most common AE reported in patients who received regorafenib and lenvatinib [3,4]; grade ≥ 3 diarrhoea occurred in 3–11% of these patients (Table 1) [2–5].

Generally, recommendations for TKI-associated diarrhoea management are consistent across tumour types (Table 3). A specific recommendation for patients with advanced HCC is to reduce any dose of concomitant lactulose [43]. Aside from this, to enable rapid reporting and management, patients should be encouraged to complete a stool diary and promptly report any concerns to their healthcare provider [45]. Diarrhoea may also be managed by making dietary changes, including avoiding caffeine, alcohol, spicy or fatty foods, dairy products and foods high in insoluble fibre [36,37,42,43,45,50,54]. Use of a food diary may be helpful for identifying particular items that exacerbate diarrhoea [51]. Additionally, consumption of bananas, rice, potatoes, apple sauce, toast and probiotics may help to alleviate symptoms [28,53]. It is important to ensure that patients with diarrhoea do not become dehydrated; fluid intake should be increased, and electrolytes monitored and replaced when necessary [36,40,43,45,47].

If diarrhoea cannot be managed by dietary changes, pharmacological intervention may be necessary. Loperamide is widely recommended; an initial dose of 4 mg should be followed by 2 mg every 4 h, or after each loose stool until symptoms subside [44,45,47,50,54]. For patients who frequently experience diarrhoea, loperamide may be taken as a pre-emptive measure 30 min before each TKI dose [43–45,51]. When diarrhoea is not controlled by loperamide, other medications such as atropine-diphenoxylate, octreotide, codeine or tincture of opium may be considered [36,37,40,44,50]. Cholestyramine should not be taken in combination with any other anti-diarrhoea medications because of the potential for drug–drug interactions [43,51]. When severe or persistent diarrhoea is unresponsive to management, TKI dose interruptions or reductions should be considered; if tolerated, the full dose can be resumed once symptoms resolve [36,39,41,43,45,49]. Some patients may need to be screened for exocrine pancreatic insufficiency, which can be treated with pancreatic enzyme replacement [55].

Fatigue

Fatigue may develop as a side effect of TKI treatment or may be a symptom of advanced HCC. In phase 3 studies, fatigue occurred in 46% of patients who received sorafenib, 40% of patients who received regorafenib, 30% of patients who received lenvatinib and 45% of those who received cabozantinib (Table 1) [2–5]. Grade ≥ 3 fatigue occurred at rates of 4–10% in these patients (Table 1).

To encourage early reporting and management to prevent worsening, it is important to educate patients about the possibility of developing treatment-related fatigue (Table 3) [42,44,53]. Patients who

experience fatigue should be advised to incorporate rest periods into their daily schedule [42,45,47]. For patients with colorectal cancer, there is evidence to suggest that physical exercise improves fatigue [56]. Although many patients with advanced HCC are not well enough to exercise strenuously, daily activity, such as walking or weightbearing exercises, should be encouraged for those who are able [41–43,45,47,49]. It is also important to ensure that patients are properly nourished and hydrated [36,40,45,49].

Fatigue may also be a consequence of malnutrition caused by other TKI-associated AEs, including diarrhoea, vomiting and nausea; management of these AEs as described in this review may alleviate fatigue. Patients with fatigue should also be assessed and treated for hypothyroidism, which is commonly associated with TKI treatments [36,37,41,43,45]. Other potential underlying causes of fatigue, such as low testosterone in men, pain, sleep dysfunction, emotional distress, depression and anaemia, should also be considered and treated as required [28,37,45,47,49]. Psychostimulants may be considered for treatment of TKI-related fatigue [45,49,50,53]. In particular, methylphenidate and modafinil have shown efficacy in patients with severe cancer-related fatigue [57,58]; however, care should be taken when using modafinil owing to the risk of interaction with TKIs through CYP3A4 (Table 4) [59].

Caffeine could be incorporated into the diet if diarrhoea or nausea and vomiting are not an issue [43,53]. Patient diaries can be useful to determine whether fatigue is cancer-related or treatment-related and may inform dose modifications [47]. To reduce daytime fatigue, patients may find it helpful to take TKIs in the evening [43]. In most cases, dose reductions or interruptions should only be considered for grade ≥ 3 fatigue (which is a major, limiting AE, not resolvable with rest), and the full dose should be resumed once fatigue is resolved [36,42,45].

Nausea and vomiting

In the phase 3 clinical trials discussed here, nausea rates were 24% for sorafenib, 17% for regorafenib, 20% for lenvatinib and 31% for cabozantinib (Table 1) [2–5]. Vomiting occurred in 15%, 13%, 16% and 26% of patients, respectively. In each study, $\leq 2\%$ of patients who received the investigational drug experienced nausea or vomiting at grade ≥ 3 (Table 1).

There is a lack of evidence-based management strategies for TKI-related nausea and vomiting. It has been suggested that avoiding chocolate, caffeine, alcohol and nicotine may be useful (Table 3) [53]. Antiemetics such as metoclopramide, levosulpiride or ondansetron may alleviate symptoms. Ondansetron should be used with caution, as it may cause QT prolongation, which is also an AE that can arise with sorafenib, cabozantinib and lenvatinib therapy [49,60–64]. Based on limited clinical experience from one centre, other pharmacological treatments that may be useful include the analgesic fentanyl and the anti-diarrhoea agent racecadotril [63].

Decreased appetite and weight loss

Decreased appetite occurred in 29% of patients who received sorafenib, 31% of patients who received regorafenib, 34% of patients who received lenvatinib and 48% of those who received cabozantinib (Table 1) [2–5]. Weight loss was experienced by 14–31% of these patients (Table 1) [2–5].

Management recommendations for decreased appetite and weight loss across TKIs and tumour types include monitoring the patient's appetite and weight in each treatment cycle and encouraging a nutritious diet (Table 3) [36,53]. For patients who experience decreased appetite, stimulants such as dronabinol or megestrol acetate may be administered [36,44]. Caution should be taken when treating patients with advanced HCC with megestrol acetate, because there is a risk of venous thrombosis [65]. Treatment of nausea as described above may

Table 4
Examples of CYP3A modulators and P-glycoprotein substrates and modulators.

CYP3A modulators				
Drug class	Strong inhibitors ^a	Moderate inhibitors ^a	Strong inducers ^b	Moderate inducers ^b
Other	Grapefruit juice, grapefruit, carambola (star fruit), Seville oranges, conivaptan,	–	St. John's wort	–
Antifungal agents	Itraconazole, ketoconazole, posaconazole, voriconazole	Clotrimazole, fluconazole	–	–
Antibiotics	Troleandomycin, clarithromycin	Ciprofloxacin, erythromycin	Rifampicin	–
Cardiovascular agents	Diltiazem	Dronedaron, verapamil	–	Bosentan
Immunosuppressants	–	Cyclosporine	–	–
HIV therapies	Cobicistat, indinavir, ritonavir, nelfinavir	–	–	Efavirenz, etravirine
Hepatitis C therapies	Boceprevir, telaprevir	–	–	–
Gastric agents	–	Aprepitant, cimetidine	–	–
Anticonvulsants and other neurologic agents	Nefazodone	Fluvoxamine, tofisopam	Carbamazepine, phenytoin	Modafinil
P-glycoprotein substrates and modulators				
Drug class	Substrates ^c	Inhibitors ^d	Inducers ^e	
Other	–	Grapefruit juice, grapefruit, carambola (star fruit), Seville oranges	St. John's wort	
Antifungal agents	–	Itraconazole	–	
Antibiotics	–	Clarithromycin	Rifampicin	
Anticoagulants	Dabigatran	–	–	
Antihistamines	Fexofenadine	–	–	
Cardiovascular agents	Digoxin	Amiodarone, carvedilol, dronedarone, propafenone, quinidine, verapamil, ranolazine	–	
Immunosuppressants	–	–	Cyclosporine, dexamethasone	
HIV therapies	–	Lopinavir, ritonavir, saquinavir, tipranavir	Tipranavir	
Hepatitis C therapies	–	Telaprevir	–	
Anticonvulsants and other neurologic agents	–	–	Carbamazepine, phenytoin, venlafaxine	

Examples from US FDA Drug Development and Drug Interactions: Table of Substrates, Inhibitors and Inducers [59] and Wessler JD *et al.* 2013 [75].

AUC: area under the curve; FDA, Food and Drug Administration.

^a Drugs that increase the AUC of sensitive index substrates of CYP3A by ≥ 5 -fold (strong) or ≥ 2 to < 5 -fold (moderate) [59].

^b Drugs that decrease the AUC of sensitive index substrates of CYP3A by $\geq 80\%$ (strong) or $\geq 50\%$ to $< 80\%$ (moderate) [59].

^c Drugs that have AUC increased by ≥ 2 -fold with verapamil or quinidine co-administration and that undergo *in vitro* transport by P-glycoprotein expression systems, but that are not extensively metabolized [59].

^d Drugs that increase the AUC of digoxin by ≥ 2 -fold with co-administration and that inhibit P-glycoprotein *in vitro* [59].

^e Drugs that decrease the AUC of digoxin or fexofenadine by > 0.2 -fold [82].

also improve appetite [28]. Some patients may experience asthenia–anorexia–cachexia syndrome, characterized by weight loss, weakness and fatigue. This condition can be treated with corticosteroids, although these may only be effective in the short term [36,66]. Patients who experience weight loss should be assessed for tumour progression and encouraged to consume a high-calorie diet and take dietary supplements [36,53]. In severe cases, if treatment-related, TKI dose reductions may be considered [36].

Hypertension

Hypertension was the most common AE experienced by patients treated with lenvatinib, with any grade hypertension occurring in 42% of patients and grade ≥ 3 hypertension occurring in 23% of patients (Table 1) [4]. Hypertension was also common in patients treated with regorafenib (any grade, 31%; grade ≥ 3 , 15%) and those treated with cabozantinib (any grade, 29%; grade ≥ 3 , 16%; Table 1) [3,5]. The overall proportion of patients who experienced hypertension while receiving sorafenib was not reported because it was below 10%; however, sorafenib-related any grade and grade ≥ 3 hypertension were reported in 5% and 2% of patients, respectively [2]. It should be noted that National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE) version 3 was used for the phase 3 trial of sorafenib, whereas version 4 was used for the other trials discussed here [67,68]. Between these versions, the grading of hypertension was significantly modified, leading to more patients being classified as having

hypertension when NCI CTCAE version 4 was used.

Recommendations for the management of hypertension in a range of tumour types are generally consistent for the four TKIs discussed here (Table 3). Blood pressure should be controlled before TKI treatment is initiated; it should then be monitored regularly for the first few months [21,39,40,42,43,45]. Monitoring can be conducted at clinic visits, and if appropriate, patients may self-monitor at home multiple times per week [40,42,43]. Angiotensin-converting enzyme (ACE) inhibitors, angiotensin receptor blockers (ARBs) and beta blockers are suitable for patients who develop hypertension while receiving TKIs [39,40,42,43,45,49]. Calcium channel blockers (CCBs) are also suitable [40,42,43,49]; however, caution should be taken when choosing which agent to use. Diltiazem and verapamil are non-dihydropyridine CCBs that inhibit CYP3A4; these and similar agents should be avoided with regorafenib and cabozantinib owing to potential drug–drug interactions (Table 4) [28,53,69]. There is also evidence that the dihydropyridine CCB nifedipine may increase VEGF secretion [70], which is undesirable, as it may counteract the effects of TKI treatment. Other dihydropyridine CCBs that do not interact with this pathway should therefore be considered. Thiazide diuretics are also an option [40,44,53], although they may increase the risk of patients experiencing diarrhoea [42]. A combinatorial treatment approach may be required to manage hypertension in some cases, although ACE inhibitors and ARBs should not be combined [71]. When antihypertensives are not effective, TKI dose reductions or interruptions may be necessary [21,38,45,51]. In persistent cases, TKI discontinuation should be considered [38,45,51].

Impact of concomitant medications and comorbidities on TKI AEs in advanced HCC

CYP3A4 modulators

Sorafenib, regorafenib, lenvatinib and cabozantinib are all metabolized by the liver enzyme CYP3A4 [60,61,64,72]; as such, concomitant use of drugs that modulate activity of CYP3A4 can affect tolerability. Pharmacokinetic evaluation showed that administration of the antifungal ketoconazole, a strong CYP3A4 inhibitor, increased plasma exposure of regorafenib and cabozantinib by approximately 33% and 38%, respectively [60,72], although ketoconazole had no clinically relevant effect on sorafenib or lenvatinib pharmacokinetics [64,73]. Based on these findings, strong CYP3A4 inhibitors should be avoided with regorafenib and cabozantinib but may be used with sorafenib and lenvatinib [60,64,72,73]. Commonly used CYP3A4 inhibitors include certain antibiotics, antifungal agents, immunosuppressants and grapefruit juice (Table 4).

Inducers of CYP3A4 activity have the potential to increase TKI metabolism and decrease efficacy. Pharmacokinetic evaluation found that administration of the antibiotic rifampicin, a strong CYP3A4 inducer, led to substantial reductions in plasma exposure of sorafenib (by 37%), regorafenib (50%) and cabozantinib (77%) [60,64,72]; however, a smaller, non-clinically relevant reduction in plasma exposure was seen for lenvatinib (18%) [74]. Commonly used CYP3A4 inducers include the anti-epileptic carbamazepine, modafinil and the herbal supplement St John's wort (Table 4). It is recommended that these agents are avoided with sorafenib, regorafenib and cabozantinib; however, they may be used with lenvatinib.

P-glycoprotein modulators and substrates

Sorafenib and cabozantinib inhibit the transport protein P-glycoprotein (P-gp), and therefore a build-up of P-gp substrates may occur following their administration [60,64]. P-gp substrates, such as the antihistamine fexofenadine, should therefore be administered with caution in patients receiving sorafenib or cabozantinib (Table 4) [60,64,75]. Two active metabolites of regorafenib are P-gp substrates [72]; therefore, concomitant use of P-gp inhibitors or inducers may affect the efficacy and tolerability of regorafenib and should be avoided. A study of 15 healthy adults receiving lenvatinib showed that P-gp inhibition occurred following a single dose of rifampicin, leading to a non-clinically meaningful increase in lenvatinib plasma exposure [74]. No data are available to exclude lenvatinib as an inducer of P-gp; therefore, P-gp substrates should be administered with caution in patients receiving lenvatinib [61]. Commonly used substrates, inducers and inhibitors of P-gp are shown in Table 4.

Gastroprotectants

Proton-pump inhibitors (PPIs) that are commonly used to treat reflux and gastric ulcers work by reducing the amount of acid produced by the stomach thus increasing the pH. There is evidence to suggest that increased stomach pH may impair the acid-dependent absorption of some TKIs. In a phase 2 study of sorafenib plus erlotinib, sorafenib plasma concentration was reduced in patients who were taking concomitant PPIs [76]. Cabozantinib absorption is also acid-dependent; however, concomitant use of the PPI esomeprazole had no effect on cabozantinib plasma exposure [77]. There are no data regarding the effects of PPIs on absorption of regorafenib or lenvatinib. It is therefore suggested that PPIs should be avoided with sorafenib and used with caution in patients treated with regorafenib or lenvatinib. Overall, there is widespread use of PPIs in patients receiving oral TKIs; however, owing to potential negative effects on TKI absorption, they should be used with care.

Liver dysfunction

Patients with advanced HCC often have reduced liver function (Child–Pugh class B or C) resulting in impaired drug metabolism and reduced albumin production. This reduced metabolism can increase the plasma concentration of TKIs that remain unbound to albumin [78,79], which may reduce TKI tolerability. The phase 3 studies discussed here were conducted in patients with relatively good liver function (Child–Pugh A) at screening. A phase 4 study of sorafenib treatment in HCC found no difference in the incidence of drug-related AEs between patients with Child–Pugh A and Child–Pugh B liver function. However, rates of serious AEs and AE-related treatment discontinuation were higher in patients with Child–Pugh B versus A liver function [80]. These findings suggest that implementation of the AE management strategies described in this article may be of particular importance to minimize AE severity and AE-related discontinuation in patients with reduced liver function. Further studies are required to characterize the efficacy and tolerability of regorafenib, lenvatinib and cabozantinib in patients with liver impairment.

Genetic variation in CYP3A4 liver enzyme function may also affect the metabolism of TKIs [81], and may therefore alter their tolerability. Over 40 CYP3A4 genetic variants have been identified, of which at least 14 alter CYP3A4 enzyme levels or activity [81]. Genetic screening for these mutations is not routine in many centres. However, in cases of severe unmanageable toxicity, the presence of functional variants of these enzymes should be considered, even if this is not easily assessable in clinical practice. Alternatively, levels of sorafenib in plasma may be measured using high-performance liquid chromatography/electrospray ionization tandem mass spectrometry, which has the potential to be useful for identifying patients with unusually high plasma drug concentrations [82]. Using this technique, the TKI dose could be modified to achieve the appropriate plasma drug concentration, although this approach has not yet been prospectively evaluated.

Tumour recurrence after liver transplantation

Patients who have undergone liver transplantation to treat HCC may experience tumour recurrence. These patients may be suitable for systemic TKI therapy; however, they were also excluded from the phase 3 trials discussed here [2–5], and therefore no robust data exist regarding TKI efficacy and safety in this population. A 2015 systematic review of case reports and small retrospective studies found that patients with recurrent HCC after liver transplantation who were treated with sorafenib ($n = 76$) experienced improved survival, compared with those who received best supportive care ($n = 54$; weighted mean [83]: 12.1 [9.95] months vs. 3.3 [2.12] months, respectively) [82,84]. However, this review also highlighted that 7/23 included studies reported AEs that required dose reductions and discontinuations in a high proportion of patients. Across the 23 studies, 197 patients received sorafenib (alone or in combination with modified immunosuppressants) after liver transplantation; of these, 42.1% required dose reductions, and 9.6% discontinued treatment due to AEs [82,84].

An additional consideration when using TKIs to treat patients who have undergone liver transplantation is the requirement for long-term immunosuppression. Given that some immunosuppressants, including cyclosporine, have the potential to interact with TKIs (Table 4) [59], careful selection of compatible therapies is important; suitable examples include tacrolimus and mycophenolate mofetil. Overall, there is a need for prospective studies to characterize the efficacy and tolerability of TKIs for recurrent HCC in these patients.

Conclusions

Sorafenib, regorafenib, lenvatinib and cabozantinib have demonstrated efficacy for the treatment of advanced HCC. However, AEs are common with all these treatments, and proactive AE management is

therefore essential to optimize tolerability and patients' quality of life. As discussed here, effective management strategies include a range of pharmacological and non-pharmacological approaches. When considering pharmacological management and treatment for comorbid conditions, it is important to consider potential interactions of concomitant medications with TKIs, as there may be tolerability implications. In cases of unmanageable tolerability issues, comorbidities such as impaired liver function and genetic variations in liver enzyme function should be considered as possible underlying factors. By employing appropriate preventative and management strategies, it should be possible to mitigate many tolerability issues associated with TKI treatment. This may lead to a reduced need for TKI treatment discontinuation and contribute to better outcomes for patients with advanced HCC.

Acknowledgements

The authors thank Heather Booth, DPhil, of Oxford PharmaGenesis, Oxford, UK for providing medical writing support, which was sponsored by Ipsen, Abingdon, UK in accordance with Good Publication Practice guidelines.

Author contributions

All authors made substantial contributions to the conception and design of the study, drafting the article or revising it critically for important intellectual content and final approval of the version to be submitted.

Declaration of interest

Lorenza Rimassa reports non-financial support and personal fees from ArQule and Ipsen, and personal fees from AbbVie, AstraZeneca, Bayer, Celgene, Eisai, Exelixis, Gilead, Lilly and Sirtex Medical.

Romano Danesi reports grants and personal fees from Pfizer, and personal fees from Celgene, Gilead, Novartis and Roche.

Tiziana Pressiani has nothing to disclose.

Philippe Merle reports grants and personal fees from Onxeo, personal fees from Bayer and Ipsen, and non-financial support from Bristol-Myers Squibb and Roche.

References

- [1] Berretta M, Rinaldi L, Di Benedetto F, Lleshi A, De Re V, Facchini G, et al. Angiogenesis inhibitors for the treatment of hepatocellular carcinoma. *Front Pharmacol* 2016;7:428.
- [2] Llovet JM, Ricci S, Mazzaferro V, Hilgard P, Gane E, Blanc JF, et al. Sorafenib in advanced hepatocellular carcinoma. *N Engl J Med* 2008;359:378–90.
- [3] Bruix J, Qin S, Merle P, Granito A, Huang YH, Bodoky G, et al. Regorafenib for patients with hepatocellular carcinoma who progressed on sorafenib treatment (RESORCE): A randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet* 2017;389:56–66.
- [4] Kudo M, Finn RS, Qin S, Han KH, Ikeda K, Piscaglia F, et al. Lenvatinib versus sorafenib in first-line treatment of patients with unresectable hepatocellular carcinoma: A randomised phase 3 non-inferiority trial. *Lancet* 2018;391:1163–73.
- [5] Abou-Alfa GK, Meyer T, Cheng AL, El-Khoueiry AB, Rimassa L, Ryoo BY, et al. Cabozantinib in patients with advanced and progressing hepatocellular carcinoma. *N Engl J Med* 2018;379:54–63.
- [6] Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394–424.
- [7] Huang Y, Wallace MC, Adams LA, MacQuillan G, Garas G, Ferguson J, et al. Rate of non-surveillance and advanced hepatocellular carcinoma at diagnosis in chronic liver disease. *J Clin Gastroenterol* 2018;52:551–6.
- [8] Kudo M, Han KH, Kokudo N, Cheng AL, Choi BI, Furuse J, et al. Liver Cancer Working Group report. *Jpn J Clin Oncol*. 2010;40(Suppl 1):i19–27.
- [9] Bruix J, Sherman M. Practice guidelines committee AASLD. Management of hepatocellular carcinoma. *Hepatology* 2005;42:1208–36.
- [10] Cheng AL, Kang YK, Chen Z, Tsao CJ, Qin S, Kim JS, et al. Efficacy and safety of sorafenib in patients in the Asia-Pacific region with advanced hepatocellular carcinoma: a phase III randomised, double-blind, placebo-controlled trial. *Lancet Oncol* 2009;10:25–34.
- [11] Finn RS, Merle P, Granito A, Huang YH, Bodoky G, Pracht M, et al. Outcomes of sequential treatment with sorafenib followed by regorafenib for HCC: Additional analyses from the phase III RESORCE trial. *J Hepatol* 2018;69:353–8.
- [12] Cho JY, Paik YH, Lim HY, Kim YG, Lim HK, Min YW, et al. Clinical parameters predictive of outcomes in sorafenib-treated patients with advanced hepatocellular carcinoma. *Liver Int* 2013;33:950–7.
- [13] Di Costanzo GG, de Stefano G, Tortora R, Farella N, Addario L, Lampasi F, et al. Sorafenib off-target effects predict outcomes in patients treated for hepatocellular carcinoma. *Future Oncol* 2015;11:943–51.
- [14] Reig M, Torres F, Rodriguez-Lope C, Forner A, Llach N, Rimola J, et al. Early, dermatologic adverse events predict better outcome in HCC patients treated with sorafenib. *J Hepatol* 2014;61:318–24.
- [15] Branco F, Alencar RS, Volt F, Sartori G, Dode A, Kikuchi L, et al. The impact of early dermatologic events in the survival of patients with hepatocellular carcinoma treated with sorafenib. *Ann Hepatol* 2017;16:263–8.
- [16] Shin SY, Lee YJ. Correlation of skin toxicity and hypertension with clinical benefit in advanced hepatocellular carcinoma patients treated with sorafenib. *Int J Clin Pharmacol Ther* 2013;51:837–46.
- [17] Shomura M, Kagawa T, Shiraishi K, Hirose S, Arase Y, Koizumi J, et al. Skin toxicity predicts efficacy to sorafenib in patients with advanced hepatocellular carcinoma. *World J Hepatol* 2014;6:670–6.
- [18] Bruix J, Merle P, Granito A, Huang YH, Bodoky G, Yokosuka O, et al. Hand-foot skin reaction (HFSR) and overall survival (OS) in the phase 3 RESORCE trial of regorafenib for treatment of hepatocellular carcinoma (HCC) progressing on sorafenib. *ASCO GI* 2018. San Francisco, CA, USA.
- [19] Koschny R, Gotthardt D, Koehler C, Jaeger D, Stremmel W, Ganten TM. Diarrhea is a positive outcome predictor for sorafenib treatment of advanced hepatocellular carcinoma. *Oncology* 2013;84:6–13.
- [20] Estfan B, Byrne M, Kim R. Sorafenib in advanced hepatocellular carcinoma: hypertension as a potential surrogate marker for efficacy. *Am J Clin Oncol* 2013;36:319–24.
- [21] Wirth LJ, Tahara M, Robinson B, Francis S, Brose MS, Habra MA, et al. Treatment-emergent hypertension and efficacy in the phase 3 Study of (E7080) lenvatinib in differentiated cancer of the thyroid (SELECT). *Cancer* 2018;124:2365–72.
- [22] Lacouture ME, Reilly LM, Gerami P, Guitart J. Hand foot skin reaction in cancer patients treated with the multikinase inhibitors sorafenib and sunitinib. *Ann Oncol* 2008;19:1955–61.
- [23] Ponten F, Ren Z, Nister M, Westermarck B, Ponten J. Epithelial-stromal interactions in basal cell cancer: the PDGF system. *J Invest Dermatol* 1994;102:304–9.
- [24] Kamimura K, Shinagawa-Kobayashi Y, Goto R, Ogawa K, Yokoo T, Sakamaki A, et al. Effective prevention of sorafenib-induced hand-foot syndrome by dried-bonito broth. *Cancer Manag Res* 2018;10:805–13.
- [25] Vakilav C, Lenihan D, Kurzrock R, Tsimberidou AM. Anti-vascular endothelial growth factor therapies and cardiovascular toxicity: What are the important clinical markers to target? *Oncologist* 2010;15:130–41.
- [26] Sanders KM, Koh SD, Ward SM. Interstitial cells of cajal as pacemakers in the gastrointestinal tract. *Annu Rev Physiol* 2006;68:307–43.
- [27] Wang D, Lehman RE, Donner DB, Matli MR, Warren RS, Welton ML. Expression and endocytosis of VEGF and its receptors in human colonic vascular endothelial cells. *Am J Physiol Gastrointest Liver Physiol* 2002;282:G1088–96.
- [28] Schmidinger M, Danesi R. Management of adverse events associated with cabozantinib therapy in renal cell carcinoma. *Oncologist* 2018;23:306–15.
- [29] Brunot A, Le Roy F, Le Sourd S, M'Sadek A, Duval M, Crouzet L, et al. Implementation of a nurse-driven educational program improves management of sorafenib's toxicities in hepatocellular carcinoma. *Cancer Nurs* 2018;41:418–23.
- [30] Scandura G, Aiello RA, Ali M, Taibi E, Sano MV, Todaro FM, et al. Appropriate management of cutaneous adverse events maximizes compliance with sorafenib treatment: a single-center experience. *Future Oncol* 2012;8:609–15.
- [31] Choueiri TK, Escudier B, Powles T, Tannir NM, Mainwaring PN, Rini BI, et al. Cabozantinib versus everolimus in advanced renal cell carcinoma (METEOR): Final results from a randomised, open-label, phase 3 trial. *Lancet Oncol* 2016;17:917–27.
- [32] Demetri GD, Reichardt P, Kang YK, Blay JY, Rutkowski P, Gelderblom H, et al. Efficacy and safety of regorafenib for advanced gastrointestinal stromal tumours after failure of imatinib and sunitinib (GRID): An international, multicentre, randomised, placebo-controlled, phase 3 trial. *Lancet* 2013;381:295–302.
- [33] Elisei R, Schlumberger MJ, Muller SP, Schoffski P, Brose MS, Shah MH, et al. Cabozantinib in progressive medullary thyroid cancer. *J Clin Oncol* 2013;31:3639–46.
- [34] Grothey A, Van Cutsem E, Sobrero A, Siena S, Falcone A, Ychou M, et al. Regorafenib monotherapy for previously treated metastatic colorectal cancer (CORRECT): An international, multicentre, randomised, placebo-controlled, phase 3 trial. *Lancet* 2013;381:303–12.
- [35] Schlumberger M, Tahara M, Wirth LJ, Robinson B, Brose MS, Elisei R, et al. Lenvatinib versus placebo in radioiodine-refractory thyroid cancer. *N Engl J Med* 2015;372:621–30.
- [36] Sastre J, Argiles G, Benavides M, Feliu J, Garcia-Alfonso P, Garcia-Carbonero R, et al. Clinical management of regorafenib in the treatment of patients with advanced colorectal cancer. *Clin Transl Oncol* 2014;16:942–53.
- [37] Mitchell J, Khokkaz T, McNeal D, Brent L. Adverse event management strategies: Optimizing treatment with regorafenib in patients with metastatic colorectal cancer. *Clin J Oncol Nurs* 2014;18:E19–25.
- [38] Melosky B. Meeting an unmet need in metastatic colorectal carcinoma with regorafenib. *Asia Pac J Oncol Nurs* 2016;3:58–65.
- [39] Markowitz JN, Fancher KM. Cabozantinib: A multitargeted oral tyrosine kinase inhibitor. *Pharmacotherapy* 2018;38:357–69.
- [40] Khan G, Moss RA, Braith F, Saltzman M. Proactive strategies for regorafenib in

- metastatic colorectal cancer: Implications for optimal patient management. *Cancer Manag Res* 2014;6:93–103.
- [41] Grothey A, George S, van Cutsem E, Blay JY, Sobrero A, Demetri GD. Optimizing treatment outcomes with regorafenib: personalized dosing and other strategies to support patient care. *Oncologist* 2014;19:669–80.
 - [42] De Wit M, Boers-Doets CB, Saettini A, Vermeersch K, de Juan CR, Ouwerkerk J, et al. Prevention and management of adverse events related to regorafenib. *Support Care Cancer* 2014;22:837–46.
 - [43] Brose MS, Frenette CT, Keefe SM, Stein SM. Management of sorafenib-related adverse events: A clinician's perspective. *Semin Oncol* 2014;41(Suppl 2):S1–16.
 - [44] Walko CM, Grande C. Management of common adverse events in patients treated with sorafenib: Nurse and pharmacist perspective. *Semin Oncol* 2014;41(Suppl 2):S17–28.
 - [45] Bolondi L, Craxi A, Trevisani F, Daniele B, Di Costanzo GG, Fagioli S, et al. Refining sorafenib therapy: Lessons from clinical practice. *Future Oncol* 2015;11:449–65.
 - [46] Bracarda S, Ruggeri EM, Monti M, Merlano M, D'Angelo A, Ferrau F, et al. Early detection, prevention and management of cutaneous adverse events due to sorafenib: Recommendations from the Sorafenib Working Group. *Crit Rev Oncol Hematol* 2012;82:378–86.
 - [47] Edmonds K, Hull D, Spencer-Shaw A, Koldenhof J, Chrysou M, Boers-Doets C, et al. Strategies for assessing and managing the adverse events of sorafenib and other targeted therapies in the treatment of renal cell and hepatocellular carcinoma: Recommendations from a European nursing task group. *Eur J Oncol Nurs* 2015;16:172–84.
 - [48] Ren Z, Zhu K, Kang H, Lu M, Qu Z, Lu L, et al. Randomized controlled trial of the prophylactic effect of urea-based cream on sorafenib-associated hand-foot skin reactions in patients with advanced hepatocellular carcinoma. *J Clin Oncol* 2015;33:894–900.
 - [49] Resteghini C, Cavalieri S, Galbiati D, Granata R, Alfieri S, Bergamini C, et al. Management of tyrosine kinase inhibitors (TKI) side effects in differentiated and medullary thyroid cancer patients. *Best Pract Res Clin Endocrinol Metab* 2017;31:349–61.
 - [50] Schott E, Ebert MP, Trojan J. Treatment of hepatocellular carcinoma with sorafenib - focus on special populations and adverse event management. *Z Gastroenterol* 2012;50:1018–27.
 - [51] Brose MS, Bible KC, Chow LQM, Gilbert J, Grande C, Worden F, et al. Management of treatment-related toxicities in advanced medullary thyroid cancer. *Cancer Treat Rev* 2018;66:64–73.
 - [52] Lacouture ME, Ciccolini K, Kloos RT, Agulnik M. Overview and management of dermatologic events associated with targeted therapies for medullary thyroid cancer. *Thyroid* 2014;24:1329–40.
 - [53] Gerendash BS, Creel PA. Practical management of adverse events associated with cabozantinib treatment in patients with renal-cell carcinoma. *Onco Targets Ther* 2017;10:5053–64.
 - [54] Grande E, Glen H, Aller J, Argenziano G, Lamas MJ, Ruszniewski P, et al. Recommendations on managing lenvatinib and everolimus in patients with advanced or metastatic renal cell carcinoma. *Expert Opin Drug Saf* 2017;16:1413–26.
 - [55] Hescot S, Vignaux O, Goldwasser F. Pancreatic atrophy – a new late toxic effect of sorafenib. *N Engl J Med* 2013;369:1475–6.
 - [56] Cheville AL, Kollasch J, Vandenberg J, Shen T, Grothey A, Gamble G, et al. A home-based exercise program to improve function, fatigue, and sleep quality in patients with Stage IV lung and colorectal cancer: a randomized controlled trial. *J Pain Symptom Manage* 2013;45:811–21.
 - [57] Jean-Pierre P, Morrow GR, Roscoe JA, Heckler C, Mohile S, Janelins M, et al. A phase 3 randomized, placebo-controlled, double-blind, clinical trial of the effect of modafinil on cancer-related fatigue among 631 patients receiving chemotherapy: A University of Rochester cancer center community clinical oncology program research base study. *Cancer* 2010;116:3513–20.
 - [58] Moraska AR, Sood A, Dakhil SR, Sloan JA, Barton D, Atherton PJ, et al. Phase III, randomized, double-blind, placebo-controlled study of long-acting methylphenidate for cancer-related fatigue: North Central Cancer Treatment Group NCCTG-N05C7 trial. *J Clin Oncol* 2010;28:3673–9.
 - [59] US Food and Drug Administration. Drug development and drug interactions: Table of substrates, inhibitors and inducers. <https://www.fda.gov/Drugs/DevelopmentApprovalProcess/DevelopmentResources/DrugInteractionsLabeling/ucm093664.htm>. [Accessed 10 May 2019].
 - [60] Cabometyx – summary of product characteristics. <https://www.medicines.org.uk/emc/product/4331/smpc>. [Accessed 10 May 2019].
 - [61] Lenvima – summary of product characteristics. <https://www.medicines.org.uk/emc/product/6840/smpc>. [Accessed 10 May 2019].
 - [62] Moffett PM, Cartwright L, Grossart EA, O'Keefe D, Kang CS. Intravenous ondansetron and the QT interval in adult emergency department patients: An observational study. *Acad Emerg Med* 2016;23:102–5.
 - [63] D'Angelo S, Secondulfo M, De Cristofano R, Sorrentino P. Selection and management of hepatocellular carcinoma patients with sorafenib: Recommendations and opinions from an Italian liver unit. *Future Oncol* 2013;9:485–91.
 - [64] Nexavar – summary of product characteristics. <https://www.medicines.org.uk/emc/product/226>. [Accessed 10 May 2019].
 - [65] Loprinzi CL, Kugler JW, Sloan JA, Mailliard JA, Krook JE, Wilwerding MB, et al. Randomized comparison of megestrol acetate versus dexamethasone versus fluoxymesterone for the treatment of cancer anorexia/cachexia. *J Clin Oncol* 1999;17:3299–306.
 - [66] Tassinari D, Maltoni M. Progestagens and corticosteroids in the management cancer cachexia. In: Mantovani G, et al. editors. *Cachexia and wasting: A modern approach*: Springer, Milano; 2006.
 - [67] National Cancer Institute. Common terminology criteria for adverse events version 3.0. https://ctep.cancer.gov/protocolDevelopment/electronic_applications/docs/ctcae3.pdf. [Accessed 10 May 2019].
 - [68] National Cancer Institute. Common terminology criteria for adverse events version 4.0. https://evs.nci.nih.gov/ftp1/CTCAE/CTCAE_4.03/CTCAE_4.03_2010-06-14_QuickReference_5x7.pdf. [Accessed 10 May 2019].
 - [69] Keefe D, Bowen J, Gibson R, Tan T, Okera M, Stringer A. Noncardiac vascular toxicities of vascular endothelial growth factor inhibitors in advanced cancer: a review. *Oncologist* 2011;16:432–44.
 - [70] Miura S, Fujino M, Matsuo Y, Tanigawa H, Saku K. Nifedipine-induced vascular endothelial growth factor secretion from coronary smooth muscle cells promotes endothelial tube formation via the kinase insert domain-containing receptor/fetal liver kinase-1/NO pathway. *Hypertens Res* 2005;28:147–53.
 - [71] James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA* 2014;311:507–20.
 - [72] Stivarga – summary of product characteristics. <https://www.medicines.org.uk/emc/product/1263>. [Accessed 10 May 2019].
 - [73] Shumaker R, Aluri J, Fan J, Martinez G, Thompson GA, Ren M. Effects of ketocazole on the pharmacokinetics of lenvatinib (E7080) in healthy participants. *Clin Pharmacol Drug Dev* 2015;4:155–60.
 - [74] Shumaker RC, Aluri J, Fan J, Martinez G, Thompson GA, Ren M. Effect of rifampicin on the pharmacokinetics of lenvatinib in healthy adults. *Clin Drug Investig* 2014;34:651–9.
 - [75] Wessler JD, Grip LT, Mendell J, Giugliano RP. The P-glycoprotein transport system and cardiovascular drugs. *J Am Coll Cardiol* 2013;61:2495–502.
 - [76] Lind JS, Dingemans AM, Groen HJ, Thunnissen FB, Bekers O, Heideman DA, et al. A multicenter phase II study of erlotinib and sorafenib in chemotherapy-naïve patients with advanced non-small cell lung cancer. *Clin Cancer Res* 2010;16:3078–87.
 - [77] Nguyen L, Holland J, Mamelok R, Laberge MK, Grenier J, Swearingen D, et al. Evaluation of the effect of food and gastric pH on the single-dose pharmacokinetics of cabozantinib in healthy adult subjects. *J Clin Pharmacol* 2015;55:1293–302.
 - [78] Oettl K, Birner-Gruenberger R, Spindelboeck W, Stueger HP, Dorn L, Stadlbauer V, et al. Oxidative albumin damage in chronic liver failure: relation to albumin binding capacity, liver dysfunction and survival. *J Hepatol* 2013;59:978–83.
 - [79] Tortorici MA, Toh M, Rahavendran SV, Labadie RR, Alvey CW, Marbury T, et al. Influence of mild and moderate hepatic impairment on axitinib pharmacokinetics. *Invest New Drugs* 2011;29:1370–80.
 - [80] Lencioni R, Kudo M, Ye SL, Bronowicki JP, Chen XP, Dagher L, et al. GIDEON (Global Investigation of therapeutic DEcisions in hepatocellular carcinoma and Of its treatment with sorafenib): second interim analysis. *Int J Clin Pract* 2014;68:609–17.
 - [81] Werk AN, Cascorbi I. Functional gene variants of CYP3A4. *Clin Pharmacol Ther* 2014;96:340–8.
 - [82] Takasaki S, Tanaka M, Kikuchi M, Maekawa M, Kawasaki Y, Ito A, et al. Simultaneous analysis of oral anticancer drugs for renal cell carcinoma in human plasma using liquid chromatography/electrospray ionization tandem mass spectrometry. *Biomed Chromatogr* 2018;32:e4184.
 - [83] Wulkersdorfer B, Zeitlinger M, Schmid M. Pharmacokinetic aspects of vascular endothelial growth factor tyrosine kinase inhibitors. *Clin Pharmacokinet* 2016;55:47–77.
 - [84] deAngelis N, Landi F, Carra MC, Azoulay D. Managements of recurrent hepatocellular carcinoma after liver transplantation: A systematic review. *World J Gastroenterol* 2015;21:11185–98.