



Research Article

Microarray analysis of transcriptional responses to salt and drought stress in *Arabidopsis thaliana*

Razieh Ghorbani, Abbas Alemzadeh^{*}, Hooman Razi

Department of Crop Production and Plant Breeding, School of Agriculture, Shiraz University, Shiraz, Iran

ARTICLE INFO

Keywords:

Agriculture
Plant biology
Ecology
Genetics
Agronomy
Horticulture
Lagenaria siceraria
Cucurbits
Harvest time
Vivipary
Seed maturation
Yield
Arabidopsis
Meta-analysis
Transcriptomics
Transcription factors

ABSTRACT

Microarray expression profile analysis is a useful approach to increase our knowledge about genes involved in regulatory networks and signal transduction pathways related to abiotic stress tolerance. Salt and drought, as two important abiotic stresses, adversely affect plant productivity in the world every year. To understand stress response mechanisms and identify genes and proteins which play critical roles in these mechanisms, the study of individual genes and proteins cannot be considered as an effective approach. On the other hand, the availability of new global data provides us an effective way to shed some light on the central role of molecules involved in stress response mechanisms in the plant. A meta-analysis of salt and drought stress responses was carried out using 38 samples of different experiments from leaves and roots of *Arabidopsis* plants exposed to drought and salt stresses. We figured out the number of differentially expressed genes (DEGs) was higher in roots under both stresses. Also, we found that the number of common DEGs under both stresses was more in roots and also the number of common DEGs in both tissues under salt stress was more than drought stress. The highest percent of DEGs was related to cell and cell part (about 87%). Around 9% and 7% of DEGs in roots and leaves encoded transcription factors, respectively. Network analysis revealed that three transcription factor families HSF, AP2/ERF and C2H2, may have critical roles in salt and drought stress response mechanisms in *Arabidopsis* and some proteins like STZ may be introduced as a new candidate gene for enhancing salt and drought tolerance in crop plants.

1. Introduction

Plants unlike animals are not able to move and escape from undesirable condition and stresses. Abiotic stresses such as drought, salt, extreme temperature, heavy metals and nutrient deficiency have undesirable effects on plants [1]. These stresses cause decrease in growth and productivity of plants. Among these stresses, drought and salt are two major environmental factors which have detrimental effects on growth, yield and crop production. There is severe water shortage in the world and the majority of these water resources are exposed to salt. Hence, it is necessary to improve water use efficiency and salt tolerance of crops through plant breeding projects using new tools like biotechnology [2].

Plant have responded to salt and drought stresses at the molecular and cellular levels and also at the physiological and biochemical levels [3]. There are different mechanisms evolved in plants to receive and transmit signals from the environment to cells which leads to response to

different stresses [4]. Molecular biology techniques together with bio-informatic studies have provided a powerful tool to identify a number of genes and proteins involving in different pathways of stress response mechanisms (SRMs) [5, 6]. Molecular response of plants to environmental stresses may actually involve a complex interaction between different molecules related to various pathways. Meta-analysis is the statistical approach which combine results from different studies to clarify simultaneously expression of thousands of genes and could be used to detect key genes in response to different stresses [7]. Recently, meta-analysis methods enable scientists to simultaneously study different molecules such as transcripts and proteins under various conditions [8].

The stress-inducible genes that their products involved in the stress may be classified into two groups: the first group including those that protect the cells against stress; and the second group including those that have crucial roles in gene expression under stress conditions [9].

Determining common proteins and their encoding genes between

^{*} Corresponding author.

E-mail address: alemzadeh@shirazu.ac.ir (A. Alemzadeh).

different stresses will allow us to shed some light on the SRMs in plants. Recent years have witnessed an upsurge in the number of functional molecular studies reporting crosstalk in different pathways involved in stress response mechanisms. There are some meta-analysis data from different plants revealing important genes which have hub roles in different abiotic stresses [6, 10]. Among the plants, *Arabidopsis* as a model plant has been subjected to a number of meta-analysis studies which leads to finding differentially expressed genes (DEGs) between stress and non-stress conditions [8, 11, 12, 13]. Detection of DEGs is the first step in understanding of SRMs in plants [13]. Recent advances in biotechnology and development of new technologies caused to facilitate the detection of new genes and determine their functions. Genomics, transcriptomics, proteomics and metabolomics may be used to identify important genes related to abiotic stress response and determine their roles in different pathways involved in SRMs [14]. One of the most important molecules which play crucial roles in these mechanisms are transcription factors [6]. There are some reports showed the roles of some transcription factors family in regulation of stress-inducible genes under stress conditions [6, 9, 15].

Gene ontology (GO) enrichment has been used as a powerful tool in many studies to determine the molecular activity of a gene (molecular function), determine the physiological role of a gene product and coordination with other genes (biological process) and determine the location of gene product in the cell (cellular component). Also, GO is usually used to annotate the function of genes and their proteins from various species especially model organisms such as *Arabidopsis*.

The aim of this study was to identify genes involved in drought and salt stress responses in *Arabidopsis* using microarray gene expression data by meta-analysis approach. In addition, we aimed to detect DEGs between stress and non-stress conditions. Moreover, in this research it was tried to figure out most important transcription factors involved in these stresses that affect the expression of various genes through their down- or up-regulation. In this paper, the functions of identified DEGs and transcription factors were discussed and using network and GO analysis, the complexity of SRMs in *Arabidopsis* was explained.

2. Materials and methods

2.1. Gene expression data collection

The raw microarray expression data was downloaded from ArrayExpress database (<https://ebi.ac.uk/arrayexpress/>) to identify drought and salt stress-responsive genes in *Arabidopsis*. The data originated from Affymetrix microarray platform. These datasets obtained from 6 independent experiments with 38 samples from leaves and roots of plants exposed to drought and salt stresses. These data were obtained from series IDs E-MEXP-1863, E-GEOD-48474 and E-GEOD-40061 (response to drought stress), and E-MEXP-2858, E-GEOD-53308 and E-GEOD-71001 (response to salt stress). All experiments were performed under 22–25 °C, 65–90% RH (relative humidity) and 16/8 h light/dark cycle in the growth chamber. Wild-type Columbia-0 was used in all experiments. In the E-MEXP-1863 experiment, 3-week-old plants were subjected to drought stress by withholding water for 5 days until plant samples reached of $52.5 \pm 7.5\%$ relative water content (RWC). In the E-GEOD-48474 experiment, drought stress was applied to 2-week-old WT plants were by withholding water for 21 days. Leaf sample were collected for transcriptome analysis in mentioned experiments. In the E-GEOD-40061 experiment, root sample were collected from 3-week-old plants which were exposure under drought stress using withholding water for 14 days (soil moisture below 30%). In the E-MEXP-2858 experiment, leaf sample were collected from one-month plants which treated with 200mM NaCl for seven days for salt treatment. In the E-GEOD-53308 experiment, leaf sample were collected from one-month plants which treated with 150 mM NaCl for four days for salt treatment. In the E-GEOD-71001 experiment, root sample were collected from 6-week-old plants which treated with 150 mM NaCl for 6 h.

2.2. Detection of DEGs

The data were retrieved from ArrayExpress database and then analyzed to obtain DEGs involved in responses to drought and salt stresses. The downloaded raw data in format CEL data were analyzed by FlexArray software version 1.6.3. Data were normalized by Robust Multi-array Average (RMA) algorithm using R version 3.2.2. To identify DEGs, the processed microarray data were subjected to two-sample student's t-test. Up-regulated genes with significant *P*-value (false discovery rate (FDR) < 0.05) with fold change value greater than 1.5 selected as DEGs [6, 11]. Venn diagram of DEGs was created by Venny 2.0 (<http://bioinfo.gp.cnb.csic.es/tools/venny/>).

2.3. Detection of transcription factors

The selected DEGs were further compared with the transcription factors genes in the PlantTFDB server <http://planttfdb.cbi.pku.edu.cn/index.php/> for identification of genes encoding transcription factors.

2.4. Gene ontology of DEGs

GO enrichment analysis of DEGs was performed using the AgriGO tool (<http://bioinfo.cau.edu.cn/agriGO/>) with default parameters such as Fisher statistical test method, Multi-test adjustment method of Yekutieli (FDR under dependency) with significant level of 0.05 and complete GO gene ontology type [16]. A Singular Enrichment Analysis (SEA) was performed using TAIR genome as reference and FDR less than <0.05.

2.5. Interaction networks between DEGs

Functional associations between DEGs during drought and salt stresses were figured out by STRING v.10 (Search Tool for the Retrieval of Interacting Genes/Proteins) database (<https://string-db.org/>). These associations were constructed based on co-expression data which is stored in the NCBI-GEO database, co-occurrence of the genes in the same organisms, text mining which is a list of significant protein interaction groups, extracted from the abstracts of scientific literature, significant protein interaction datasets from other experiments, individual gene fusion events per species, significant protein interaction groups gathered from curated databases and neighborhood which is runs of genes that occur repeatedly in close neighborhood in genomes. Confidence score for interactions was medium score (above 0.4). The gene network obtained from STRING v.10 was imported into Cytoscape version 3.6.0 for further analysis and display.

3. Results and discussion

3.1. Identification of DEGs in *Arabidopsis thaliana* under stress conditions

This study was carried out to achieve more information about the changes in gene expression in response to drought and salt stresses in leaves and roots of *Arabidopsis*. Through comprehensive comparison analysis, we identified a set of 4540 up-regulated DEGs in the leaf tissue which 1643 (36%) and 2897 (64%) of them up-regulated in response to drought and salt stress, respectively, whereas in the root tissue 6906 DEGs were up-regulated that 3150 (46%) and 3756 (54%) of them up-regulated in response to drought and salt stress, respectively (supplementary 1). The number of DEGs which up-regulated in roots was 1.5-fold higher than leaves that under drought stress was 1.9-fold, whereas under salt stress was less than 1.3-fold. A four-way Venn diagram was used to represent co-occurrence of DEGs under salt and drought stresses in root and leaf tissues. The diagram represented a list of 584 (13%) DEGs was up-regulated under salt and drought stresses in leaf tissue, but 1186 (17%) DEGs were common between drought and salt stresses in roots (Fig. 1). In the other words, the number of DEGs involved in both stresses in roots was 2-fold higher than leaves. Also, the common DEGs between

roots and leaves under drought and salt stresses were determined using the Venn diagram. Under salt stress, 1266 (11%) DEGs were common between root and leaf tissues, but the number of common DEGs under drought stress was only 302 (2.6%), that means the number of common DEGs in both tissues under salt stress was 4.2-fold higher than drought stress (Fig. 1). It can be concluded that the number of common pathways between roots and leaves induced by salt stress is more than those of induced by drought stress.

As mentioned above, the number of DEGs was up-regulated under stress conditions in roots was higher than leaves that point to the important role of roots in abiotic SRMs in *Arabidopsis*. In some plants like cotton it has been reported that the number of DEGs in leaves was higher than roots under salt stress [17], but there are some reports in other plants which showed the number of up-regulated DEGs in roots were higher than leaves under drought stress [18]. It indicates that different species may use different tolerance mechanisms in response to different stresses. The root is the first tissue to sense stress conditions and it may be explained why, in some plants like *Arabidopsis*, the number of genes involved in the stress in root tissue is higher than other tissues, but in some plants like cotton, probably, leaves have a central role in the control of water loss and hence, more genes involved in the stress in this tissue [19]. According to their stress response mechanisms, plants may be divided into classes: (i) plants that root has a central role and (ii) plants that the leaf has a central role.

The expression changes of DEGs was from 1.5 to 137-fold respect to the control (supplementary 1). The percent of DEGs which up-regulated more than 10-fold in response to drought stress in the root tissue (3.1%) was more than the leaf tissue (1.6%), but it was reverse in the case of salt stress, in roots, 1.6% of DEGs up-regulated more than 10-fold, while it was 2% for leaves (supplementary 1). These results indicate that the genes in leaves and roots have different expression patterns in response to salt and drought stresses.

3.2. Gene ontology

Gene ontology (GO) enrichment analysis was conducted to detect the function of up-regulated DEGs in response to drought and salt stresses in root and leaf tissues in *Arabidopsis*. A total of 1617 and 2854 DEGs

respectively in drought and salt stresses in leaves and 3026 and 3628 DEGs respectively in drought and salt stresses in roots were analyzed for GO analysis. Based on GO analysis, up-regulated DEGs were categorized into 3 groups of biological process (BP), molecular function (MF) and cellular component (CC).

The aforementioned groups, BP, MF and CC consisted of 17, 11 and 12 categories, respectively (Figs. 2 and 3). In both tissues, in response to both stresses, for CC most enriched GO terms were cell (about 87%), cell part (about 87%) and organelle (about 65%), whereas the two highly most enriched ones for MF were binding (49%) and catalytic activity (42%) (Figs. 2 and 3). Furthermore, for BP the highest frequency of DEGs function was belonged to cellular process (53%), metabolic process (50%) and response to stimulus (36%), respectively (Figs. 2 and 3), which agreement with results of other researches about microarray meta-analysis in plants in response to biotic and abiotic stresses [6, 20].

In BP, the significant DEGs in cellular process (GO:0009987) were assigned to cellular response to stimulus, regulation of cellular process, multi-organism cellular process, cellular metabolic process, single-organism cellular process and positive regulation of cellular process (supplementary 2). For the second group in BP, metabolic process (GO:0008152), the significant ones were assigned to GO terms such as single-organism metabolic process, cellular metabolic process, organic substance metabolic process and catabolic process.

In BP group, for signaling (GO:0023052) and response to stimulus (GO:0050896), the number of DEGs in each tissue in salt stress was higher than drought stress. In signaling group, 9.5 and 9.7 percent of DEGs in the leaf and 9.1 and 10.3 percent of DEGs in the root were up-regulated in response to drought and salt stresses, respectively (Figs. 2 and 3). Gene ontology of up-regulated DEGs showed that signaling category included DEGs involvement in hormone-activated signaling pathways, signal transduction, MAP kinase kinase activity, ATPase activity, calcium-dependent protein serine/threonine kinase activity and receptor signaling protein serine/threonine kinase activity. Phytohormones such as abscisic acid, auxin, cytokinin and gibberellic acid play important roles in different process of growth and development and also response to biotic and abiotic stresses in plants [21, 22]. In Chai et al (2019) study, Expression of *AtNCED3* (nine-cis-epoxycarotenoid dioxygenase) which is an important gene in ABA biosynthesis, increased about




Fig. 1. Venn diagram of differentially expressed genes in response to drought and salt stresses in leaves and roots of *Arabidopsis thaliana* which created by Venny 2.0.




Fig. 2. Gene ontology analysis in leaves. Frequency of most representative biological process terms in *Arabidopsis thaliana* under salt and drought stresses. Gene ontology analysis was made in the AgriGO platform (FDR = 5%). More details in Supplementary 2.

4.6-fold in *Arabidopsis* plants after salt stress. Also, in our study salt stress increased expression of this gene about 1.8-fold in leaf tissue of *Arabidopsis* plant (supplementary 1).

Another major part of up-regulated DEGs in BP group, was belonged to response to stimulus category. Based on GO results, 37 and 38 percent of DEGs in leaves and 31 and 36 percent of DEGs in roots were up-regulated in response to drought and salt stresses, respectively (Figs. 2 and 3). Components of this group were up-regulated in response to abiotic stimulus, such as temperature, salt and drought stresses, response to endogenous stimulus such as phytohormones.

Components of MF group have important roles in response to abiotic stresses. In this category, catalytic activity and binding groups have more

up-regulated DEGs in both tissues. Majority of up-regulated DEGs (>46%) was belonged to binding category (GO:0005488) and more than 40% was belonged to catalytic activity category (GO:0003824) in each tissue (Figs. 2 and 3). The results were coincidence with that obtained by Gao *et al* (2008) in response to drought stress in chickpea [23].

Some significant GO terms of binding group were ion binding, lipid binding, cofactor binding and protein binding and significant GO terms of catalytic activity group were oxidoreductase activity and hydrolase activity. Catalytic activity role in response to abiotic stresses especially osmotic stress was demonstrated [24]. Based on previous studies, anti-oxidant activity plays an important role in abiotic and even biotic stresses tolerance, but despite its important role in plant defense against osmotic




Fig. 3. Gene ontology analysis in roots. Frequency of most representative biological process terms in *Arabidopsis thaliana* under salt and drought stresses. Gene ontology analysis made in the AgriGO platform (FDR = 5%). More details in Supplementary 2.

stress, amount of up-regulated DEGs in this group was low in each tissue and in each stress, so that amount of this group was 1.4 and 0.94 percent in leaf and 0.85 and 1.1 percent in root respectively under drought and salt stresses. This kind of results has also been reported in earlier studies [25]. Genes encoded antioxidant activity are one of the most important genes in response to abiotic stresses. In plants during of abiotic and biotic stresses, reactive oxygen species (ROS) generated which severe damage to whole cells and organisms. Antioxidant enzymes such as superoxide dismutase (SOD), catalase (CAT), ascorbate peroxidase (APX), glutathione peroxidase (GPX), and glutathione reductase (GR) scavenge ROS components in plants [25, 26]. Kanesaki *et al* (2002) identified which salt stress caused to increase the expression of superoxide dismutase in *Synecchocystis*. In our study, this gene up-regulated in response to drought and salt stress in leaf tissue [27]. Furthermore, Ghaffari *et al* (2013) reported an increase of ascorbate peroxidase 2-like protein and Cu/Zn superoxide dismutase in a tolerant line of sunflower roots in response to drought stress [28]. In our study, the expression of ascorbate peroxidase gene increased in root tissue in response to drought and salt stress. Based on results of Li *et al* (2012), osmotic stress using polyethylene glycol (PEG), increased the expression of monodehydroascorbate reductase (MDHAR) in leaves of wild type tomato. MDHAR play important role in plant antioxidant system by regeneration of ascorbate. In our study, the expression of this gene also up-regulated in leaf tissue in response to salt stress.

Transcription factors play important roles in plant responses to stresses via regulation of downstream genes by binding to *cis*-acting elements in the promoter region of target genes [29]. The percent of these proteins were low in our study as previously reported by Khojasteh *et al* (2018).

The results of GO analysis showed a similarity pattern of DEGs function for leaf and root tissues in response to drought and salt stresses but differences were in specific DEGs and number of significant DEGs in each group between two stresses in each tissue (supplementary 2). In roots, approximately in most categories of BP and MF groups, amount of up-regulated DEGs under salt stress were higher than drought stress, but in the CC group, the amount of up-regulated DEGs was approximately equal under both stresses except for membrane, membrane part and organelle part categories which amount of up-regulated DEGs was higher in drought stress. Whereas in leaf tissue, the amount of up-regulated DEGs in different categories of gene ontology was not so much different between two stresses. These results may indicate that roots respect to salt stress is more sensitive than drought stress.

3.3. Identification of DEGs encoding transcription factors in *Arabidopsis* in response to drought and salt stresses

Transcription factors (TFs) play important roles in plant tolerance to environmental stresses. These proteins as *trans*-acting elements bind to *cis* elements in promoter region of various genes involved in different stresses and as a result cause to activate or repress the expression of target genes and subsequently increase the plants tolerance [30]. With regard to the important role of transcription factors in plant response to different stresses, we further identified and compared the expression profiles of these proteins in root and leaf tissues of *Arabidopsis* under drought and salt stresses.

Based on the results, 109 (about 6.6%) and 223 (about 7.7%) genes in leaves and 234 (about 7.4%) and 396 (about 10.5%) genes in roots respectively in response to drought and salt stress were identified as TFs (Fig. 4). Identified TFs were classified in 28 and 34 groups in leaves, and 37 and 40 groups in roots in response to drought and salt stress, respectively (supplementary 3). In each tissue under both stresses, the greatest TF families were belonged to AP2/ERF, MYB and NAC families. Moreover, other important TF families such as bZIP, HSF, C2H2 and WRKY were also detected. In addition, 54 and 98 TFs respectively in leaf and root tissues were identified as common TFs between drought and salt stresses (supplementary 4). Common TFs in leaves were divided into 19

groups. The greatest number of these TFs was belonged to NAC family followed by AP2/ERF, HSF, MYB, C2H2 and bZIP accounting for 68% of the total TFs (Fig. 5). Furthermore, common TFs in roots were divided into 23 groups which the greatest number of them was belonged to MYB family followed by AP2/ERF, NAC, DOF, bZIP and HSF accounting for 64% of the total TFs (Fig. 6).

AP2/ERF family is one of the most important transcription factors family in plants which plays important roles in response to biotic and abiotic stresses [31, 32, 33]. In the present study, CRF7 (AT1G22985, CYTOKININ RESPONSE FACTOR 7), a member of ERF family, was up-regulated in response to drought and salt stresses in leaves whereas its expression in roots increased only under salt stress (supplementary 3). The analysis of *Arabidopsis* microarray data previously showed that the expression of CRF genes changed in response to abiotic stresses [34]. Overexpression of *HARDY* (AT2g36450) gene, encoding a member of this family, improved drought and salt tolerance in transgenic *Trifolium alexandrinum* L [35]. Our microarray analysis results also showed that the expression of this gene up-regulated in roots in response to drought stress (supplementary 3). RAP2.1 (AT1G46768) which up-regulated in response to both stresses in leaves (supplementary 3), encodes a member of the DREB subfamily. It has previously shown that drought and cold stresses induced expression of RAP2.1 gene through ABA-independent pathway [36]. The expression of RAP2.1 gene is negatively regulated by cold and drought stresses in *Arabidopsis*. Moreover, RAP2.6 (AT1G43160) and DREB19 (AT2G38340) which encodes a member of ERF and DREB subfamilies, respectively, up-regulated in roots in response to drought and salt stresses, whereas up-regulated only in response to salt stress in leaves (supplementary 3). In addition, it has been reported that the expression of these genes significantly up-regulated under salt stress in transgenic plants [36]. It has been previously reported that the expression of DREB2A (AT5G05410), a gene which encodes a member of DREB subfamily, increased in response to salt, drought and heat stresses [32, 37]. One of the earliest responses to drought stress is activation of ERF genes such as ERF8 (AT1G53170) which results in the induction of ethylene biosynthesis [38]. The expression of both of these genes, DREB2A and ERF8, was up-regulated in response to salt and drought stress in leaves, whereas up-regulated only under salt stress in roots (supplementary 3). *Arabidopsis* transgenic plants contain *TaERF1* showed tolerance to different stresses such as drought, salt and low-temperature stresses [39]. It has been previously shown that ERF104 (AT5G61600) and ERF105 (AT5G51190) which in this study up-regulated in leaves under both stresses (supplementary 3), play a critical role in response to high light and freezing stresses [40].

Another important transcription factor family in plants is NAC family. There are some reports that showed the members of this family has key roles in response to biotic and abiotic stresses [41]. In *Arabidopsis*, 105 redundant putative NAC genes were identified [42]. Our results showed that NAC019 (AT1G52890), NAC02 (AT5G04410) and NAC072 (AT4G27410, RD26) were up-regulated in leaves in response to both stresses, while their expression increased in roots only under salt stress, whereas NAC01 (AT1G01010) and NAC032 (AT1G77450) were up-regulated in roots in response to both stresses and NAC032 was up-regulated only in response to salt stress in leaves. NAC047 (AT3G04070) was up-regulated in both tissues in response to both stresses (supplementary 3). It has been shown that the introduction of NAC019 in *Arabidopsis* plants increased the tolerance to drought, salt, and low-temperature stresses in transgenic plants [43]. Different studies showed that other members of this family such as OsNAC1, OsNAC2, OsNAC10 and AtNAC2 have important roles in various abiotic stresses [44, 45]. Furthermore, it has been also demonstrated the role of some members of this family in ABA-dependent stress-signaling pathway [44, 46].

bZIP family is another large family of transcription factor in plants and 75 members of this family was recognized in *Arabidopsis* [47]. Microarray analysis of *Arabidopsis* in response to drought stress showed that the expression of bZIP44 (AT1G75390) increased significantly in




Fig. 4. Venn diagram of differentially expressed genes encoding transcription factors in response to drought and salt stresses in leaves and roots of *Arabidopsis thaliana* which created by Venny 2.0.

shoot tissue [48]. Our results indicated that its expression increased in response to both stresses in roots, whereas in leaves, up-regulated only in response to drought stress (supplementary 3). Moreover, in this study, other members of bZIP family such as *AtbZIP50/TGA7* (AT1G77920), *AtbZIP55/GBF3* (AT2G46270) were detected in response to drought and salt stresses in leaves, whereas *AtbZIP56/HY5* (AT5G11260) up-regulated in response to both stresses only in roots and in leaves up-regulated only under salt stress (supplementary 3).

MYB and WRKY are two families of transcription factors involved in response to abiotic stresses in plants [48]. For example, it has been shown that the expression of *AtMYB108/BOS1* gene induced in response to oxidative stress, drought and salt stresses. *AtMYB108* also involved in the crosstalk between abiotic and biotic stresses [49]. *AtMYB65* (AT3G11440) is a member of this family which involves in GA signaling in growth and flowering processes [50] and here, up-regulated in response to both stresses in roots (supplementary 3). *AtMYB108* (AT3G06490) regulated filament elongation and anther dehiscence by JA and GA signaling pathways. *AtMYB108* is involved in the crosstalk

between abiotic and biotic stress signaling. Furthermore, the expression of *AtMYB108/BOS1* was induced in response to oxidative stress, drought and salt stresses [49]. Our work also showed that the expression of this gene increased in roots in response to both stresses whereas, in leaves up-regulated only in response to salt stress (supplementary 3). The results of the study showed that the expression of AT5G60890 (*AtMYB34*) up-regulated in root tissue in response to both stresses, whereas in leaves, its expression increased only in response to salt stress. Up-regulation of this gene was identified in *Arabidopsis* plant in Bhargava *et al* (2013) study by microarray and RNA-Seq analysis.

It has been shown that there are 74 members for WRKY family in *Arabidopsis* [51]. *AtWRKY25* (AT2G30250), a member of WRKY family, expresses under different abiotic stresses such as heat, drought, salt and oxidative stresses [52, 53, 54]. Here also, its expression increased in response to both stresses only in leaves but in roots, its expression increased just in response to salt stress (supplementary 3). In other species, it has also reported that the members of these families have important roles in abiotic stress responses. *OsWRKY23* (01g53260), as




Fig. 5. Common transcription factors in leaves of *Arabidopsis thaliana* which up-regulated in response to drought and salt stresses.




Fig. 6. Common transcription factors in roots of *Arabidopsis thaliana* which up-regulated in response to drought and salt stresses.

member of these family in rice, involved in the regulation of resistance to salt, ABA, H_2O_2 , osmotic and dark stresses [55]. They showed that the overexpression of *OsWRKY23* in *Arabidopsis* caused to accelerate the leaf senescence in darkness.

Another important transcription factor family is HSF family. There are 21 genes which encodes the members of this family in *Arabidopsis* [56]. Our results indicated that some members of HSF family, *HSFA2*, *HSFB2B*, *HSFA4A*, *HSFA6A* and *HSFB2A* in leaves and *HSFA2*, *HSFB3*, *HSFA6B* and *HSFA3* in roots were up-regulated in response to both of drought and salt stresses (supplementary 3). These results are in accord with what others reported. Overexpression of *AtHSFA2* in *Arabidopsis*, increased the tolerance of plants against abiotic stresses [57]. These transcription factors are regulated by other proteins, for example, the transcription of *AtHSFA3* is regulated by *DREB2A* gene via binding to two DRE core elements of *DREB2A* in the *AtHSFA3* promoter region [58]. These TFs have important roles in abiotic stresses in other species, it has been shown that the expression of tomato *HSFA3* and wheat *HSF3* increased thermotolerance of *Arabidopsis* transgenic plants [59, 60]. The result of this study showed that the transcription factors family of AP2/ERF, MYB, NAC, bZIP, HSF and WRKY may play important roles in response to drought and salt stresses in *Arabidopsis*.

3.4. Gene network analysis

In this study, a network was generated between common TFs in leaves and roots in response to drought and salt stresses in *Arabidopsis*, using STRING 10.5. Only 113 of 152 TFs had interactions (supplementary 5) and so all the disconnected nodes in the network were removed. Between the TFs, seven of them, *HSFA2*, *HSF4*, *MYB47*, *MYB32*, *NAC047*, *NF-YA5* and *AT1G76580* were common in both tissues and both stresses. Subsequently, the TFs which had interactions together, were subjected to Cytoscape v2.8.2 for more analysis. Based on the results of Cytoscape, a total of 550 interactions (edges) were detected between 113 TFs (nodes) (Fig. 7, supplementary 6). In this network, hub genes were identified based on their outdegree (the number of genes which was affected by the same gene) and those that had an outdegree equal or more than 10 considered as a hub gene which includes *HSFB2A* (belongs to HSF family), *DREB2A* (belongs to AP2/ERF family), *ZAT6* (belongs to C2H2 family), *RHL41* (belongs to C2H2 family), *HSFB2B* (belongs to HSF family) and *HSFA4A* (belongs to HSF family) (supplementary 6). These results indicated that the members of HSF, AP2/ERF and C2H2 families appear to play critical roles in salt and drought stress response mechanisms.

HSFB2A had the highest outdegree in the network, so that 16 genes, *RHL41*, *ZAT6*, *DREB2A*, *HSFA2*, *HSFA6B*, *HSFA6A*, *HSFA4A*, *HSFB2B*, *HSFB3*, *HSF4*, *NAC046*, *NAC032*, *NAC087*, *ZF2*, *CZF1*, *ZF2* were directly affected by this TF (Fig. 7, supplementary 6). This is the first report that showed this protein is a hub TF in response to salt and drought stresses.

DREB2A was affected by *RHL41*, *HSFB2A*, *NAC06* and *NAC096* whereas 15 genes such as *NAC019*, *RD26*, *HSFA4A*, *HSFA3*, *HSFA2*, *AT-HSFB2B* and etc. were affected by this TF (Fig. 7, supplementary 6). DREB (dehydration responsive element binding) is a subfamily of AP2/ERF transcription factors family that regulate ABA-independent signal transduction pathway [61]. *DREB2A* has one AP2/ERF domain and includes DRE-binding proteins (DREBs) which binds to dehydration-responsive element/C-repeat (DRE/CRT) in promoter of abiotic stress-responsive genes and regulates their expression [32, 37]. It has been previously shown that the expression of *DREB2A* was induced in response to dehydration, salt and heat stresses (5, 32, 37). Based on the gene network analysis, *HSFB2A*, *RHL41*, *DREB2A*, *ERF104* and *HY5* genes affected *ZAT6* expression (Fig. 7, supplementary 6). This TF activated the expression of some genes encoding TFs such as *RD26*, *HSFA4A*, *ERF-1*, *WRKY25* and *NAC047* (Fig. 7). *ZAT6* belongs to C2H2 subfamily of Zinc-finger transcription factors family [62]. Based on the results of Liu et al (2013), salt and osmotic stresses regulated *ZAT6* expression in *Arabidopsis*. The expression of *RHL41* directly activated by *ERF104* and *HSFB2A* genes, and affected 14 genes such as *DREB2A*, *HSFA2*, *HSFA4A*, *WRKY25*, *ZAT6* *ERF-1* and etc (Fig. 7, supplementary 6).

RHL41 (AT5G59820) which also known as *ZAT12* is a member of C2H2 zinc finger transcription factors family [63]. The expression of *ZAT12* induced by light, low and high temperature, wounding, osmotic, salt and oxidative stresses [64, 65]. It has been also proved *ZAT12* involved in the up-regulation of *WRKY25* in response to oxidative stress, which is one of the genes in ROS signal transduction [66]. Based on our knowledge, there are no reports to indicate that, *AT-HSFB2B* and *HSFA4A* play important roles in salt and drought stresses, and this is the first report to show these members of HSF family may play as hub proteins in these stresses.

Network analysis interestingly showed that *STZ* (salt tolerance zinc finger), a TF from C2H2 family, is affected by 17 other TFs which is the highest number in this network, but has no effect on other TFs (outdegree equal zero) (Fig. 7, supplementary 6). It has been previously shown that this protein acts as a transcription repressor to elevate abiotic stress tolerance in plants [67]. Hence, it can be resulted that *STZ* has a critical role in stress tolerance mechanism in *Arabidopsis* and can be considered as a suitable candidate to enhance salt and drought tolerance in crop




Fig. 7. Network analysis of differentially expressed genes encoding transcription factors detected in salt and drought stresses. Network produced by STRING (version 10) with a confidence score >0.4 , and visualized by Cytoscape 3.6.0. Node colors refer to the family of transcription factor. Node size corresponds to the number of outdegrees.

plants through genetic engineering approach.

4. Conclusion

Drought and salt stresses are two main groups of abiotic stresses which affect agriculture productivity in worldwide. Furthermore, there is a crosstalk between abiotic and biotic stresses in plants. Therefore, identification of molecular mechanisms in response to them is helpful to increase simultaneously tolerance to drought and salt stresses. To achieve deeper understanding of molecular mechanisms in *Arabidopsis* and identification of responsive genes to drought and salt stresses, we identified up-regulated DEGs in leaf and root tissues in response to these stresses using microarray data analysis approach. We found that there are some differences between roots and leaves in terms of DEGs. Generally, the number of DEGs which up-regulated in response to stresses in roots was 1.5-fold higher than leaves, especially for drought stress, whereas it can be concluded that the number of common pathways between roots and leaves induced by salt stress is more than those of induced by drought stress. Various proteins involved in stress response mechanisms that transcription factors are most important of them; among these kind of proteins, three families such as HSF, AP2/ERF and C2H2, appear to play critical roles in salt and drought stress response mechanisms in *Arabidopsis*. In addition, the results of this study introducing some new proteins such as *STZ*, a member of AP2/ERF family, which plays a critical role in stress tolerance mechanisms in *Arabidopsis* and can be considered as a suitable candidate for increasing salt and drought tolerance in crop plants.

Declarations

Author contribution statement

Razieh Ghorbani: Performed the experiments; Analyzed and interpreted the data; Contributed reagents, materials, analysis tools or data; Wrote the paper.

Abbas Alemzadeh: Conceived and designed the experiments; Analyzed and interpreted the data.

Hooman Razi: Contributed reagents, materials, analysis tools or data.

Funding statement

This work was supported by Shiraz University.

Competing interest statement

The authors declare no conflict of interest.

Additional information

Supplementary content related to this article has been published online at <https://doi.org/10.1016/j.heliyon.2019.e02614>.

References

- [1] L. Rastgoo, A. Alemzadeh, Biochemical responses of gouan (*Aeluropus littoralis*) to heavy metals stress, *Aust. J. Crop. Sci.* 5 (2011) 375–383.

- [2] D. Golldack, I. Lüking, O. Yang, Plant tolerance to drought and salt: stress regulating transcription factors and their functional significance in the cellular transcriptional network, *Plant Cell Rep.* 30 (2011) 1383–1391.
- [3] L. Rastgo, A. Alemzadeh, A. Afsharifar, Isolation of two novel isoforms encoding zinc- and copper-transporting P1B-ATPase from Gouan (*Aeluropus littoralis*), *Plant Omics* 4 (2011) 377–383.
- [4] Z. Sami, A. Alemzadeh, Isolation and molecular characterization of a novel Na^+/H^+ antiporter gene, *AlNHX2*, from *Aeluropus littoralis* and comparison of *AlNHX1* and *AlNHX2*, *Plant Omics* 9 (2016) 205–212.
- [5] Z. Zinati, A. Alemzadeh, A.H. Kayvanjoo, Computational approaches for classification and prediction of P-type ATPase substrate specificity in *Arabidopsis*, *Physiol. Mol. Biol. Plants* 22 (2016) 163–174.
- [6] E. Ashrafi-Dehkordi, A. Alemzadeh, N. Tanaka, H. Razi, Meta-analysis of transcriptomic responses to biotic and abiotic stress in tomato, *Peer. J.* 6 (2018).
- [7] G.C. Tseng, D. Ghosh, E. Feingold, Comprehensive literature review and statistical considerations for microarray meta-analysis, *Nucleic Acids Res.* 40 (2012) 3785–3799.
- [8] J.S. Rest, O. Wilkins, W. Yuan, M.D. Purugganan, J. Gurevitch, Meta-analysis and meta regression of transcriptomic responses to water stress in *Arabidopsis*, *Plant J.* 85 (2016) 548–560.
- [9] M. Seki, M. Narusaka, J. Ishida, T. Nanjo, M. Fujita, Y. Oono, A. Kamiya, M. Nakajima, A. Enju, T. Sakurai, M. Satou, K. Akiyama, T. Tajii, K. Yamaguchi-Shinozaki, P. Carninci, J. Kawai, Y. Hayashizaki, K. Shinozaki, Monitoring the expression profiles of 7000 *Arabidopsis* genes under drought, cold and high-salt stresses using a full-length cDNA microarray, *Plant J.* 31 (2002) 279–292.
- [10] B. Balan, F.P. Marra, T. Caruso, F. Martinelli, Transcriptomic responses to biotic stresses in *Malus x domestica*: a meta-analysis study, *Sci. Rep.* 8 (2018).
- [11] A. Bhargava, I. Clabaugh, J.P. To, B.B. Maxwell, Y.H. Chiang, G.E. Schaller, A. Loraine, J.J. Kieber, Identification of cytokinin-responsive genes using microarray meta-analysis and RNA-seq in *Arabidopsis*, *Plant Physiol.* 162 (2013) 272–294.
- [12] R. Shaik, W. Ramakrishna, Genes and co-expression modules common to drought and bacterial stress responses in *Arabidopsis* and rice, *PLoS One* 8 (2013).
- [13] P.C. Shen, A.L. Hour, L.Y.D. Liu, Microarray meta-analysis to explore abiotic stress-specific gene expression patterns in *Arabidopsis*, *Bot. Stud.* 58 (2017).
- [14] T.S. Gechev, J. Hille, Molecular basis of plant stress, *Cell. Mol. Life Sci.* 69 (2012) 3161–3163.
- [15] Q. Liu, M. Kasuga, Y. Sakuma, H. Abe, S. Miura, K. Yamaguchi-Shinozaki, K. Shinozaki, Two transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought- and low temperature-responsive gene expression, respectively, in *Arabidopsis*, *Plant. Cell.* 10 (1998) 1391–1406.
- [16] Z. Du, X. Zhou, Y. Ling, Z. Zhang, Z. Su, AgriGO: a GO analysis toolkit for the agricultural community, *Nucleic Acids Res.* 38 (2010).
- [17] F. Zhang, G. Zhu, L. Du, X. Shang, C. Cheng, B. Yang, Y. Hu, C. Cai, W. Guo, Genetic regulation of salt stress tolerance revealed by RNA-Seq in cotton diploid wild species, *Gossypium davidsonii*, *Sci. Rep.* 6 (2016).
- [18] C. Zhang, L. Zhang, S. Zhang, S. Zhu, P. Wu, Y. Chen, M. Li, H. Jiang, G. Wu, Global Analysis of gene expression profiles in physic nut (*Jatropha curcas* L.) seedlings exposed to drought stress, *BMC Plant Biol.* 15 (2015).
- [19] C.S. Galvan-Ampudia, C. Testerink, Salt stress signals shape the plant root, *Curr. Opin. Plant Biol.* 14 (2011) 296–302.
- [20] M. Khojasteh, B. Khahani, M. Taghavi, E. Tavakol, Identification and characterization of responsive genes in rice during compatible interactions with pathogenic pathogens of *Xanthomonas oryzae*, *Eur. J. Plant Pathol.* 151 (2018) 141–153.
- [21] Z. Peleg, E. Blumwald, Hormone balance and abiotic stress tolerance in crop plants, *Curr. Opin. Plant Biol.* 14 (2011) 290–295.
- [22] W.W. Chai, W.Y. Wang, Q. Ma, H.J. Yin, S. Hepworth, S.M. Wang, Comparative transcriptome analysis reveals unique genetic adaptations conferring salt tolerance in a *xerohalophyte*, *Funct. Plant Biol.* 46 (2019) 670–683.
- [23] W.R. Gao, X.S. Wang, Q.Y. Liu, H. Peng, C. Chen, J.G. Li, H. Ma, Comparative analysis of ESTs in response to drought stress in chickpea (*C. arietinum* L.), *BBRC (Biochem. Biophys. Res. Commun.)* 376 (3) (2008) 578–583.
- [24] I.S. Jeong, A. Fukudome, E. Aksoy, W.Y. Bang, S. Kim, Q. Guan, J.D. Bahk, K.A. May, W.K. Russell, J. Zhu, H. Koiwa, Correction: regulation of abiotic stress signaling by *Arabidopsis* C-terminal domain phosphatase-like 1 requires interaction with a K-homology domain-containing protein, *PLoS One* 10 (2015).
- [25] J.B. de Abreu Neto, M. Frei, Microarray meta-analysis focused on the response of genes involved in redox homeostasis to diverse abiotic stresses in rice, *Front. Plant Sci.* 6 (2016) 1260.
- [26] Y. Kanesaki, I. Suzuki, S.I. Allakhverdiev, K. Mikami, N. Murata, Salt stress and hyperosmotic stress regulate the expression of different sets of genes in *Synechocystis* sp. PCC 6803, *Biochem. Biophys. Res. Commun.* 290 (2002) 339–348.
- [27] M. Ghaffari, M. Toorchi, M. Valizadeh, S. Komatsu, Differential response of root proteome to drought stress in drought sensitive and tolerant sunflower inbred lines, *Funct. Plant Biol.* 40 (6) (2013) 609–617.
- [28] F. Li, Q.Y. Wu, M. Duan, X.C. Dong, B. Li, Q.W. Meng, Transgenic tomato plants overexpressing chloroplastic monodehydroascorbate reductase are resistant to salt- and PEG-induced osmotic stress, *Photosynthetica* 50 (1) (2012) 120–128.
- [29] K. Yamaguchi-Shinozaki, K. Shinozaki, Transcriptional regulatory networks in cellular responses and tolerance to dehydration and cold stresses, *Annu. Rev. Plant Biol.* 57 (2006) 781–803.
- [30] W.J. Chen, T. Zhu, Networks of transcription factors with roles in environmental stress response, *Trends Plant Sci.* 9 (2004) 591–596.
- [31] K.H. Sohn, S.C. Lee, H.W. Jung, J.K. Hong, B.K. Hwang, Expression and functional roles of the pepper pathogen-induced transcription factor *RAV1* in bacterial disease resistance, and drought and salt stress tolerance, *Plant Mol. Biol.* 61 (2006) 897.
- [32] Y. Sakuma, K. Maruyama, F. Qin, Y. Osakabe, K. Shinozaki, K. Yamaguchi-Shinozaki, Dual function of an *Arabidopsis* transcription factor DREB2A in water-stress-responsive and heat-stress-responsive gene expression, *Proc. Natl. Acad. Sci. U.S.A.* 103 (2006) 18822–18827.
- [33] A. Karaba, S. Dixit, R. Greco, A. Aharoni, K.R. Trijatmiko, N. Marsch-Martinez, A. Krishnan, K.N. Nataraja, M. Udayakumar, A. Pereira, Improvement of water use efficiency in rice by expression of *HARDY*, an *Arabidopsis* drought and salt tolerance gene, *Proc. Natl. Acad. Sci. U.S.A.* 104 (2007) 15270–15275.
- [34] C.T. Argueso, F.J. Ferreira, J.J. Kieber, Environmental perception avenues: the interaction of cytokinin and environmental response pathways, *Plant Cell Environ.* 32 (2009) 1147–1160.
- [35] G.M. Abogadallah, R.M. Nada, R. Malinowski, P. Quick, Overexpression of *HARDY*, an AP2/ERF gene from *Arabidopsis*, improves drought and salt tolerance by reducing transgenic *Trifolium alexandrinum* L., *J. Appl. Genet.* 233 (2011) 1265–1276.
- [36] S.S. Krishnaswamy, S. Srivastava, M. Mohammadi, M.H. Rahman, M.K. Deyholos, N.N. Kav, Transcriptional profiling of pea ABR17 mediated changes in gene expression in *Arabidopsis thaliana*, *BMC Plant Biol.* 8 (2008) 91.
- [37] Y. Sakuma, K. Maruyama, Y. Osakabe, F. Qin, M. Seki, K. Shinozaki, K. Yamaguchi-Shinozaki, Functional analysis of an *Arabidopsis* transcription factor, DREB2A, involved in drought-responsive gene expression, *Plant. Cell.* 18 (2006) 1292–1309.
- [38] M. Dubois, H. Claeys, L. Van den Broeck, D. Inzé, Time of day determines *Arabidopsis* transcriptome and growth dynamics under mild drought, *Plant Cell Environ.* 40 (2017) 180–189.
- [39] Z.S. Xu, L.Q. Xia, M. Chen, X.G. Cheng, R.Y. Zhang, L.C. Li, Y.X. Zhao, Y. Lu, Z.Y. Ni, L. Liu, Z.G. Qiu, Isolation and molecular characterization of the *Triticum aestivum* L. ethylene-responsive factor 1 (*TaERF1*) that increases multiple stress tolerance, *Plant. Mol. Biol.* 65 (2007) 719–732.
- [40] S. Bolt, E. Zuther, S. Zintl, D.K. Hinch, T. Schmölling, ERF105 is a transcription factor gene of *Arabidopsis thaliana* required for freezing tolerance and cold acclimation, *Plant Cell Environ.* 40 (2017) 108–120.
- [41] K. Nakashima, L.S.P. Tran, D. Van Nguyen, M. Fujita, K. Maruyama, D. Todaka, Y. Ito, N. Hayashi, K. Shinozaki, K. Yamaguchi-Shinozaki, Functional analysis of a NAC-type transcription factor OsNAC6 involved in abiotic and biotic stress-responsive gene expression in rice, *Plant J.* 51 (2007) 617–630.
- [42] H. Ooka, K. Satoh, K. Doi, T. Nagata, Y. Otomo, K. Murakami, K. Matsubara, N. Osato, J. Kawai, P. Carninci, Y. Hayashizaki, K. Suzuki, K. Kojima, Y. Takahara, K. Yamamoto, S. Kikuchi, Comprehensive analysis of NAC family genes in *Oryza sativa* and *Arabidopsis thaliana*, *DNA Res.* 10 (2003) 239–247.
- [43] L.S.P. Tran, K. Nakashima, Y. Sakuma, S.D. Simpson, Y. Fujita, K. Maruyama, M. Fujita, M. Seki, K. Shinozaki, K. Yamaguchi-Shinozaki, Isolation and functional analysis of *Arabidopsis* stress-inducible NAC transcription factors that bind to a drought-responsive cis-element in the early responsive to dehydration stress 1 promoter, *Plant. Cell.* 16 (2004) 2481–2498.
- [44] X.J. He, R.L. Mu, W.H. Cao, Z.G. Zhang, J.S. Zhang, S.Y. Chen, AtNAC2, a transcription factor downstream of ethylene and auxin signaling pathways, is involved in salt stress response and lateral root development, *Plant J.* 44 (2005) 903–916.
- [45] J.S. Jeong, Y.S. Kim, K.H. Baek, H. Jung, S.H. Ha, Y.D. Choi, M. Kim, C. Reuzeau, J.K. Kim, Root-specific expression of *OsNAC10* improves drought tolerance and grain yield in rice under field drought conditions, *Plant Physiol.* 153 (1) (2010) 185–197.
- [46] M. Fujita, Y. Fujita, K. Maruyama, M. Seki, K. Hiratsu, M. Ohme-Takagi, L.S.P. Tran, K. Yamaguchi-Shinozaki, K. Shinozaki, A dehydration-induced NAC protein, RD26, is involved in a novel ABA-dependent stress-signaling pathway, *Plant J.* 39 (2004) 863–876.
- [47] M. Jakoby, B. Weisshaar, W. Dröge-Laser, J. Vicente-Carbajosa, J. Tiedemann, T. Kroj, F. Parcy, bZIP transcription factors in *Arabidopsis*, *Trends Plant Sci.* 7 (2002) 106–111.
- [48] S. Rasheed, K. Bashir, A. Matsui, M. Tanaka, M. Seki, Transcriptomic analysis of soil-grown *Arabidopsis thaliana* roots and shoots in response to a drought stress, *Front. Plant Sci.* 7 (2016) 180.
- [49] T. Mengiste, X. Chen, J. Salmeron, R. Dietrich, The *BOTRYTIS SUSCEPTIBLE1* gene encodes an R2R3MYB transcription factor protein that is required for biotic and abiotic stress responses in *Arabidopsis*, *Plant. Cell.* 15 (2003) 2551–2565.
- [50] G.F.W. Gocal, C.C. Sheldon, F. Gubler, T. Moritz, D.J. Bagnall, C.P. MacMillan, S.F. Li, R.W. Parish, E.S. Dennis, D. Weigel, R.W. King, *GAMYB-like* genes, flowering, and gibberellin signaling in *Arabidopsis*, *Plant Physiol.* 127 (2001) 1682–1693.
- [51] A. Ülker, I.E. Somssich, WRKY transcription factors: from DNA binding towards biological function, *Curr. Opin. Plant Biol.* 7 (2004) 491–498.
- [52] S. Li, Q. Fu, W. Huang, D. Yu, Functional analysis of an *Arabidopsis* transcription factor *WRKY25* in heat stress, *Plant Cell Rep.* 28 (2009) 683–693.
- [53] S. Li, Q. Fu, L. Chen, W. Huang, D. Yu, *Arabidopsis thaliana* *WRKY25*, *WRKY26*, and *WRKY33* coordinate induction of plant thermotolerance, *Planta* 233 (2011) 1237–1252.
- [54] Y. Jiang, M.K. Deyholos, Functional characterization of *Arabidopsis* NaCl-inducible *WRKY25* and *WRKY33* transcription factors in abiotic stresses, *Plant Mol. Biol.* 69 (2009) 91–105.
- [55] S. Jing, X. Zhou, Y. Song, D. Yu, Heterologous expression of *OsWRKY23* gene enhances pathogen defense and dark-induced leaf senescence in *Arabidopsis*, *Plant Growth Regul.* 58 (2009) 181–190.

- [56] K.D. Scharf, T. Berberich, I. Ebersberger, L. Nover, The plant heat stress transcription factor (Hsf) family: structure, function and evolution, *Biochim. Biophys. Acta. Gene. Regul. Mech.* 1819 (2012) 104–119.
- [57] A. Nishizawa, Y. Yabuta, E. Yoshida, T. Maruta, K. Yoshimura, S. Shigeoka, Arabidopsis heat shock transcription factor A2 as a key regulator in response to several types of environmental stress, *Plant J.* 48 (2006) 535–547.
- [58] Y. Yoshida, Y. Sakuma, D. Todaka, K. Maruyama, F. Qin, J. Mizoi, S. Kidokoro, Y. Fujita, K. Shinozaki, K. Yamaguchi-Shinozaki, Functional analysis of an Arabidopsis heat-shock transcription factor *HsfA3* in the transcriptional cascade downstream of the *DREB2A* stress-regulatory system, *Biochem. Biophys. Res. Commun.* 368 (2008) 515–521.
- [59] Z. Li, L. Zhang, A. Wang, X. Xu, J. Li, Ectopic overexpression of *SlHsfA3*, a heat stress transcription factor from tomato, confers increased thermotolerance and salt hypersensitivity in germination in transgenic Arabidopsis, *PLoS One* 8 (2013), e54880.
- [60] S. Zhang, Z.S. Xu, P. Li, L. Yang, Y. Wei, M. Chen, L. Li, G. Zhang, Y. Ma, Overexpression of *TaHSF3* in transgenic Arabidopsis enhances tolerance to extreme temperatures, *Plant Mol. Biol. Report.* 31 (2013) 688–697.
- [61] P.K. Agarwal, P. Agarwal, M.K. Reddy, S.K. Sopory, Role of DREB transcription factors in abiotic and biotic stress tolerance in plants, *Plant Cell Rep.* 25 (2006) 1263–1274.
- [62] X.M. Liu, X.C. Nguyen, K.E. Kim, H.J. Han, J. Yoo, K. Lee, W.S. Chung, Phosphorylation of the zinc finger transcriptional regulator *ZAT6* by *MPK6* regulates Arabidopsis seed germination under salt and osmotic stress, *Biochem. Biophys. Res. Commun.* 430 (3) (2013) 1054–1059.
- [63] C.C. Englbrecht, H. Schoof, S. Böhm, Conservation, diversification and expansion of C2H2 zinc finger proteins in the *Arabidopsis thaliana* genome, *BMC Genomics* 5 (2004) 39.
- [64] S. Davletova, K. Schlauch, J. Coutu, R. Mittler, The zinc-finger protein *Zat12* plays a central role in reactive oxygen and abiotic stress signaling in Arabidopsis, *Plant Physiol.* 139 (2005) 847–856.
- [65] L. Rizhsky, S. Davletova, H. Liang, R. Mittler, The zinc finger protein *Zat12* is required for cytosolic ascorbate peroxidase 1 expression during oxidative stress in Arabidopsis, *J. Biol. Chem.* 279 (2004) 11736–11743.
- [66] G. Miller, V. Shulaev, R. Mittler, Reactive oxygen signaling and abiotic stress, *Physiol. Plant.* 133 (2008) 481–489.
- [67] H. Sakamoto, K. Maruyama, Y. Sakuma, T. Meshi, M. Iwabuchi, K. Shinozaki, K. Yamaguchi-Shinozaki, Arabidopsis CYS2/HIS2-type zinc-finger proteins function as transcription repressors under drought, cold, and high-salt stress conditions, *Plant Physiol.* 136 (2004) 2734–2746.