



Contents lists available at ScienceDirect

Diabetes & Metabolic Syndrome: Clinical Research & Reviews

journal homepage: www.elsevier.com/locate/dsx

Review

Molecular mechanisms of trehalose in modulating glucose homeostasis in diabetes

Habib Yaribeygi^{a, **}, Alijan Yaribeygi^b, Thozhukat Sathyapalan^c, Amirhossein Sahebkar^{d, e, f, *}^a Research Center of Physiology, Semnan University of Medical Sciences, Semnan, Iran^b Chronic Kidney Disease Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran^c Department of Academic Diabetes, Endocrinology and Metabolism, Hull York Medical School, University of Hull, Hull, HU3 2JZ, UK^d Biotechnology Research Center, Pharmaceutical Technology Institute, Mashhad University of Medical Sciences, Mashhad, Iran^e Neurogenic Inflammation Research Center, Mashhad University of Medical Sciences, Mashhad, Iran^f School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

ARTICLE INFO

Article history:

Received 24 April 2019

Accepted 22 May 2019

Keywords:

Diabetes mellitus

Trehalose

Oxidative stress

Glucose

Homeostasis

ABSTRACT

Diabetes mellitus is the most prevalent metabolic disorder contributing to significant morbidity and mortality in humans. Many preventative and therapeutic agents have been developed for normalizing glycemic profile in patients with diabetes. In addition to various pharmacologic strategies, many non-pharmacological agents have also been suggested to improve glycemic control in patients with diabetes. Trehalose is a naturally occurring disaccharide which is not synthesized in human but is widely used in food industries. Some studies have provided evidence indicating that it can potentially modulate glucose metabolism and help to stabilize glucose homeostasis in patients with diabetes. Studies have shown that trehalose can significantly modulate insulin sensitivity via at least 7 molecular pathways leading to better control of hyperglycemia. In the current study, we concluded about possible anti-hyperglycemic effects of trehalose suggesting trehalose as a potentially potent non-pharmacological agent for the management of diabetes.

© 2019 Diabetes India. Published by Elsevier Ltd. All rights reserved.

1. Introduction

The global prevalence of diabetes mellitus (DM) is rising exponentially [1]. This chronic disorder has paramount effects on various metabolic pathways as well as on the physiologic function of most organs and hence responsible for a wide variety of diabetes-induced tissue dysfunctions known as diabetic complications [2,3]. DM acts as a potent upstream event for many pathophysiologic mechanisms such as oxidative stress, inflammation, apoptosis and fibrosis which in turn induce the onset and progression of various forms of diabetic complications [4,5]. Diabetes and its related complications impose a significant economic burden

to the health systems and are responsible for many cases of disabilities and death in human [6,7]. Therefore, normalizing glycemic profile and management of chronic hyperglycemia by pharmacological and non-pharmacological approaches are of paramount importance [8–10].

Trehalose is a disaccharide carbohydrate which is synthesized in many organisms ranging from bacteria to plants [11]. Although this natural sweetener molecule could not be synthesized in humans, it has some biological roles and effects on various metabolic pathways after consumption [12–14]. It is used extensively in the food industry, biotechnology and pharmaceutical industry. There is some evidence suggesting that it can potentially modulate glucose homeostasis [15,16]. In the current review, we assessed the possible links between trehalose usage and DM in humans and whether it can modulate hyperglycemia toward the physiologic state.

2. Trehalose, biochemistry and biology

Trehalose is a naturally-occurring, non-reducing sugar formed by two D-glucose molecules linked by α -1,1-glycosidic bindings

* Corresponding author. Department of Medical Biotechnology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, P.O. Box: 91779-48564, Iran.

** Corresponding author. Research Center of Physiology, Semnan University of Medical Sciences, Semnan, Iran

E-mail addresses: habib.yari@yahoo.com (H. Yaribeygi), sahebkar@mums.ac.ir, amir_saheb2000@yahoo.com (A. Sahebkar).

[17,18]. This unreactive disaccharide is synthesized by a wide variety of species ranging from prokaryotes, fungi, invertebrates, insects, plants, and yeast and has significant metabolic roles including storage of energy [18]. The synthetic form of Trehalose was produced by enzymatic processes for the first time in 1995 [19], and at present is extensively used in food industries as a stabilizer for protecting foods against dryness, freezing and osmotic pressure stresses [16]. In comparison with sucrose, trehalose has lower sweetener potency and is broken into two glucose molecules by the enzyme trehalase and provides a high amount of energy [18,20].

It also acts as an osmo-protectant especially against ethanol and osmotic pressures. It provides a stabilizing effect for the cellular membrane by increasing the plasma membrane tolerance to dehydration, shriveling and temperature shock [21]. It has suggested that non-reactivity and high stability properties of trehalose are due to the very low energy of the glycoside oxygen bindings to its two hexose rings [21]. Although trehalase enzyme is widely expressed in human and founded in the brush border of mucosal intestinal cells, no evidence exists about the trehalose production in human [18].

3. Molecular mechanisms involved in insulin resistance

Insulin resistance; which is a key event in the development of dysmetabolic states such as non-alcoholic fatty liver disease, DM and metabolic syndrome; is a complicated condition involving several molecular mechanisms [22,23]. In a previous review we found that at least 10 molecular pathways are included in peripheral resistance to insulin including reduced response to circulatory insulin such as up-regulation of PTP1B (protein-tyrosine phosphatase 1B), inflammatory mediators and adipokines secretion, oxidative stress, ER (endoplasmic reticulum) stress, lower IRS-1 (insulin receptor substrates-1) phosphorylation, adipocytes metabolism, augmented insulin degradation, defect in GLUT-4 (glucose transporter-4) activity, lower capacity of receptors to binding to insulin and mitochondrial dysfunction [23]. As a result, any pharmacologic or non-pharmacological agents which modulate these molecular mechanisms may be also able to regulate insulin sensitivity [8,23–25].

4. Possible links between trehalose and diabetes mellitus

Some recent evidence have shown that trehalose can potentially modulate glucose homeostasis (Fig. 1) [15,16]. In following sections, we examine these potentials of trehalose (Table 1).

4.1. Trehalose and insulin sensitivity

Trehalose can potentially increase insulin sensitivity and improve insulin resistance in diabetic milieu [26]. This effect is exerted either directly on glucose signaling pathways or indirectly via alleviating pathophysiological pathways such as oxidative stress,




Fig. 1. Molecular mechanisms by which trehalose prevents and reverses the effects of DM on glucose homeostasis.

inflammation and/or improvement in lipid metabolism [26]. Arai et al., in 2010 demonstrated that trehalose induces insulin sensitivity by several mechanisms including a reduction in inflammatory mediators such as MCP-1 (monocyte chemotactic protein-1), TNF- α (tumor necrosis factor-alpha), and PAI-1 (plasminogen activator inhibitor-1) and improvement in adipocytes function by inhibition of lipid accumulation [26]. Arai and coworkers in 2013 presented further evidence indicating that 8 weeks trehalose consumption mitigates insulin resistance by adiponectin secretion, lipid profile correction and up-regulation of the IRS-1 and IRS-2 mRNA expression in obese mice [27]. Mizote and colleagues in 2016 found in a clinical trial that trehalose improved insulin resistance by adiponectin release and PAI-1 down-regulation in people at risk of metabolic syndrome [15]. Higgins et al., in 2018 showed that trehalose is a potent analog for ALOXE3 (Epidermis-type Lipooxygenase 3) which in turn induces insulin sensitivity via PPAR- γ (peroxisome proliferator-activated receptor gamma) dependent mechanism in mice [28]. These evidence strongly suggest that trehalose induces insulin sensitivity in hyperlipidemia or in the hyperglycemic milieu.

4.2. Trehalose and postprandial glucose/insulin secretion

Reducing insulin secretion is the most known effects of trehalose on glucose homeostasis and is supported by various studies [26,29,30]. Arai and coworkers in 2010 observed that trehalose markedly reduced the fasting insulin levels in comparison with glucose and maltose in rats [26]. Oku et al., in 2000 reported that trehalose intake has remarkable effects on blood glucose in people with higher levels of trehalase activity than people with low trehalase activity. Trehalose had a potent inhibitory effect on insulin release in these subjects [29]. Also, Maki et al., in 2009 reported that trehalose intake in obese men elicited lower postprandial insulin secretory response than to that of glucose [30]. Yoshizane and colleagues in 2017 observed in healthy volunteers that trehalose ingestion did not induce a rapid rise in postprandial glucose and did not stimulate GIP (gastric inhibitory protein) secretion [16]. These evidence clearly confirm that trehalose has weaker stimulatory effects on postprandial insulin release in comparison with other sugars [26,30]. It has also been suggested that the trehalase acts in a manner dependent on insulin level demonstrating the intricate relationship between insulin levels and trehalase activity [31]. These finding also showed that trehalose reduces insulin dependent postprandial adipocyte accumulation which is regularly seen after carbohydrate consumption [16,26].

4.3. Trehalose and glucose metabolism

Trehalose has shown to normalize glucose metabolic pathways [32,33]. Sato et al., in 1999 demonstrated that animals consuming parenteral trehalose have improved glucose metabolism and nutritional indices than after consuming other disaccharides [32]. Also, van Can and coworkers in 2012 provided data indicating trehalose improves glucose metabolism in patients with glucose intolerance [34]. Yasugi et al., in 2017 suggested that animals using trehalose as nutrient have improved metabolism which is gained via adaptation [35]. These evidence implied that trehalose may directly modulate glucose metabolism but the exact mechanisms need to be elucidated in further studies.

4.4. Trehalose and lipid metabolism

Correction of the lipid profile may be one of the other possible mechanisms by which trehalose improves glucose homeostasis [26]. There is growing evidence that trehalose directly modulates

Table 1

Various influences of trehalose on DM, trehalose consumption can modulate glucose homeostasis via at least 7 molecular mechanisms.

Molecular Pathway	Influence of Trehalose	References
Insulin Sensitivity	Improves insulin sensitivity by several mechanisms including IRS up-regulation, PPAR- γ dependent mechanism and ameliorating other pathophysiological pathways	[26–28]
Insulin Secretion	Reduces postprandial insulin release	[26; 29; 30]
Glucose Metabolism	Improves glucose metabolism and regulates postprandial glucose levels	[32–35]
Lipid Metabolism	Corrects lipid metabolism by modulating postprandial insulin release	[15; 26; 27]
Islet Function	Improves beta cell function	[36–40]
Oxidative Stress	Attenuates oxidative stress leading to improved insulin sensitivity	[42–46]
Inflammatory Cytokines	Prevent/suppresses inflammatory responses	[26; 45; 50; 51]

IRSs = insulin receptor substrates, PPAR- γ = Peroxisome proliferator-activated receptor gamma.

secretion of adipokines and increases adiponectin release leading to improvement in lipid metabolism [15,27]. Since sub-optimal lipid profile is closely linked to insulin resistance disorders as well as DM; we suggest that any improvement in lipid metabolism could also effectively improve insulin sensitivity [15].

4.5. Trehalose and improvement in islet function

Trehalose consumption could improve pancreatic islet function in several ways [36,37]. Emerging evidence indicates that trehalose prevents apoptotic processes, which are involved in beta cell dysfunction [36–38]. Beattie and coworkers in 1997 provided first direct evidence suggesting long-term trehalose consumption improves pancreatic beta cells' efficiency in patients with diabetes [37]. Pan et al., in 2018 demonstrated that trehalose is able to suppress apoptosis in diabetic animals [38]. Lin et al., in 2016 established an experimental study confirming trehalose diminishes pathogenic mechanisms such as apoptosis and pyroptosis probably by its anti-oxidant and anti-inflammatory potentials in pancreatic cells of diabetic rats [39]. This beneficial effect was previously suggested by Xu and coworkers in 2013 in neuronal cells where they showed that trehalose improved diabetes-induced neuronal defects by modulating the autophagy process [40]. Chen and colleagues in 2016 reported that lower concentration of trehalose markedly improves islet function by ameliorating islet amyloid polypeptide synthesis in patients with diabetes [36]. These data strongly suggest that trehalose consumption may improve islet function and glucose homeostasis.

4.6. Trehalose and oxidative stress

Free radical overload and oxidative stress occur in patients with diabetes [41]. Metabolic derangements which develop during DM markedly induces free radical generation which in turn results in an increase in peripheral resistance to insulin [32,41]. Some experiments suggest that trehalose can potentially reduce oxidative stress [42]. Mizunoe and colleagues in 2018 showed that trehalose protects against oxidative damages by Nrf2 (nuclear factor erythroid 2-related factor 2; a nuclear factor regulates antioxidant elements expression)-dependent antioxidant elements up-regulation in cell lines of mouse hepatocytes [42]. Alvarez-Peral et al., in 2002 demonstrated that trehalose has a protective role against oxidative stress in *Candida albicans* [43]. Moreover, Tang et al., in 2017 reported that trehalose diminished mitochondria-induced oxidative stress by regulation of the autophagy process in patients with osteoarthritis [44]. Echigo et al., in 2012 found that trehalose improved oxidative damages in the brain after subarachnoid bleeding in experimental studies, both in vitro and in vivo [45]. These data in addition to other evidence demonstrates potent antioxidative effects of trehalose which contributes to the potential

antihyperglycaemic effects of this non-reducing disaccharide [46].

4.7. Trehalose and inflammatory cytokines

Inflammatory mediators have significant roles in the pathophysiology of many complications of DM [47,48]. It has been well established that the expression/release of inflammatory mediators is up-regulated in patients with insulin resistance [4,49]. There is growing evidence that trehalose inhibits inflammatory responses by ameliorating inflammatory mediators [26,50]. Consumption of trehalose markedly reduced inflammatory cytokines such as TNF- α , MCP-1 and PAI-1 in experimental studies [26,51]. In addition, Echigo et al., in 2012 showed that trehalose ameliorated the inflammation in in-vitro and in-vivo models of subarachnoid hemorrhage [45]. This suggests that intake of trehalose improves insulin sensitivity at least partly via attenuating inflammatory responses [51].

4.8. Other possible links

There is some evidence suggesting other potential relationships between trehalose consumption and insulin sensitivity such as thermogenesis [52], stabilizing insulin [53], and improvement in mitochondrial dysfunction [44]. But these molecular pathways need further investigation by in vivo and in vitro studies.

Trehalose has been shown to inhibit multiple glucose transporter (GLUT) receptors through binding in the inward open conformation of these receptors [54,55]. By inhibiting GLUT function trehalose could potentially prevent excess glucose absorption in the intestine. It has also reported that trehalose reduces postprandial glycemic excursions in patients with impaired glucose tolerance [56]. This suggests that ingestion of trehalose with daily meals could potentially prevent hyperglycemia and development of diabetes.

5. Conclusion

Trehalose is a non-reducing carbohydrate widely produced by different species but not in the human. It has lower sweetener potency than sucrose and is broken down into two molecules of glucose by trehalase enzyme activity. This enzyme is expressed in humans and is localized in the brush border of endothelial cells of the intestine. In addition to providing energy in other species, trehalose can modulate glucose homeostasis by at least 7 molecular pathways. Data suggests that trehalose consumption improve hyperglycemic milieu by ameliorating pathophysiological mechanisms such as oxidative stress, inflammation and apoptosis, improving beta cell function, reducing postprandial insulin release and normalizing lipid profile. These molecular mechanisms suggest the role of trehalose as a potential non-pharmacological agent for the

management of glycaemia in patients with diabetes, however, more clinical trials are needed to assess this.

Author's contribution

AS conceived the study. HY and AY wrote the manuscript draft. AS and TS critically revised the draft. All authors approved the manuscript for submission.

Conflicts of interest

The authors clearly declare that have no conflict of interest in this study.

References

- Mayer-Davis EJ, Lawrence JM, Dabelea D, Divers J, Isom S, Dolan L, et al. Incidence trends of type 1 and type 2 diabetes among youths, 2002–2012. *N Engl J Med* 2017;376:1419–29.
- Yaribeygi H, Butler AE, Barreto GE, Sahebkar A. Antioxidative potential of antidiabetic agents: a possible protective mechanism against vascular complications in diabetic patients. *J Cell Physiol* 2019;234:2436–46.
- Abraham TM, Pencina KM, Pencina MJ, Fox CS. Trends in diabetes incidence: the framingham heart study. *Diabetes Care* 2015;38:482–7.
- Yaribeygi H, Atkin SL, Pirro M, Sahebkar A. A review of the anti-inflammatory properties of antidiabetic agents providing protective effects against vascular complications in diabetes. *J Cell Physiol* 2019;234:8286–94.
- Yaribeygi H, Butler AE, Sahebkar A. Aerobic exercise can modulate the underlying mechanisms involved in the development of diabetic complications. *J Cell Physiol* 2019;234:12508–15.
- Association AD. Economic costs of diabetes in the US in 2017. *Diabetes Care* 2018;41:917–28.
- Rosella L, Lebenbaum M, Fitzpatrick T, O'reilly D, Wang J, Booth G, et al. Impact of diabetes on healthcare costs in a population-based cohort: a cost analysis. *Diabet Med* 2016;33:395–403.
- Yaribeygi H, Zare V, Butler AE, Barreto GE, Sahebkar A. Antidiabetic potential of saffron and its active constituents. *J Cell Physiol* 2019;8610–7.
- Yaribeygi H, Atkin SL, Simental-Mendia LE, Sahebkar A. Molecular mechanisms by which aerobic exercise induces insulin sensitivity. *J Cell Physiol* 2019;234:12385–92.
- Yaribeygi H, Atkin SL, Ramezani M, Sahebkar A. A review of the molecular pathways mediating the improvement in diabetes mellitus following caloric restriction. *J Cell Physiol* 2019;234:8436–42.
- Elbein AD, Pan Y, Pastuszak I, Carroll D. New insights on trehalose: a multifunctional molecule. *Glycobiology* 2003;13:17R–27R.
- Figuerola CM, Feil R, Ishihara H, Watanabe M, Kölling K, Krause U, et al. Trehalose 6-phosphate coordinates organic and amino acid metabolism with carbon availability. *Plant J* 2016;85:410–23.
- Eleutherio E, Panek A, De Mesquita JF, Trevisol E, Magalhães R. Revisiting yeast trehalose metabolism. *Curr Genet* 2015;61:263–74.
- Wang X, Du Y, Yu D. Trehalose phosphate synthase 5-dependent trehalose metabolism modulates basal defense responses in *Arabidopsis thaliana*. *J Integr Plant Biol* 2019;61:509–27.
- Mizote A, Yamada M, Yoshizane C, Arai N, Maruta K, Arai S, et al. Daily intake of trehalose is effective in the prevention of lifestyle-related diseases in individuals with risk factors for metabolic syndrome. *J Nutr Sci Vitaminol* 2016;62:380–7.
- Yoshizane C, Mizote A, Yamada M, Arai N, Arai S, Maruta K, et al. Glycemic, insulinemic and incretin responses after oral trehalose ingestion in healthy subjects. *Nutr J* 2017;16:9.
- Thammahong A, Puttikamonkul S, Perfect JR, Brennan RG, Cramer RA. Central role of the trehalose biosynthesis pathway in the pathogenesis of human fungal infections: opportunities and challenges for therapeutic development. *Microbiol Mol Biol Rev* 2017;81: e00053-16.
- Feofilova E, Usov A, Mysyakina I, Kochkina G. Trehalose: chemical structure, biological functions, and practical application. *Microbiology* 2014;83:184–94.
- Sugimoto T. Production of trehalose by starch-glycation method with enzymes and its use. *Food Industry* 1995;38:34–9.
- Moriano ME, Alamprese C. Honey, trehalose and erythritol as sucrose-alternative sweeteners for artisanal ice cream. A pilot study. *LWT-Food Science and Technology* 2017;75:329–34.
- Schiraldi C, Di Lernia I, De Rosa M. Trehalose production: exploiting novel approaches. *Trends Biotechnol* 2002;20:420–5.
- Freeman AM, Pennings N. Insulin resistance. *StatPearls* [internet]. StatPearls Publishing; 2018.
- Yaribeygi H, Farrokhi FR, Butler AE, Sahebkar A. Insulin resistance: review of the underlying molecular mechanisms. *J Cell Physiol* 2019;234:8152–61.
- Yaribeygi H, Katsiki N, Behnam B, Iranpanah H, Sahebkar A. MicroRNAs and type 2 diabetes mellitus: molecular mechanisms and the effect of antidiabetic drug treatment. *Metabolism* 2018;87:48–55.
- Yaribeygi H, Atkin SL, Sahebkar A. Mitochondrial dysfunction in diabetes and the regulatory roles of antidiabetic agents on the mitochondrial function. *J Cell Physiol* 2019;234:8402–10.
- Arai C, Arai N, Mizote A, Kohno K, Iwaki K, Hanaya T, et al. Trehalose prevents adipocyte hypertrophy and mitigates insulin resistance. *Nutr Res (NY)* 2010;30:840–8.
- Arai C, MIYAKE M, Matsumoto Y, Mizote A, Yoshizane C, Hanaya Y, et al. Trehalose prevents adipocyte hypertrophy and mitigates insulin resistance in mice with established obesity. *J Nutr Sci Vitaminol* 2013;59:393–401.
- Higgins CB, Zhang Y, Mayer AL, Fujiwara H, Stothard AI, Graham MJ, et al. Hepatocyte ALOXE3 is induced during adaptive fasting and enhances insulin sensitivity by activating hepatic PPAR γ . *JCI insight* 2018;3.
- Oku T, Nakamura S. Estimation of intestinal trehalase activity from a laxative threshold of trehalose and lactulose on healthy female subjects. *Eur J Clin Nutr* 2000;54:783.
- Maki KC, Kanter M, Rains TM, Hess SP, Geohas J. Acute effects of low insulinemic sweeteners on postprandial insulin and glucose concentrations in obese men. *Int J Food Sci Nutr* 2009;60:48–55.
- Richards A, Krakowka S, Dexter L, Schmid H, Wolterbeek A, Waalkens-Berendsen D, et al. Trehalose: a review of properties, history of use and human tolerance, and results of multiple safety studies. *Food Chem Toxicol* 2002;40:871–98.
- Sato S, Okamoto K, Minami R, Kohri H, Yamamoto S. Trehalose can be used as a parenteral saccharide source in rabbits. *J Nutr* 1999;129:158–64.
- Seo Y, Kingsley S, Walker G, Mondoux MA, Tissenbaum HA. Metabolic shift from glycogen to trehalose promotes lifespan and healthspan in *Caenorhabditis elegans*. *Proc Natl Acad Sci Unit States Am* 2018;115:E2791–800.
- van Can JG, van Loon LJ, Brouns F, Blaak EE. Reduced glycaemic and insulinemic responses following trehalose and isomaltulose ingestion: implications for postprandial substrate use in impaired glucose-tolerant subjects. *Br J Nutr* 2012;108:1210–7.
- Yasugi T, Yamada T, Nishimura T. Adaptation to dietary conditions by trehalose metabolism in *Drosophila*. *Sci Rep* 2017;7:1619.
- Chen C-H, Yao T, Zhang Q, He Y-M, Xu L-H, Zheng M, et al. Influence of trehalose on human islet amyloid polypeptide fibrillation and aggregation. *RSC Adv* 2016;6:15240–6.
- Beattie GM, Crowe JH, Lopez AD, Cirulli V, Ricordi C, Hayek A. Trehalose: a cryoprotectant that enhances recovery and preserves function of human pancreatic islets after long-term storage. *Diabetes* 1997;46:519–23.
- Pan H, Ding Y, Yan N, Nie Y, Li M, Tong L. Trehalose prevents sciatic nerve damage to and apoptosis of Schwann cells of streptozotocin-induced diabetic C57BL/6j mice. *Biomed Pharmacother* 2018;105:907–14.
- Lin C-F, Kuo Y-T, Chen T-Y, Chien C-T. Quercetin-rich Guava (*Psidium guajava*) juice in combination with trehalose reduces autophagy, apoptosis and pyroptosis formation in the kidney and pancreas of type II diabetic rats. *Molecules* 2016;21:334.
- Xu C, Li X, Wang F, Weng H, Yang P. Trehalose prevents neural tube defects by correcting maternal diabetes-suppressed autophagy and neurogenesis. *Am J Physiol Endocrinol Metab* 2013;305:E667–78.
- Yaribeygi H, Atkin SL, Sahebkar A. A review of the molecular mechanisms of hyperglycemia-induced free radical generation leading to oxidative stress. *J Cell Physiol* 2019;234:1300–12.
- Mizunoe Y, Kobayashi M, Sudo Y, Watanabe S, Yasukawa H, Natori D, et al. Trehalose protects against oxidative stress by regulating the Keap1–Nrf2 and autophagy pathways. *Redox biology* 2018;15:115–24.
- Alvarez-Peral FJ, Zaragoza O, Pedreno Y, Argüelles J-C. Protective role of trehalose during severe oxidative stress caused by hydrogen peroxide and the adaptive oxidative stress response in *Candida albicans*. *Microbiology* 2002;148:2599–606.
- Tang Q, Zheng G, Feng Z, Chen Y, Lou Y, Wang C, et al. Trehalose ameliorates oxidative stress-mediated mitochondrial dysfunction and ER stress via selective autophagy stimulation and autophagic flux restoration in osteoarthritis development. *Cell Death Dis* 2017;8: e3081.
- Echigo R, Shimohata N, Karatsu K, Yano F, Kayasuga-Kariya Y, Fujisawa A, et al. Trehalose treatment suppresses inflammation, oxidative stress, and vasospasm induced by experimental subarachnoid hemorrhage. *J Transl Med* 2012;10:80.
- Hu J-H, Zan L-S, Zhao X-L, Li Q-W, Jiang Z-L, Li Y-K, et al. Effects of trehalose supplementation on semen quality and oxidative stress variables in frozen-thawed bovine semen. *J Anim Sci* 2010;88:1657–62.
- Yaribeygi H, Katsiki N, Butler AE, Sahebkar A. Effects of antidiabetic drugs on NLRP3 inflammasome activity, with a focus on diabetic kidneys. *Drug Discov Today* 2019;24:256–62.
- Yaribeygi H, Atkin SL, Sahebkar A. Interleukin-18 and diabetic nephropathy: a review. *J Cell Physiol* 2019;234:5674–82.
- Yaribeygi H, Mohammadi MT, Rezaee R, Sahebkar A. Crocin improves renal function by declining Nox-4, IL-18, and p53 expression levels in an experimental model of diabetic nephropathy. *J Cell Biochem* 2018;119:6080–93.
- Arai C, Kohguchi M, Akamatsu S, Arai N, Yoshizane C, Hasegawa N, et al. Trehalose suppresses lipopolysaccharide-induced osteoclastogenesis bone marrow in mice. *Nutr Res* 2001;21:993–9.
- Taya K, Hirose K, Hamada S. Trehalose inhibits inflammatory cytokine production by protecting I κ B- α reduction in mouse peritoneal macrophages. *Arch Oral Biol* 2009;54:749–56.
- Zhang Y, Higgins CB, Mayer AL, Mysorekar IU, Razani B, Graham MJ, et al.

- TFEB-dependent induction of thermogenesis by the hepatocyte SLC2A inhibitor trehalose. *Autophagy* 2018;14:1959–75.
- [53] Mansfield KM, Maynard HD. Site-specific insulin-trehalose glycopolymer conjugate by grafting from strategy improves bioactivity. *ACS Macro Lett* 2018;7:324–9.
- [54] Mayer AL, Higgins CB, Heitmeier MR, Kraft TE, Qian X, Crowley JR, et al. SLC2A8 (GLUT8) is a mammalian trehalose transporter required for trehalose-induced autophagy. *Sci Rep* 2016;6:38586.
- [55] DeBosch BJ, Heitmeier MR, Mayer AL, Higgins CB, Crowley JR, Kraft TE, et al. Trehalose inhibits solute carrier 2A (SLC2A) proteins to induce autophagy and prevent hepatic steatosis. *Sci Signal* 2016;9:ra21.
- [56] van Can JG, van Loon LJ, Brouns F, Blaak EE. Reduced glycaemic and insulinaemic responses following trehalose and isomaltulose ingestion: implications for postprandial substrate use in impaired glucose-tolerant subjects. *Br J Nutr* 2012;108:1210–7.