



Glutamate receptor metabotropic 7 (*GRM7*) gene polymorphisms in mood disorders and attention deficit hyperactive disorder



Rezvan Noroozi^{a,b}, Mohammad Taheri^{c,*}, Mir Davood Omrani^c, Soudeh Ghafouri-Fard^{a,**}

^a Department of Medical Genetics, Shahid Beheshti University of Medical Sciences, Tehran, Iran

^b Pytochemistry Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

^c Urogenital Stem Cell Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

ARTICLE INFO

Keywords:

Glutamate receptor metabotropic 7
Bipolar I disorder
Bipolar II disorder
Major depressive disorder
Attention deficit hyperactive disorder

ABSTRACT

L-glutamate is the chief excitatory neurotransmitter in the central nervous system (CNS) which activates metabotropic receptors including the metabotropic glutamate receptor GRM7. Single nucleotide polymorphisms (SNPs) within *GRM7* gene have been associated with several psychiatric conditions. In the present study, we assessed association between two *GRM7* SNPs (rs6782011 and rs779867) and two neuropsychiatric disorders including attention deficit hyperactive disorder (ADHD) and mood disorders. There were no significant differences in genotype, allele and haplotypes frequencies of the rs6782011 and rs779867 between bipolar disorder 1 (BPD1) patients and controls. The CC genotype of the rs6782011 was significantly associated with BPD2 in recessive model (OR (95% CI) = 1.78 (1.09–2.91), adjusted P value = 0.04) and with ADHD in dominant and co-dominant models (OR (95% CI) = 1.98 (1.11–3.53), adjusted P value = 0.04; OR (95% CI) = 2.27 (1.23–4.17), adjusted P value = 0.04 respectively). The C G haplotype (rs6782011 and rs779867 respectively) was more prevalent among both BPD2 patients (OR (95%CI) = 2.03 (1.36–3.01), adjusted P value = 0.002) and MDD patients (OR (95%CI) = 2.08 (1.37–3.16), adjusted P value = 0.002) compared with controls. The current study provides further evidences for participation of *GRM7* variants in conferring risk of neuropsychiatric disorders.

1. Introduction

L-glutamate as the chief excitatory neurotransmitter in the central nervous system (CNS) can induce both ligand gated ion channels (ionotropic receptors) and G-protein coupled (metabotropic) receptors. The latter include three groups of receptors with distinct functions. While group I receptors stimulate phospholipase C, group II and III receptors mostly suppress the cyclic AMP cascade (Kew and Kemp, 2005). Glutamatergic neurotransmission participates in many features of normal brain activities and can be disturbed in many neurological diseases (Zhou and Danbolt, 2014). Defect in glutamate signaling has been suggested as an underlying pathology in some patients with attention deficit hyperactive disorder (ADHD) (Adler et al., 2012). Supporting evidences for participation of glutamate signaling in ADHD have been emerged from human association studies indicating association between ADHD and glutamate receptor subunit gene *GRIN2B* variants (Dorval et al., 2007), animal studies showing dys-regulation of glutamate effect of the dopamine system in ADHD rat model (Warton

et al., 2009) and the reported lower concentration of glutamate and glutamine in the basal ganglia of ADHD adults compared with normal individuals (Maltezos et al., 2014). Dys-regulation of glutamatergic system is invariably involved in mood disorders including major depressive disorder (MDD) and bipolar disorder (BPD) in a way that the glutamate hypothesis of mood disorders is predicted to supplement the predominant monoamine theory (Jun et al., 2014). Depressive and manic episodes have been associated with alteration of the glutamine/glutamate ratio in divergent ways (Yuksel and Ongur, 2010). Other clues to “glutamate hypothesis” of mood disorders have been provided by the predominance of glutamatergic neurons and synapses in many brain areas, the role of this neurotransmitter in the cognition and emotion, reported aberration of this pathway in many limbic/cortical zones of depressed patients and association between abnormal glutamatergic signaling and both structural and functional alterations in the brain (Yuksel and Ongur, 2010).

The metabotropic glutamate receptor GRM7 is the most greatly conserved group III receptors has a protective role against neuron

* Corresponding author.

** Corresponding author.

E-mail addresses: mohammad_823@yahoo.com (M. Taheri), s.ghafourifard@sbmu.ac.ir (S. Ghafouri-Fard).

Table 1

The detailed information of SNPs.

SNP	Position	Minor Allele	Minor allele frequency	Minor allele count	Type
rs6782011	Chr 3:7457960	T	0.41	2046	Intron variant
rs779867	Chr 3:7442784	A	0.36	1785	Intron variant

Table 2

Nucleotide sequence of primer pairs and PCR protocol.

SNP	Primer sequence	Tm	Annealing temperature	PCR product size (bp)
rs6782011	Forward inner primer (T allele): GCTCTGACCAAATTACAAAATATATGTGGT	63 °C	57 °C	191 bp (T allele)
	Reverse inner primer (C allele): CACTCTGAATATTAGTACTCAAACAGGGG	63 °C		261 bp (C allele)
	Forward outer primer: GTTAGAACATTTGGACTATAAGCATGGC	63 °C		392 bp (two outer primers)
	Reverse outer primer: ATAATAAACCCAGTCTTCTGCATCAACGT	63 °C		
rs779867	Forward inner primer (A allele): AAACCAGGGTTTCCACTCTCATGTAAA	65 °C	58 °C	163 bp (A allele)
	Reverse inner primer (G allele): CATTAATCCAAGAGCATCTGTAAGCCC	65 °C		244 bp (G allele)
	Forward outer primer: GATCAAGATGATATAAGGGGAAACAGG	65 °C		353 bp (two outer primers)
	Reverse outer primer: CTAGGTTTCATCCAGGAAGGGACTAAAG	65 °C		

excitotoxicity via suppression of the second messenger adenylate cyclase and reduction of N-methyl-D-aspartic acid (NMDA) receptor function (Gu et al., 2012). Several linkage studies have reported a risk locus for BPD disorder in adjacency of the *GRM7* genomic locus (Fallin et al., 2005; Etain et al., 2006; Tang et al., 2011). Others have linked this locus with MDD (Pergadia et al., 2011). We previously assessed association between two *GRM7* single nucleotide polymorphisms (SNPs) (rs6782011 and rs779867) and autism spectrum disorder (ASD) in Iranian population and found supportive evidence of such associations (Noroozi et al., 2016). In the current study, we genotyped these SNPs in a population of BPD, MDD and ADHD patients and age-/sex-matched healthy subjects to find their association with these mental disorders in Iranian population.

2. Material and methods

2.1. Subjects

Sample size was calculated based on the usual expectations for a case-control study i.e. 95% confidence ($\alpha = 0.05$), 70 or 80% power ($\beta = 20$ or %30) in 1:1 ratio, and least extreme Odds Ratio to be detected as 2.0. The hypothetical proportion of controls with exposure to risk allele was supposed to be 0.36 according to the minor allele frequency (MAF) of the rs779867 as provided by the dbSNP database (Dean et al., 2009). As a result, sample size was calculated around 106 persons in case and control groups to reach study power of 70% and 135 to reach power of 80%. A total of 364 patients (male = 139, female = 225) with mood disorders (BPD1, BPD2 and MDD), 250 age-matched controls (male = 84, female = 166), 108 ADHD patients (male = 81, female = 27) and 164 age-matched controls (male = 123, female = 41) participated in this study. These patients were recruited from the Farshchian Hospital, Hamadan, Iran. The MDD patients had at

least two distinct major depressive episodes. Patients' evaluation and diagnosis were based on the criteria of the Diagnostic and Statistical Manual of Mental Disorders, 4th edition (DSM-IV). The exclusion criteria were history of any sort of substance abuse, organic brain syndrome, or any general medical disease with psychiatric signs. The control samples were collected from both the above-mentioned hospital and primary and secondary schools. Exclusion criteria for the control samples included mood disorder or ADHD diagnosis and any chronic medical condition. Written informed consent forms were obtained from all study participants or their guardians. The study protocol was approved by the ethics committees of both Hamadan and Shahid Beheshti Universities of Medical Sciences.

2.2. SNP genotyping

Table 1 shows the detailed information of SNPs.

Both SNPs were genotyped using tetra-primer amplification-refractory mutation system (ARMS)-PCR technique and the results were verified by Sanger sequencing of 10% of samples in ABI 3730xl DNA analyzer (Macrogen, Korea). Table 2 shows the nucleotide sequences of primers used for genotyping, the corresponding annealing temperatures in the PCR and the expected amplicon sizes. The PCR program consisted an initial denaturing at 95 °C for 5 min; 35 cycles at 95 °C for 30 s, specific annealing temperature for 30 s, 72 °C for 1 min and a final extension at 72 °C for 5 min.

2.3. Statistical analysis

SNP Analyzer 2.0 online tool was used for appraisal of agreement of genotype distributions with Hardy-Weinberg equilibrium, haplotype approximation, linkage disequilibrium (LD) blocking and association analysis (Yoo et al., 2008). Genetic associations between mood

Table 3

Demographic data of study participants. Comparison of genotype, allele and haplotypes frequencies between patients and controls.

Variable	BPD1	BPD2	MDD	Control	ADHD	Control
Age (mean \pm SD)	41 \pm 3.2	44 \pm 8.9	49 \pm 1.0	45 \pm 8.5	10 \pm 3.4	10 \pm 8.1
Gender						
Female	62	89	74	166	27	41
Male	41	56	42	84	81	123
Total	103	145	116	250	108	164

Table 4
The detailed data of associations between genotypes and alleles of the mentioned SNPs and BPD1, BPD2, MDD and ADHD disorders respectively (Minor alleles of SNPs are shown in lower-case letters, Control[#]: the control for BPD1, BPD2, and MDD; Control[§], the control for ADHD; significant P-values are presented in boldface).

SNP	Model	Control [#] N (%)	BPD1 N (%)	OR (95% CI)	P value	Adjusted P value	BPD2 N (%)	OR (95% CI)	P value	Adjusted P value	
rs6782011	Allele	C vs. t	203 (41) 297 (59)	85 (41) 121 (59)	1.03 (0.74–1.43)	1.00	141 (49) 149 (51)	1.39 (1.03–1.85)	0.87	1.00	
	Co-dominant	CC vs. tt	44 (17.6)	18 (17)	0.97 (0.49–1.89)	1.00	40 (27.6)	1.89 (1.08–3.33)	0.96	1.00	
		CT vs. tt	115 (46)	49 (48)	1.08 (0.64–1.78)	1.00	61 (42.1)	0.91 (0.57–1.47)	0.8	1.00	
	Dominant	CC + Ct vs. tt	159 (63.6)	67 (65)	1.06 (0.66–1.72)	1.00	101 (69.7)	1.31 (0.85–2.04)	0.98	1.00	
		CC vs. Ct + tt	91 (36.4)	36 (35)	0.99 (0.54–1.81)	1.00	44 (30.3)	1.78 (1.09–2.91)	0.54	1.00	
	rs779867	Allele	a vs. G	206 (82.4) 253 (51)	85 (83) 98 (48)	0.89 (0.64–1.23)	0.93	105 (72.4) 141 (49)	0.92 (0.69–1.23)	0.46	0.93
Co-dominant		aa vs. GG	61 (24.4)	22 (21.4)	0.78 (0.4–1.52)	1.00	33 (22.8)	0.85 (0.47–1.54)	0.75	1.00	
		aG vs. GG	131 (52.4)	54 (52.4)	0.88 (0.51–1.54)	1.00	75 (51.7)	0.90 (0.54–1.47)	0.55	1.00	
Dominant		aG + aa vs. GG	192 (76.8)	76 (73.8)	0.85 (0.50–1.44)	1.00	108 (74.5)	0.88 (0.55–1.42)	0.54	1.00	
		aa vs. aG + GG	58 (23.2)	27 (26.2)	0.84 (0.48–1.46)	1.00	37 (25.5)	0.91 (0.56–1.48)	0.71	1.00	
rs6782011		Adjusted P value	MDD	113 (49)	110 (51)	1.39 (1.02–1.9)	0.04	0.08	151 (46)	1.22 (0.86–1.72)	0.26
	N (%)		119 (51)	106 (49)	1.89 (1.04–3.45)	0.08	0.17	177 (54)	0.69 (0.34–1.42)	0.02	0.04
	Adjusted P value		49 (42.2)	64 (59.3)	1.32 (0.82–2.12)	0.24	0.49	111 (67.7)	2.27 (1.23–4.17)	0.02	0.04
	N (%)		81 (69.8)	87 (80.6)	1.78 (1.06–3.01)	0.03	0.06	53 (32.3)	1.98 (1.11–3.53)	0.55	1.00
	Adjusted P value		32 (27.6)	23 (21.3)	0.82 (0.60–1.12)	0.22	0.43	40 (24.4)	0.84 (0.47–1.50)	0.67	1.00
	N (%)		84 (72.4)	102 (47)	0.63 (0.32–1.23)	0.31	0.61	161 (49)	0.93 (0.66–1.31)	0.11	0.22
	Adjusted P value	MDD	126 (54)	114 (53)	1.03 (0.60–1.75)	0.58	1.00	35 (21.3)	1.12 (0.57–2.21)	0.13	0.26
		N (%)	20 (17.2)	28 (25.9)	0.87 (0.52–1.44)	0.12	0.25	91 (55.5)	0.66 (0.38–1.13)	0.38	0.76
		Adjusted P value	86 (74.1)	74 (68.5)	0.64 (0.37–1.13)	0.12	0.25	126 (76.8)	1.29 (0.73–2.28)	0.38	0.76
		N (%)	30 (25.9)	34 (31.5)				38 (23.2)			
		Adjusted P value	20 (17.2)	28 (25.9)				35 (21.3)			
		N (%)	96 (82.8)	80 (74.1)				80 (74.1)			

disorders. Contribution of *GRM7* in the pathogenesis of ADHD has also been assessed in Korean population through transmission disequilibrium test which demonstrated biased transmission of the G allele of the rs3792452. Moreover, the GG genotype in the rs37952452 was associated with higher mean T-scores for omission errors on the continuous performance test as well as higher State-Trait Anxiety Inventory for Children (STAIC)-T and STAIC-S scores (Park et al., 2013).

We also reported higher prevalence of the C G haplotype (rs6782011 and rs779867 respectively) among both BPD2 patients and MDD patients compared with controls. While other predicted haplotype blocks of these two SNPs were associated with ASD, this haplotype was not associated with ASD in our previous study (Noroozi et al., 2016) which might reflect distinct roles for *GRM7* variants in the pathogenesis of ASD compared with mood disorder/ADHD. This hypothesis is further supported by the dissimilarity in allele/genotype frequencies of the mentioned SNPs between ASD and mood disorder/ADHD patients. Previous genome wide association studies have shown associations between *GRM7* genomic variants and both BPD (The Wellcome Trust Case Control et al., 2007; Alliey-Rodriguez et al., 2011; Sklar et al., 2008) and MDD (Sullivan et al., 2009; Shi et al., 2011; Shyn et al., 2011) in different populations. In addition, the *GRM7* rs9814881 has been linked with MDD in the Chinese Han population (Niu et al., 2017). Based on the results of the previous studies in different populations, the mentioned haplotype of these intronic variants might have functional effects on the encoded protein possibly through regulation of alternative splicing or instead contain another functional variant which is responsible for alterations in *GRM7* expression or function. Using HaploReg as a tool for discovering chromatin states, conservation, and regulatory motif modifications resulting from genomic variants (Ward and Kellis, 2012), we identified that the rs6782011 SNP would change regulatory motifs for DMRT7, Mef2 and Pax-3. The role of Pax genes in the process of neurodevelopment (Thompson and Ziman, 2011) and contribution of Mef2 in the pathogenesis of Angelman-like syndrome with neurologic manifestations (Luk, 2016) support a functional role for the rs6782011 in the assessed neuropsychiatric disorders. However, future studies are needed to verify this hypothesis.

In brief, in the current study we demonstrated associations between *GRM7* variants and both mood disorder and ADHD conditions. Such study further highlights the role of glutaminergic pathways in the pathogenesis of psychiatric disorders and warrants future studies to elaborate the underlying mechanism.

5. Conflicts of interest

The authors declare they have no conflict of interest.

Acknowledgement

The current study was supported by a grant from Shahid Beheshti University of Medical Sciences.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.neuint.2019.104483>.

References

Adler, L.A., Kroon, R.A., Stein, M., Shahid, M., Tarazi, F.I., Szegedi, A., Schipper, J., Cazorla, P., 2012. A translational approach to evaluate the efficacy and safety of the novel AMPA receptor positive allosteric modulator org 26576 in adult attention-deficit/hyperactivity disorder. *Biol. Psychiatry* 72, 971–977.

Alliey-Rodriguez, N., Zhang, D., Badner, J.A., Lahey, B.B., Zhang, X., Dinwiddie, S., Romanos, B., Pleny, N., Liu, C., Gershon, E.S., 2011. Genome-wide association study of personality traits in bipolar patients. *Psychiatr. Genet.* 21, 190–194.

Barnett, J.H., Smoller, J.W., 2009. The genetics of bipolar disorder. *Neuroscience* 164, 331–343.

Dean, A., Sullivan, K., Soe, M., 2009. Open Source Epidemiologic Statistics for Public

Health. version. Updated 2009/19/09. Available from: <http://www.openepi.com> (10 Nov 2010).

Dorval, K.M., Wigg, K.G., Crosbie, J., Tannock, R., Kennedy, J.L., Ickowicz, A., Pathare, T., Malone, M., Schachar, R., Barr, C.L., 2007. Association of the glutamate receptor subunit gene *GRIN2B* with attention-deficit/hyperactivity disorder. *Genes Brain Behav.* 6, 444–452.

Elia, J., Glessner, J.T., Wang, K., Takahashi, N., Shtir, C.J., Hadley, D., Sleiman, P.M., Zhang, H., Kim, C.E., Robison, R., Lyon, G.J., Flory, J.H., Bradford, J.P., Imielinski, M., Hou, C., Frackelton, E.C., Chiavacci, R.M., Sakurai, T., Rabin, C., Middleton, F.A., Thomas, K.A., Garris, M., Mentch, F., Freitag, C.M., Steinhausen, H.C., Todorov, A.A., Reif, A., Rothenberger, A., Franke, B., Mick, E.O., Roeyers, H., Buitelaar, J., Lesch, K.P., Banaschewski, T., Ebstein, R.P., Mulas, F., Oades, R.D., Sergeant, J., Sonuga-Barke, E., Renner, T.J., Romanos, M., Romanos, J., Warnke, A., Walitza, S., Meyer, J., Palmason, H., Seitz, C., Loo, S.K., Smalley, S.L., Biederman, J., Kent, L., Asherson, P., Anney, R.J., Gaynor, J.W., Shaw, P., Devoto, M., White, P.S., Grant, S.F., Buxbaum, J.D., Rapoport, J.L., Williams, N.M., Nelson, S.F., Faraone, S.V., Hakonarson, H., 2011. Genome-wide copy number variation study associates metabotropic glutamate receptor gene networks with attention deficit hyperactivity disorder. *Nat. Genet.* 44, 78–84.

Etain, B., Mathieu, F., Rietschel, M., Maier, W., Albus, M., Mckeon, P., Roche, S., Kealey, C., Blackwood, D., Muir, W., Bellivier, F., Henry, C., Dina, C., Gallina, S., Gurling, H., Malafosse, A., Preisig, M., Ferrero, F., Cichon, S., Schumacher, J., Ohlraun, S., Borrmann-Hassenbach, M., Propping, P., Abou Jamra, R., Schulze, T.G., Marusic, A., Dernovsek, Z.M., Giros, B., Bourgeron, T., Lemaingué, A., Bacq, D., Betard, C., Charon, C., Nothen, M.M., Lathrop, M., Leboyer, M., 2006. Genome-wide scan for genes involved in bipolar affective disorder in 70 European families ascertained through a bipolar type I early-onset proband: supportive evidence for linkage at 3p14. *Mol. Psychiatry* 11, 685–694.

Fallin, M.D., Lasseter, V.K., Avramopoulos, D., Nicodemus, K.K., Wolyniec, P.S., Mcgrath, J.A., Steel, G., Nestadt, G., Liang, K.Y., Haganir, R.L., Valle, D., Pulver, A.E., 2005. Bipolar I disorder and schizophrenia: a 440-single-nucleotide polymorphism screen of 64 candidate genes among Ashkenazi Jewish case-parent trios. *Am. J. Hum. Genet.* 77, 918–936.

Faraone, S.V., Biederman, J., Wozniak, J., 2012. Examining the comorbidity between attention deficit hyperactivity disorder and bipolar I disorder: a meta-analysis of family genetic studies. *Am. J. Psychiatry* 169, 1256–1266.

Gu, Z., Liu, W., Wei, J., Yan, Z., 2012. Regulation of N-methyl-D-aspartic acid (NMDA) receptors by metabotropic glutamate receptor 7. *J. Biol. Chem.* 287, 10265–10275.

Jun, C., Choi, Y., Lim, S.M., Bae, S., Hong, Y.S., Kim, J.E., Lyoo, I.K., 2014. Disturbance of the glutamatergic system in mood disorders. *Exp. Neurobiol.* 23, 28–35.

Kew, J.N., Kemp, J.A., 2005. Ionotropic and metabotropic glutamate receptor structure and pharmacology. *Psychopharmacology (Berl)* 179, 4–29.

Luk, H.M., 2016. Angelman-like syndrome: a genetic approach to diagnosis with illustrative cases. *Case Rep Genet* 2016, 9790169.

Maltezos, S., Horder, J., Coghlan, S., Skirrow, C., O'gorman, R., Lavender, T.J., Mendez, M.A., Mehta, M., Daly, E., Xenitidis, K., Paliokosta, E., Spain, D., Pitts, M., Asherson, P., Lythgoe, D.J., Barker, G.J., Murphy, D.G., 2014. Glutamate/glutamine and neuronal integrity in adults with ADHD: a proton MRS study. *Transl. Psychiatry* 4, e373.

Mcquillin, A., Bass, N., Anjorin, A., Lawrence, J., Kandaswamy, R., Lydall, G., Moran, J., Sklar, P., Purcell, S., Gurling, H., 2011. Analysis of genetic deletions and duplications in the University College London bipolar disorder case control sample. *Eur. J. Hum. Genet.* 19, 588–592.

Niu, W., Huang, X., Yu, T., Chen, S., Li, X., Wu, X., Cao, Y., Zhang, R., Bi, Y., Yang, F., Wang, L., Li, W., Xu, Y., He, L., He, G., 2017. Association study of *GRM7* polymorphisms with major depressive disorder in the Chinese Han population. *Psychiatr. Genet.* 27, 78–79.

Noroozi, R., Taheri, M., Movafagh, A., Mirfakhraie, R., Solgi, G., Sayad, A., Mazdeh, M., Darvish, H., 2016. Glutamate receptor, metabotropic 7 (*GRM7*) gene variations and susceptibility to autism: a case-control study. *Autism Res.* 9, 1161–1168.

Park, S., Jung, S.W., Kim, B.N., Cho, S.C., Shin, M.S., Kim, J.W., Yoo, H.J., Cho, D.Y., Chung, U.S., Son, J.W., Kim, H.W., 2013. Association between the *GRM7* rs3792452 polymorphism and attention deficit hyperactivity disorder in a Korean sample. *Behav. Brain Funct.* 9, 1.

Pergadia, M.L., Glowinski, A.L., Wray, N.R., Agrawal, A., Saccone, S.F., Loukola, A., Broms, U., Korhonen, T., Penninx, B.W., Grant, J.D., Nelson, E.C., Henders, A.K., Schrage, A.J., Chou, Y.L., Keskitalo-Vuokko, K., Zhu, Q., Gordon, S.D., Vink, J.M., De Geus, E.J., Macgregor, S., Liu, J.Z., Willemsen, G., Medland, S.E., Boomsma, D.I., Montgomery, G.W., Rice, J.P., Goate, A.M., Heath, A.C., Kaprio, J., Martin, N.G., Madden, P.A., 2011. A 3p26-3p25 genetic linkage finding for DSM-IV major depression in heavy smoking families. *Am. J. Psychiatry* 168, 848–852.

Saus, E., Brunet, A., Armengol, L., Alonso, P., Crespo, J.M., Fernandez-Aranda, F., Guitart, M., Martin-Santos, R., Menchon, J.M., Navines, R., Soria, V., Torrens, M., Urrutavizcaya, M., Valles, V., Gratacos, M., Estivill, X., 2010. Comprehensive copy number variant (CNV) analysis of neuronal pathways genes in psychiatric disorders identifies rare variants within patients. *J. Psychiatr. Res.* 44, 971–978.

Shi, J., Potash, J.B., Knowles, J.A., Weissman, M. m., Coryell, W., Scheftner, W.A., Lawson, W.B., Depaulo JR., J.R., Gejman, P.V., Sanders, A.R., Johnson, J.K., Adams, P., Chaudhury, S., Jancic, D., Evgrafov, O., Zvinjatskovskiy, A., Ertman, N., Gladis, M., Neimanas, K., Goodell, M., Hale, N., Ney, N., Verma, R., Mirel, D., Holmans, P., Levinson, D.F., 2011. Genome-wide association study of recurrent early-onset major depressive disorder. *Mol. Psychiatry* 16, 193–201.

Shyn, S.I., Shi, J., Kraft, J.B., Potash, J.B., Knowles, J.A., Weissman, M.M., Garriock, H.A., Yokoyama, J.S., Mcgrath, P.J., Peters, E.J., Scheftner, W.A., Coryell, W., Lawson, W.B., Jancic, D., Gejman, P.V., Sanders, A.R., Holmans, P., Slager, S.L., Levinson, D.F., Hamilton, S.P., 2011. Novel loci for major depression identified by genome-wide association study of Sequenced Treatment Alternatives to Relieve Depression

- and meta-analysis of three studies. *Mol. Psychiatry* 16, 202–215.
- Sklar, P., Smoller, J.W., Fan, J., Ferreira, M.A., Perlis, R.H., Chambert, K., Nimgaonkar, V.L., McQueen, M.B., Faraone, S.V., Kirby, A., De Bakker, P.I., Ogdie, M.N., Thase, M.E., Sachs, G.S., Todd-Brown, K., Gabriel, S.B., Sougnez, C., Gates, C., Blumenstiel, B., Defelice, M., Ardlie, K.G., Franklin, J., Muir, W.J., Mcghee, K.A., Macintyre, D.J., Mclean, A., Vanbeck, M., Mcquillin, A., Bass, N.J., Robinson, M., Lawrence, J., Anjorin, A., Curtis, D., Scolnick, E.M., Daly, M.J., Blackwood, D.H., Gurling, H.M., Purcell, S.M., 2008. Whole-genome association study of bipolar disorder. *Mol. Psychiatry* 13, 558–569.
- Sullivan, P.F., De Geus, E.J., Willemsen, G., James, M.R., Smit, J.H., Zandbelt, T., Arolt, V., Baune, B.T., Blackwood, D., Cichon, S., Coventry, W.L., Domschke, K., Farmer, A., Fava, M., Gordon, S.D., He, Q., Heath, A.C., Heutink, P., Holsboer, F., Hoogendijk, W.J., Hottenga, J.J., Hu, Y., Kohli, M., Lin, D., Lucae, S., Macintyre, D.J., Maier, W., Mcghee, K.A., McGuffin, P., Montgomery, G.W., Muir, W.J., Nolen, W.A., Nothen, M.M., Perlis, R.H., Piro, K., Posthuma, D., Rietschel, M., Rizzu, P., Schosser, A., Smit, A.B., Smoller, J.W., Tzeng, J.Y., Van Dyck, R., Verhage, M., Zitman, F.G., Martin, N.G., Wray, N.R., Boomsma, D.I., Penninx, B.W., 2009. Genome-wide association for major depressive disorder: a possible role for the presynaptic protein piccolo. *Mol. Psychiatry* 14, 359–375.
- Tang, B., Thornton-Wells, T., Askland, K.D., 2011. Comparative linkage meta-analysis reveals regionally-distinct, disparate genetic architectures: application to bipolar disorder and schizophrenia. *PLoS One* 6 e19073.
- The Wellcome Trust Case Control, C., Burton, P.R., Clayton, D.G., Cardon, L.R., Craddock, N., Deloukas, P., Duncanson, A., Kwiatkowski, D.P., McCarthy, M.I., Ouwehand, W.H., Samani, N.J., Todd, J.A., Donnelly, P., Barrett, J.C., Burton, P.R., Davison, D., Donnelly, P., Easton, D., Evans, D., Leung, H.-T., Marchini, J.L., Morris, A.P., Spencer, C.C.A., Tobin, M.D., Cardon, L.R., Clayton, D.G., Attwood, A.P., Boorman, J.P., Cant, B., Everson, U., Hussey, J.M., Jolley, J.D., Knight, A.S., Koch, K., Meech, E., Nutland, S., Prowse, C.V., Stevens, H.E., Taylor, N.C., Walters, G.R., Walker, N.M., Watkins, N.A., Winzer, T., Todd, J.A., Ouwehand, W.H., Jones, R.W., Mcardle, W.L., Ring, S.M., Strachan, D.P., Pembrey, M., Breen, G., St Clair, D., Caesar, S., Gordon-Smith, K., Jones, L., Fraser, C., Green, E.K., Grozeva, D., Hamshere, M.L., Holmans, P.A., Jones, I.R., Kirov, G., Moskvina, V., Nikolov, I., O'donovan, M.C., Owen, M.J., Craddock, N., Collier, D.A., Elkin, A., Farmer, A., Williamson, R., McGuffin, P., Young, A.H., Ferrier, I.N., Ball, S.G., Balmforth, A.J., Barrett, J.H., Bishop, D.T., Iles, M.M., Maqbool, A., Yuldasheva, N., Hall, A.S., Braund, P.S., Burton, P.R., Dixon, R.J., Mangino, M., Stevens, S., Tobin, M.D., Thompson, J.R., Samani, N.J., Bredin, F., Tremelling, M., Parkes, M., Drummond, H., Lees, C.W., Nimmo, E.R., Satsangi, J., Fisher, S.A., Forbes, A., Lewis, C.M., et al., 2007. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* 447, 661.
- Thompson, J.A., Ziman, M., 2011. Pax genes during neural development and their potential role in neuroregeneration. *Progress in Neurobiology* 95, 334–351.
- Ward, L.D., Kellis, M., 2012. HaploReg: a resource for exploring chromatin states, conservation, and regulatory motif alterations within sets of genetically linked variants. *Nucleic Acids Res.* 40, D930–D934.
- Warton, F.L., Howells, F.M., Russell, V.A., 2009. Increased glutamate-stimulated release of dopamine in substantia nigra of a rat model for attention-deficit/hyperactivity disorder—lack of effect of methylphenidate. *Metab. Brain Dis.* 24, 599–613.
- Yang, Y., Pan, C.H., 2013. Role of metabotropic glutamate receptor 7 in autism spectrum disorders: a pilot study. *Life Sci.* 92, 149–153.
- Yoo, J., Lee, Y., Kim, Y., Rha, S.Y., Kim, Y., 2008. SNPAnalyzer 2.0: a web-based integrated workbench for linkage disequilibrium analysis and association analysis. *BMC Bioinf.* 9, 290.
- Yuksel, C., Ongur, D., 2010. Magnetic resonance spectroscopy studies of glutamate-related abnormalities in mood disorders. *Biol. Psychiatry* 68, 785–794.
- Zhou, Y., Danbolt, N.C., 2014. Glutamate as a neurotransmitter in the healthy brain. *J. Neural Transm.* 121, 799–817.