



## Neuromuscular activity during stair descent in ACL reconstructed patients: A pilot study<sup>☆</sup>

Aglaja Busch<sup>a,b,\*</sup>, Angela Blasimann<sup>a</sup>, Philipp Henle<sup>c</sup>, Heiner Baur<sup>a</sup>

<sup>a</sup> Department Health & Physiotherapy, Bern University of Applied Sciences, Murtenstrasse 10, CH-3008 Bern, Switzerland

<sup>b</sup> University Outpatient Clinic, Sports Medicine & Sports Orthopedics, University of Potsdam, Am Neuen Palais 10, DE-14469 Potsdam, Germany

<sup>c</sup> Department of Knee Surgery and Sports Traumatology, Sonnenhof Orthopaedic Center, Buchserstrasse 30, CH-3006 Bern, Switzerland

### ARTICLE INFO

#### Article history:

Received 13 July 2018

Received in revised form 20 November 2018

Accepted 21 December 2018

#### Keywords:

Knee

Neuromuscular control

Sensorimotor control

Electromyography

Rehabilitation

### ABSTRACT

**Background:** The anterior cruciate ligament (ACL) rupture is a severe knee injury. Altered kinematics and kinetics in ACL reconstructed (ACL-R) patients compared to healthy participants (ACL-I) are known and attributed to an altered sensorimotor control. However, studies on neuromuscular control often lack homogeneous patient cohorts. The objective was to examine neuromuscular activity during stair descent in patients one year after ACL reconstruction.

**Method:** Neuromuscular activity of vastus medialis (VM) and lateralis (VL), biceps femoris (BF) and semitendinosus (ST) was recorded by electromyography in 10 ACL-R (age:  $26 \pm 10$  years; height:  $175 \pm 6$  cm; mass:  $75 \pm 14$  kg) and 10 healthy matched controls (age:  $31 \pm 7$  years; height:  $175 \pm 7$  cm; mass:  $68 \pm 10$  kg). A 10-minute walking treadmill warm-up was used for submaximal normalization. Afterwards participants descended 10 times a six-step stairway at a self-selected speed. The movement was separated into pre-activation (PRE), weight acceptance (WA) and push-off phase (PO). Normalized root mean squares for each muscle, limb and movement phase were calculated. Kruskal-Wallis ANOVA compared ACL-R injured and contralateral leg and the ACL-I leg ( $\alpha = 0.05$ ).

**Results:** Significant increased normalised activity in ST during WA in ACL-R injured leg compared to ACL-I and during PO in VL in the ACL-R contralateral leg compared to ACL-I. Decreased activity was shown in VM in ACL-R injured compared to contralateral leg ( $p < 0.05$ ).

**Conclusion:** Altered neuromuscular activations are present one year after ACL reconstruction compared to the contralateral and healthy matched control limb. Current standard rehabilitation programs may not be able to fully restore sensorimotor control and demand further investigations.

© 2018 Elsevier B.V. All rights reserved.

## 1. Introduction

The anterior cruciate ligament (ACL) rupture is a common sports injury which is followed by short-term and long-term effects such as pain, limited mobility, reduced sports participation and finally joint degeneration. To prevent an early onset of degener-

<sup>☆</sup> **Acknowledgement:** The study is supported by the Swiss National Science Foundation SNSF (project number: 176060, <http://p3.snf.ch/project-176060>).

\* Corresponding author at: University Outpatient Clinic, University of Potsdam, Am Neuen Palais 10, D-14469 Potsdam, Germany.

E-mail addresses: [agbusch@uni-potsdam.de](mailto:agbusch@uni-potsdam.de), (A. Busch), [angela.blasimann@bfh.ch](mailto:angela.blasimann@bfh.ch), (A. Blasimann), [philipphenle@sonnenhof.ch](mailto:philipphenle@sonnenhof.ch), (P. Henle), [heiner.baur@bfh.ch](mailto:heiner.baur@bfh.ch), (H. Baur).

ative knee osteoarthritis the surgical reconstruction is still very often the treatment of choice [1], despite similar outcomes for conservatively or surgically treated ACL patients [2,3].

It is already stated in many different studies that kinematic and kinetic analyses show differences regarding movement patterns during walking, running or jumping in participants with ACL deficiency or reconstruction still years after rehabilitation [4–6]. These differences can also be seen in normal activities of daily living, as a systematic review and meta-analysis found some evidence that patients with reconstructed ACL (ACL-R) had lower external knee flexion moments during stair descent compared to controls and the contralateral leg. Furthermore, trends show that the ACL reconstructed knee had lower varus moments compared to the contralateral limb but not compared to controls [7]. This indicates that, after ACL reconstruction and rehabilitation, the movement pattern is not fully restored. Several articles and reviews reporting high reinjury rates after reconstruction and return to sport also confirmed this fact which points out a possible lack in the surgical or conservative treatment of ACL injuries, rehabilitation scheme itself or insufficient compliance to it [8,9].

It is assumed that the loss of the ACL and its incorporated receptors emerges into an altered sensorimotor control [10–13]. The sensorimotor control implements the interaction between the sensory input (afferent), the processing of the information and the output (efferent) as a motor behaviour, which can be measured e.g. biomechanically [14]. Hall and colleagues [15] showed that the neuromuscular activity in ACLR patients during stair descent was significantly higher in the *M. gluteus maximus*, *medius* and *M. semimembranosus*. Moreover, these patients showed a reduced activity in the *M. rectus femoris* compared to the control group. These results underline the previously described indication of an altered neuromuscular control. Nevertheless, the research group included a broad range of patients with time after surgery from one to 18 years. This limits the interpretation of neuromuscular deficits towards a specific time point of for example return to full sport participation.

To the authors' knowledge Hall et al. [15] published the first study investigating the neuromuscular activity during stair walking in ACLR patients. Therefore, the aim of this cross-sectional pilot study was to examine the neuromuscular activity in ACLR participants one-year post-surgery. The hypothesis was that the ACLR participants show lower quadriceps and higher hamstring activity compared to healthy controls. Furthermore, it is expected that one-year post-surgery no differences are present between the injured and contralateral sides. The results should contribute to the knowledge about specific neuromuscular consequences after ACL rupture and reconstruction during activities of daily living and should lead to more insights into the neuromuscular deficits after an ACL injury.

## 2. Material and method

### 2.1. Participants

In total, 20 participants were included in this study. Ten participants (ACL-R) suffered one year ( $13.6 \pm 2$  months) prior to the study from an ACL rupture. All patients were treated with a quadriceps tendon graft by the same surgeon. The participants had additional injuries which needed surgery, such as suture of the meniscus ( $N = 5$ ), partial tear of the medial collateral ligament ( $N = 3$ ), partial meniscectomy ( $N = 1$ ) and one had no additional injuries.

Ten healthy participants (ACL-I) without prior injury of the knee were selected as controls. They were matched according to age, height, weight, gender, (sports) activity level and leg dominance. General inclusion criteria were: Age between 18 and 65 years, physically active (Tegner Activity Score (TAS) = min. 4) [16] and ACL status intact or reconstructed. Exclusion criteria for both groups were: cardiac, neurological, peripheral and vascular diseases, musculoskeletal disorders, acute infection, acute pain, effusion, other acute lower limb/trunk injuries, thrombosis, alcohol abuse and pregnancy.

All participants complete a questionnaire regarding their quantity of weekly sports participation (sports discipline and time spent), the Knee injury and Osteoarthritis Outcome Score (KOOS) with a subscale asking for pain, other symptoms, function in daily living, function in sport and recreation and knee related quality of life [17], the TAS with a score range from 0 (sick leave or disability pension) to 10 (competitive sport on professional level) [16] and a visual analogue scale (VAS) [18] (0–10 cm) describing the general well-being (gwb) and acute pain. The study was ethically approved by the local legal authority.

### 2.2. Procedures

After providing informed consent and filling out the questionnaires, participants were prepared for electromyographic (EMG) measurements. The skin was shaved, roughed with sandpaper and cleaned with alcohol for optimal detection of the muscle signals. The electrodes were attached on the *M. vastus medialis* (VM) and *lateralis* (VL), *M. biceps femoris* (BF) and *M. semitendinosus* (ST) of both limbs and a reference electrode was placed on the right patella according to the recommendations of the Surface ElectroMyoGraphy for the Non-Invasive Assessment of Muscles Project (SENIAM) [19]. Bipolar electrodes (Blue Sensor®, Ambu, Denmark, Type P-00-S, inter-electrode distance: 20 mm) were used and the inter-electrode impedance was controlled and kept equal or below 2 k $\Omega$  (Impedance meter D175, Digitimer®, Hertfordshire, UK). Transmission of the signal occurred across a differential-preamplifier (gain: 500, Input Impedance: 4000 M, Common Mode Rejection 90 dB at 60 Hz) to a telemetric main amplifier (PowerPack, Pfittec®, Endingen, Germany; Band-Pass Filter: 10 Hz to 1 kHz, gain: 5.0, resultant overall gain: 2500), where it was recorded at 2000 Hz. An analogue/digital conversion was conducted (NI PCI 6255, National Instruments®, Austin, USA; 1.25 MS/s, 16 Bit), before labVIEW®-based software (Imago Record, Pfittec®, Endingen, Germany) registered the signals.

A 10-minute warm-up on a treadmill (hp cosmos®, QUASARmed, Nussdorf-Traunstein, D) at a speed of 5 km/h (1.39 m/s), preceded the measurements to prepare participants for the upcoming protocol: First they walked on the same treadmill with same speed for three minutes, whereby electromyographic signals were recorded in the last minute. The treadmill was equipped with two force transducers (Typ KMB52, 10 kN, Megatron Elektronik AG & Co. KG, Putzbrunn, Germany) underneath it to detect the initial heel strike of each gait cycle. During post-processing these recordings were used for individual submaximal EMG normalization of each subject [20].

After the warm-up and measurement for normalization, the participants performed the stair climbing task. They descended 10 times at a self-selected speed a six-step stairway. Two multi-component force plates (Type 9286BA, Kistler®, Winterthur, CH) were embedded in the third and fourth steps. They were supported by a heavy metal frame independent from the stairs. The participants had to start always with the right foot, to have the same limb on the third, respectively fourth force plate to detect the initial foot contact. The inclination of the stairway was 30.6° with a step height of 17.1 cm and step depth of 29 cm [21].

### 2.3. Data analysis

Stair descent movement cycles were divided into three movement phases: pre-activation (PRE), weight acceptance (WA) and push-off (PO). The pre-activation phase was defined from 150 ms prior to initial foot strike until foot strike. From initial foot strike until the lowest applied vertical ground reaction force lasted the weight acceptance phase (“braking phase” until anterior–posterior force crosses the zero line). The push-off phase follows until vertical ground reaction forces decline to zero (“propulsion phase”).

Electromyographic data was further processed using labVIEW®-based software (Imago Process Master, Pfittec®, Emdingen, Germany). The raw EMG signals were band-pass filtered at 10–500 Hz (Butterworth, 2nd order) and the root mean squares (RMS) for each phase and muscle were calculated and exported in an Excel spread sheet. The activation of the mean gait cycle for each muscle from normal walking was used to (submaximal) normalize the three phases during stairway walking. This secured comparability of individual results between injured and contralateral limbs as well as between injured limb and matched leg from controls.

Furthermore, the average and standard deviation of the 10 single trails in each participant were calculated and checked for plausibility. Individual values higher than twice the standard deviation were tracked back to the original data and corrected if possible.

### 2.4. Statistical analysis

The data was checked for normal distribution using the Shapiro–Wilk Test, revealing a deviation from normal distribution, however for a better comparison with other studies the data is also presented with means and standard deviations. The participant characteristics (age, height and weight) were tested for significant differences using an unpaired t-test. A Kruskal–Wallis analysis of variance (ANOVA) was conducted for the ACL-R injured and contralateral leg and ACL-I matched leg for the three different movement phases (PRE, WA & PO) and the four recorded muscles (Statistical Package for the Social Science (SPSS) software version 23.0, IBM, SPSS; Inc. Chicago, IL, USA, 2015). Additionally, a Dunn–Bonferroni post-hoc pairwise comparison was performed. The  $\alpha$ -level was set at 0.05.


## 3. Results

An unpaired t-test comparison of age, height and weight revealed no significant differences between the ACL-R and ACL-I groups ( $p > 0.05$ ). The ACL-R group showed a reduced total score of the KOOS (approx. - 10% of total KOOS,  $p = 0.001$ ) compared to the ACL-I control group. The VAS asking for pain was similar in before and after measurements and the VAS regarding the

**Table 1**  
Baseline characteristics of participants.

Characteristics	ACL-I (N = 10)	ACL-R (N = 10)
Age – years	31 ± 7	26 ± 10
Height – cm	175 ± 8	175 ± 7
Weight – kg	68 ± 10	75 ± 14
Female (%)	30	30
KOOS (total max: 168)	166 ± 2	148 ± 7
VAS – pain pre/post	0.1/0.7	1/1
VAS – gwb pre/post	0.7/0.9	2/4
Tegner score (max: 10)	6 ± 1	7 ± 2

Legend: Knee Injury and Osteoarthritis Outcome Score; Visual analogue scale; indicates values on a 0–10 cm scale; Tegner activity score (pre-injury) range from 0 (sick leave or disability pension) to 10 (competitive sport on professional level).



**Figure 1.** Normalized neuromuscular activity (in % of the submaximal voluntary contraction (subMVC)) comparing ACL-R injured and contralateral leg and ACL-I matched control leg. Displayed in a boxplot showing from top to down: outlier, upper extreme, upper quartile, median, lower quartile, lower extreme, outlier. Legend: PRE= pre-activation; WA = weight acceptance; PO = push off; VM = vastus medialis; VL = vastus lateralis; BF = biceps femoris; ST = semitendinosus.

general well-being was slightly increased in both groups after protocol completion. A Tegner score of averaged 6 in the ACL-I group indicates a higher recreational sports activity in the disciplines of e.g. soccer, basketball, tennis or jogging up to five times per week. The ACL-R participants with a Tegner score of 7 (averaged) were slightly more active, with participation in competitive sports in e.g. tennis, athletics, basketball and orientation run. Further details of the participant characteristics can be found in Table 1.

The comparison of neuromuscular activity during stair descent of the ACL-R injured and contralateral leg and the ACL-I matched control leg revealed significant differences of the groups in three conditions. During the WA in the semitendinosus muscle ( $p = 0.031$ ) (Figure 1), post-hoc analysis resulting in a significant higher activity of 61% in the ACL-R injured leg compared to the ACL-I matched leg ( $p = 0.047$ ) (see Table 2). Furthermore, statistically significant differences were also observed in the vastus medialis and vastus lateralis muscles during the PO phase ( $p = 0.007$ ,  $p = 0.008$ , respectively) (Figure 1). The post-hoc analysis showed for the vastus medialis significantly reduced activities of about 70% in the ACL-R injured leg compared to the contralateral

**Table 2**

Normalized neuromuscular activity (mean and standard deviation in % of the submaximal voluntary contraction (subMVC)) of the ACL-R injured and contralateral leg and ACL-I matched control leg.

Phase	Muscle	ACL-R injured (subMVC)	ACL-R contralateral (subMVC)	ACL-I matched (subMVC)	p-value
Pre-activation	VM	128 ± 70	152 ± 76	125 ± 36	0.89
	VL	150 ± 62	135 ± 72	122 ± 44	0.5
	BF	154 ± 122	121 ± 50	149 ± 57	0.55
	ST	242 ± 182	147 ± 89	148 ± 61	0.54
Weight acceptance	VM	275 ± 95	256 ± 131	270 ± 138	0.81
	VL	290 ± 187	186 ± 31	215 ± 82	0.2
	BF	70 ± 46	113 ± 57	64 ± 34	0.09
	ST	158 ± 110	120 ± 60	61 ± 55	0.03
Push off	VM	109 ± 39	359 ± 213	158 ± 97	0.007
	VL	199 ± 130	343 ± 160	127 ± 72	0.008
	BF	149 ± 111	71 ± 37	89 ± 34	0.17
	ST	191 ± 146	120 ± 88	88 ± 65	0.09

Legend: ACL-R = ACL reconstructed group; ACL-I = ACL matched control group; VM = vastus medialis; VL = vastus lateralis; BF = biceps femoris; ST = semitendinosus.

leg ( $p = 0.008$ ) (Table 2). The vastus lateralis had significantly higher neuromuscular activation (approx. +63%) comparing the ACL-R contralateral leg and the ACL-I matched leg ( $p = 0.006$ ) (Table 2). Additional details of the mean and standard deviation of the neuromuscular activity of each muscle in the three movement phases can be found in Table 2.

#### 4. Discussion

The aim of this study was to investigate the neuromuscular activity during stair descent in patients one-year after ACL reconstruction. Results showed differences of the muscular activation comparing the ACL-R injured and contralateral leg and the ACL-I matched leg during the three movement phases. Especially, the higher activities of the ST during WA in the ACL-R injured leg and the reduced neuromuscular activities of the VM of the ACL-R injured limb and higher activation of the VL of the ACL-R contralateral leg in the PO phase differed significantly. These results partly confirm the hypothesis of lower quadriceps and higher hamstring activity in the ACL-R participants compared to the matched control leg.

The participant characteristics show that the ACL-R participants are still functionally affected by the ACL injury, but can complete the movement tasks without greater decline or increase of the general well-being or pain, respectively. Furthermore, the Tegner score of averaged 7 indicates a return to high physical activity, also to competitive sports. These findings are comparable to a systematic review and meta-analysis revealing that the majority of ACL patients following ACL reconstruction return to some kind of sport, 63% returned to their preinjury level and less than the half returns to competitive sport [22].

The recorded higher neuromuscular activation of the hamstrings during the weight acceptance can also be found in the study by Hall [15]. A significant higher activity of the semimembranosus muscle was found during one to 50% of stance phase during stair descent compared to the control group. In contrast, a study focusing on the kinematics during stair ambulation comparing the ACL-R injured to the contralateral limb found significantly lower peak knee flexion moments from weight acceptance until push off during stair descent in the leg with ACL reconstruction surgery [23]. Taking the primary purpose of the ACL – to prevent an extensive anterior tibia translation during extension – into account, higher activities of the hamstring muscles point to a supportive adaptation of the synergistic muscles in favour of the reconstructed ACL [24]. Thus, rehabilitation might not only focus on the recovery of quadriceps strength (knee extension) but also on the strengthening and neuromuscular control of the hamstrings.

Findings of significant reduction of the muscular activity in the VM during push off are also indicated in other studies [15,23]. Concerning the injury mechanism of the ACL rupture with extreme knee valgus, this reduction might be a preventive act of adaptation against intense stress on the reconstructed ACL during movement. Furthermore, a reduced activity of the quadriceps might be a consequence of the surgical treatment with the quadriceps tendon graft. A study examining balance performance, isokinetic strength, squat jumps and stiffness in participants having a surgical repair of the ruptured ACL with a bone-patella-bone tendon graft or semitendinosus/gracilis autograft, showed only significant differences in squat jump height comparing both groups one year after surgery [25]. However, insights into neuromuscular adaptations of the respective surgical techniques would complete the knowledge about the particular consequences.

The significantly higher activation of the VL and greater activation of the VM during PO in the contralateral leg compared to controls indicates a compensative strategy to ensure adequate function during movement [23]. Nonetheless, the higher activation might lead to an early onset of fatigue of the joint stabilizing muscles, which could explain high ACL rupture rates of the contralateral leg after the initial rupture [8].

Overall, the presented results prove and are in line with further conclusions that also years after surgery and complete rehabilitation process movement strategies and neuromuscular activation patterns are still impaired in both limbs [26]. This can also be seen in the high reinjury rates after ACL rupture and reconstruction [2]. However, causes of reinjury rates are still not clearly identified. Different consequences in both limbs, as shown in this pilot study, do not support the general process of often used

return to sport (RTS) criteria using limb symmetry indexes. A recent case control study supports these assumptions with bilateral deficits in hop jumps comparing ACL reconstructed and healthy participants [27]. Furthermore, trends of higher hamstring activity during WA could lead to new orientation shifting the focus from pure quadriceps strength, representing a general aspect of physical function, to a more differentiated approach by analysing neuromuscular activity in physical function [28]. This may complement a holistic approach towards ACL injury needs integrating strength and neuromuscular control of the hamstring to support the ACL graft. Additionally, a systematic review showed very inconsistent RTS strength criteria and isokinetic protocols to evaluate muscular strength [29]. Time until RTS is considered an important factor which is still discussed and investigated [30]. Thus, RTS guidelines are currently under revision pointing towards a multifactorial and optimized criterion-based RTS approach with a clear demand to include sensorimotor based criteria [31].

The demand to include sensorimotor based criteria is supported by the findings of the present pilot study and speaks in favour of an integrated rehabilitation process. A recent systematic review and multidisciplinary consensus points out the need of neuromuscular training to complete rehabilitation besides range of motion and muscular strength recovery [9]. Nevertheless, the neuromuscular approach might not be fully sufficient. First studies shift their attention from the outcomes of sensorimotor control (e.g. neuromuscular activity or biomechanical output) to the processing level (brain activity). Literature shows alteration in the cortical activation patterns, also years after ACL reconstruction and rehabilitation [32,33]. The ability of the brain to adapt morphologically and functionally to permanent changes is described as plasticity and provides a perspective to further explain the consequences of an ACL injury. Therefore, an ACL injury might not only be considered as a musculoskeletal but also be considered as a neurophysiological injury [34] and therefore more knowledge on central and peripheral adaptations is needed.

The study design with the integrated force plates underneath the stairway contributes to a standardised and precise analysis of initial contact and subsequent EMG data; however, there are some limitations. First, the participants could descend the stairs at a self-selected speed. This method can also be found in other studies [23,24] but might have a decisive impact on the recorded EMG activities during movement. Nonetheless, a standardised speed, for example with a metronome indicating when to take the next step, could highly modify the normal gait pattern of the participants. Furthermore, the submaximal normalization needs to be considered when comparing the results with other studies. Moreover, it is critically discussed whether the maximal voluntary isometric contraction (MVIC) measurement would have been the better option for EMG normalization. Literature has shown, that without training, the MVIC is reduced by 20–40% compared to a trained MVIC. Also, the static contraction could lead to different results from the values recorded during movement, where the muscles move relatively to the skin [35]. Moreover, the used study protocol is in line with a previous conducted study measuring neuromuscular activity in participants with an acute ACL rupture and is therefore relevant for further comparisons [36]. The statistical analysis with the Kruskal–Wallis ANOVA was conducted to compare the muscles and movement phases of the groups. However, due to the small number of participants the statistically significant results need to be handled with caution. Nevertheless, the percent differences between conditions were quite substantial reaching sufficient clinical relevance. A further gender specific analysis of EMG data is warranted as soon as sufficient sample sizes per gender are measured, as studies on differences in kinematics and neuromuscular activity during single leg squats in healthy young participants indicate [37,38]. This is also highlighted by reported gender specific injury risk [39].

## 5. Conclusion

The neuromuscular activity comparing the injured and contralateral legs of the ACL reconstructed group to a healthy matched control group differs during stair descent one-year after surgery. More participants are needed to draw relevant conclusions but the pilot results indicate the special importance of neuromuscular control mechanisms to further enhance rehabilitation and return to sports decisions. Analysis of neuromuscular activity by electromyography may serve as a valuable quality control tool to monitor this process.

## Declaration of interest

None.

## References

- [1] Thomas AC, Villwock M, Wojtys EM, Palmieri-Smith RM. Lower extremity muscle strength after anterior cruciate ligament injury and reconstruction. *J Athl Train* 2013;48:610–20. <https://doi.org/10.4085/1062-6050-48.3.23>.
- [2] Paterno MV, Rauh MJ, Schmitt LC, Ford KR, Hewett TE. Incidence of contralateral and ipsilateral anterior cruciate ligament (ACL) injury after primary ACL reconstruction and return to sport. *Clin J Sport Med* 2012;22:116–21. <https://doi.org/10.1097/JSM.0b013e318246ef9e>.
- [3] Meunier A, Odensten M, Good L. Long-term results after primary repair or non-surgical treatment of anterior cruciate ligament rupture: a randomized study with a 15-year followup. *Scand J Med Sci Sports* 2007;17:230–7. <https://doi.org/10.1111/j.1600-0838.2006.00547.x>.
- [4] Nyland J, Klain S, Caborn D. Lower extremity compensatory neuromuscular and biomechanical adaptations 2 to 11 years after anterior cruciate ligament reconstruction. *Arthroscopy* 2010;26:1212–25. <https://doi.org/10.1016/j.arthro.2010.01.003>.
- [5] Zhao C, Lin C, Wang W, Zeng C, Fang H, Pan J, et al. Kinematics of anterior cruciate ligament-deficient knees in a Chinese population during stair ascent. *J Orthop Surg Res* 2016;11:89. <https://doi.org/10.1186/s13018-016-0423>.
- [6] Markström JL, Tengman E, Häger CK. ACL-reconstructed and ACL-deficient individuals show differentiated trunk, hip, and knee kinematics during vertical hops more than 20 years post-injury. *Knee Surg Sports Traumatol Arthrosc* 2017. <https://doi.org/10.1007/s00167-017-4528-4>.
- [7] Hajizadeh M, Oskouei A, Ghalich F, Sole G. Knee kinematics and joint moments during stair negotiation in participants with anterior cruciate ligament deficiency and reconstruction: a systematic review and meta-analysis. *PM R* 2016;8. <https://doi.org/10.1016/j.pmrj.2016.01.014>.

- [8] Wiggins AJ, Grandhi RK, Schneider DK, Stanfield D, Webster KE, Myer GD. Risk of secondary injury in younger athletes after anterior cruciate ligament reconstruction: a systematic review and meta-analysis. *Am J Sports Med* 2016;44:1861–76. <https://doi.org/10.1177/0363546515621554>.
- [9] van Melick N, van Cingel REH, Brooijmans F, Neeter C, van Tienen T, Hullegie W, Nijhuis-van de Sanden MWG. Evidence-based clinical practice update: practice guidelines for anterior cruciate ligament rehabilitation based on a systematic review and multidisciplinary consensus. *Br J Sports Med* 2016;1506–15. <https://doi.org/10.1136/bjsports-2015-095898>.
- [10] Engelhardt M, Freiwald J, Reuter I, Mortier J, Huth D. Beeinflussung der Sportfähigkeit durch neuromuskuläre Veränderungen nach Trauma und Operation am Kniegelenk. *Art Ther* 2000;13:302–6.
- [11] Ageberg E. Consequences of a ligament injury on neuromuscular function and relevance to rehabilitation – using the anterior cruciate ligament-injured knee as model. *J Electromyogr Kinesiol* 2002;12:205–12.
- [12] Barrack R, Munn B. Effects of knee ligament injury and reconstruction on proprioception. *Proprioception Neuromuscul. Control Jt. Stab.*; 2000.
- [13] Relph N, Herrington L, Tyson S. The effects of ACL injury on knee proprioception: a meta-analysis. *Physiotherapy* 2014;100:187–95. <https://doi.org/10.1016/j.physio.2013.11.002>.
- [14] Baumeister J. What the brain can tell us in musculoskeletal rehabilitation. *J Sports Med Doping Stud* 2012;2:2–3. <https://doi.org/10.4172/2161-0673.1000e106>.
- [15] Hall M, Stevermer CA, Gillette JC. Muscle activity amplitudes and co-contraction during stair ambulation following anterior cruciate ligament reconstruction. *J Electromyogr Kinesiol* 2015;25:298–304. <https://doi.org/10.1016/j.jelekin.2015.01.007>.
- [16] Orthopaedics C, Lule YT. Rating Systems in the Evaluation of Knee Ligament Injuries; 2017. <https://doi.org/10.1097/00003086-198509000-00007>.
- [17] Roos E, Lohmander L. The Knee injury and Osteoarthritis Outcome Score (KOOS): from joint injury to osteoarthritis. *Health Qual Life Outcomes* 2003;1:64.
- [18] Freyd MT. The graphic rating scale. *J Educ Psychol* 1923;14:83–102.
- [19] Hermens H, Freriks B, Disselhorst-Klug C, Rau G. Development of recommendations for SEMD sensor and sensor placement procedures. *J Electromyogr Kinesiol* 2000;10:361–74.
- [20] Baur H, Muller S, Hirschmuller A, Cassel M, Weber J, Mayer F. Comparison in lower leg neuromuscular activity between runners with unilateral mid-portion Achilles tendinopathy and healthy individuals. *J Electromyogr Kinesiol* 2011;21:499–505. <https://doi.org/10.1016/j.jelekin.2010.11.010>.
- [21] Leitner M, Schmid S, Hilfiker R, Radlinger R. Test–retest reliability of vertical ground reaction forces during stair climbing in the elderly population. *Gait Posture* 2011;34. <https://doi.org/10.1016/j.gaitpost.2011.06.014>.
- [22] Ardern CL, Taylor NF, Feller JA, Webster KE. Return-to-sport outcomes at 2 to 7 years after anterior cruciate ligament reconstruction surgery. *Am J Sports Med* 2012;40:41–8. <https://doi.org/10.1177/0363546511422999>.
- [23] Zabala ME, Favre J, Scanlan SF, Donahue J, Andiracchi TP. Three-dimensional knee moments of ACL reconstructed and control subjects during gait, stair ascent and stair descent. *J Biomech* 2013;46:515–20. <https://doi.org/10.1016/j.jbiomech.2012.10.010.Three-Dimensional>.
- [24] Gao B, Cordova ML, Nigel N. Three-dimensional joint kinematics of ACL-deficient and ACL-reconstructed knees during stair ascent and descent. *Hum Mov Sci* 2012;31:222–35. <https://doi.org/10.1016/j.humov.2011.04.009>.
- [25] Pasquini A, Jacopetti M, Pogliacomi F, Ramazzina I, Costantino C. Neuromuscular recovery in ACL reconstruction with bone–tendon–patellar–bone and semitendinosus–gracilis autograft. 88; 2017; 62–8. <https://doi.org/10.23750/abm.v88i4>.
- [26] Ward S, Pearce AJ, Pietrosimone B, Bennell K, Clark R, Bryant AL. Neuromuscular deficits after peripheral joint injury: a neurophysiological hypothesis. *Muscle Nerve* 2015;327–32. <https://doi.org/10.1002/mus.24463>.
- [27] Zult T, Gokeler A, van Raay JJAM, Brouwer RW, Zijdewind I, Hortobágyi T. An anterior cruciate ligament injury does not affect the neuromuscular function of the non-injured leg except for dynamic balance and voluntary quadriceps activation. *Knee Surg Sports Traumatol Arthrosc* 2017;25:172–83. <https://doi.org/10.1007/s00167-016-4335-3>.
- [28] Lepley AS, Gribble PA, Thomas AC, Tevald MA, Sohn DH, Pietrosimone BG. Longitudinal evaluation of stair walking biomechanics in patients with ACL injury. *Med Sci Sports Exerc* 2015;7–15. <https://doi.org/10.1249/MSS.0000000000000741>.
- [29] Undheim MB, Cosgrave C, King E, Strike S, Marshall B, Falvey E, et al. Isokinetic muscle strength and readiness to return to sport following anterior cruciate ligament reconstruction: is there an association? A systematic review and a protocol recommendation. *Br J Sports Med* 2015;49:1305–10. <https://doi.org/10.1136/bjsports-2014-093962>.
- [30] Welling W, Benjaminse A, Seil R, Lemmink K, Zaffagnini S, Gokeler A. Low rates of patients meeting return to sport criteria 9 months after anterior cruciate ligament reconstruction: a prospective longitudinal study. *Knee Surg Sports Traumatol Arthrosc* 2018;0:1–9. <https://doi.org/10.1007/s00167-018-4916-4>.
- [31] Dingenen B, Gokeler A. Optimization of the return-to-sport paradigm after anterior cruciate ligament reconstruction: a critical step back to move forward. *Sports Med* 2017;47:1487–500. <https://doi.org/10.1007/s40279-017-0674-6>.
- [32] Baumeister J, Reinecke K, Schubert M, Weiß M. Altered electrocortical brain activity after ACL reconstruction during force control. *J Orthop Res* 2011;29:1383–9. <https://doi.org/10.1002/jor.21380>.
- [33] Grooms DR, Page SJ, Nichols-larsen DS, Chaudhari AMW, White SE, Onate JA. Neuroplasticity associated with anterior cruciate ligament reconstruction. 47; 2017; 180–9. <https://doi.org/10.2519/jospt.2017.7003>.
- [34] Kapreli E, Athanasopoulos S, Gliatis J, Papatheanasiou M, Peeters R, Strimpakos N. Anterior cruciate ligament deficiency causes brain plasticity: a functional MRI study; 2009; 2419–26. <https://doi.org/10.1177/0363546509343201>.
- [35] Hug F. Can muscle coordination be precisely studied by surface electromyography?; 21; 2011; 1–12. <https://doi.org/10.1016/j.jelekin.2010.08.009>.
- [36] Blasimant A, Busch A, Freiburghaus S, Marolf M, Boesch L, Henle P, et al. Altered muscle activity in patients with ACL tear compared to healthy controls during stair descent. *Phys Ther Sport* 2017;28:e1ee25.
- [37] Zeller BL, McCrory JL, Kibler WB, Uhl T. Differences in kinematics and women during the single-legged squat\*. *Am J Sports Med* 2003;31:449–56.
- [38] Gehring D, Melnyk M, Gollhofer A. Gender and fatigue have influence on knee joint control strategies during landing. *Clin Biomech* 2009;24:82–7. <https://doi.org/10.1016/j.clinbiomech.2008.07.005>.
- [39] Pappas GP, Vogelsohn MA, Staroswiecki E, Gold GE, Safran MR. Magnetic resonance imaging of asymptomatic knees in collegiate basketball players: the effect of one season of play. *Clin J Sport Med* 2016;26:483–9. <https://doi.org/10.1097/JSM.0000000000000283>.