



# Correction to: Clinical and radiological outcomes of unilateral biportal endoscopic lumbar interbody fusion (ULIF) compared with conventional posterior lumbar interbody fusion (PLIF): 1-year follow-up

Man-Kyu Park<sup>1</sup> · Soo-An Park<sup>2,3</sup> · Sang-Kyu Son<sup>1</sup> · Weon-Wook Park<sup>2</sup> · Seung-Hyun Choi<sup>2</sup>

Published online: 24 June 2019  
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

## Correction to: Neurosurgical Review

<https://doi.org/10.1007/s10143-019-01114-3>

The original publication of this article has incorrect presentation of one of the author names. Instead of Sangu-Kyu Son, it should have been Sang-Kyu Son.

The original article has been corrected.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---

The online version of the original article can be found at <https://doi.org/10.1007/s10143-019-01114-3>

---

✉ Soo-An Park  
[soan.park@gmail.com](mailto:soan.park@gmail.com)

<sup>1</sup> Department of Neurosurgery, Parkweonwook Hospital, Busan, South Korea

<sup>2</sup> Department of Orthopaedic Surgery, Parkweonwook Hospital, 4, Gwangan-ro, Suyeong-gu, Busan 48298, South Korea

<sup>3</sup> Department of Orthopaedic Surgery, Chung General Hospital, 76, Sujeong-ro, Sujeong-gu, Seongnam-si, Gyeonggi-do 13316, South Korea