

Alternaria alternata infection causing rhinosinusitis and orbital involvement in an immunocompetent patient

Z. Hattab¹, N. Ben Lasfar¹, M. Abid¹, F. Bellazreg¹, A. Fathallah², W. Hachfi¹ and A. Letaief¹

1) Infectious Diseases Department, CHU Farhat Hached Hospital, Sousse, Tunisia and 2) Parasitology and Mycology Department, CHU Farhat Hached Hospital, Sousse, Tunisia

Abstract

Invasive fungal rhinosinusitis is a rare infection that occurs primarily in immunocompromised patients. The fungal pathogen *Alternaria alternata* is rarely associated with rhinosinusitis. We report a case of *A. alternata* rhinosinusitis in an immunocompetent patient.

© 2019 Published by Elsevier Ltd.

Keywords: Cerebral parenchyma, fungus, mycelian filaments, sinus

Original Submission: 25 August 2018; **Revised Submission:** 14 February 2019; **Accepted:** 30 April 2019

Article published online: 16 May 2019

Corresponding author: N. Ben Lasfar, MD, CHU Farhat Hached Hospital, Sousse, Tunisia..

E-mail: benlasfar.nadia@yahoo.fr

Introduction

Alternaria is an ubiquitous fungus that is considered to be a nonpathogenic contaminant of the clinical specimen unless isolated by repeated culture and correlated with clinical findings. However, it is a rare cause of human infection, especially in immunocompromised patients but even more rarely in healthy hosts [1].

We report a case of *Alternaria alternata* in a healthy patient. Cutaneous infections are the most frequent diseases caused by *Alternaria*. However, infections in other organs, such as facial osteomyelitis [2], sinusitis [3] and keratomycosis [4], have also been reported. Initially patients may report nonspecific symptoms similar to chronic rhinosinusitis. Complications related to the orbital extension include preseptal cellulitis and orbital cellulitis. The lesion grows contiguously, possibly invading the central nervous system [5].

Case report

A 19-year-old man with allergic rhinitis, with no history of corticoid therapy or nasal drug use, sought care for haemicranial headache with purulent rhinorrhea, anosmia, right exophthalmia, and vision impairment without fever, which had evolved for 9 months. His neurologic examination revealed nothing abnormal. Ocular motility was normal. Examination of the fundus found a papillary edema. Computed tomographic scan revealed obliteration of the maxillary and frontal sinuses with destruction of the orbital wall and intraorbital extension (Fig. 1). Histologic examination of sinus biopsy material showed an inflammatory process of the nasal mucosa chorion with periodic acid–Schiff mycelian filaments in the mucus and fibrin leucocyte material.

Culture of the biopsy sample identified *A. alternata*. Anti-fungal sensitivity could not be assessed. The bronchial tree at fibroscopy was normal; the mycelian filaments were not found in alveolar lavage fluid. The patient underwent debridement, septoplasty and sphenoidotomy and was treated with amphotericin B 37 mg/kg per day for 15 days, followed by oral itraconazole 200 mg per day for 2 months. HIV serology was negative. Immunoglobulin assay revealed no deficit. The patient




FIG. 1. Coronal-view preoperative computed tomographic scan revealing intracranial and intraocular extension.

underwent septoplasty, sphenoidectomy, polypectomy and meatotomy. There were no complications at 1 year.

Conclusion

Our patient presented rhinosinusitis due to *Alternaria* involving the cerebral parenchyma and had no apparent predisposing factor outside of a history of allergic rhinosinusitis, which is considered to be a risk factor for fungal sinusitis [6].

The confirmation of diagnosis is based on histopathologic examination of a sample with periodic acid–Schiff staining. The histologic appearance of *Alternaria* is characterized by a dense accumulation of hyphae, infiltration of surrounding tissue, and occasional branching and chains of conidia. The treatment of *Alternaria* and appropriate antifungal therapy doses are not standardized in the literature. The optimal treatment should

consist of broad surgical debridement of involved tissues with prolonged systemic antifungal medication. *Alternaria* is sensitive to amphotericin B and miconazole [7]. Therefore, itraconazole was cited in the literature as the treatment of choice in *Alternaria* infections based on its minimum inhibitory concentration values.

Alternaria is a cause of chronic fungal sinusitis. Histopathologic examination and microbiologic identification can help in diagnosis.

Conflict of interest

None declared.

References

- [1] Loveless MO, Winn RE, Campbell M, Jones SR. Mixed invasive infection with *Alternaria* species and *Curvularia* species. *Am J Clin Pathol* 1981;76:491–3.
- [2] Chhabra V, Rastogi S, Barua M, Kumar S. *Alternaria alternata* infection associated osteomyelitis of maxilla: a rare disease entity. *Indian J Dent Res* 2013;24:639–41.
- [3] Shugar MA, Montgomery WW, Hyslop Jr NE. *Alternaria* sinusitis. *Ann Otol Rhinol Laryngol* 1981;90:251–4.
- [4] Zahra LV, Mallia D, Hardie JG, Bezzina A, Fenech T. Keratomycosis due to *Alternaria alternata* in a diabetic patient [case report]. *Mycoses* 2002;45:512–4.
- [5] Ingley AP, Parikh SL, DelGaudio JM. Orbital and cranial nerve presentations and sequelae are hallmarks of invasive fungal sinusitis caused by *Mucor* in contrast to *Aspergillus*. *Am J Rhinol* 2008;22:155–8.
- [6] Kliimek L, Bardenhewer C, Spielhaupter M, Harai C, Becker K, Pfaar O. [Local allergic rhinitis to *Alternaria alternata*: evidence for local IgE production exclusively in the nasal mucosa]. *HNO* 2015;63:364–72.
- [7] Valera FCP, de Lago T, Tamashiro E, Yassuda CC, Silveira F, Anselmo-Lima WT. Prognosis of acute invasive fungal rhinosinusitis related to underlying disease. *Int J Infect Dis* 2011;15:e841–4.