



Applied nutritional investigation

Breast milk: To each his own. From metabolomic study, evidence of personalized nutrition in preterm infants



Serafina Perrone M.D., Ph.D. ^{a,*}, Mariangela Longini Biologist ^a, Ilaria Zollino M.D. ^a,
 Francesco Bazzini Biologist ^a, Maria Tassini Biologist ^b, Antonio Vivi Biologist ^b, Carlotta Bracciali M.D. ^a,
 Marco Calderisi Ph.D. ^c, Giuseppe Buonocore M.D. ^a

^a Department of Molecular and Developmental Medicine, University of Siena, Siena, Italy

^b NMR Center, University of Siena, Siena, Italy

^c Kode Solutions, Pisa, Italy

ARTICLE INFO

Article History:

Received 24 August 2018

Received in revised form 21 November 2018

Accepted 20 December 2018

Keywords:

Human milk

Metabolomics

Preterm

NMR spectroscopy

Nutrition

ABSTRACT

Objectives: The composition of milk from mothers delivering prematurely differs from that of full-term mature milk and changes over time. The aim of this study is to test the hypothesis that changes in milk metabolomic profile from mothers delivering prematurely persist over time when compared with mothers delivering at term.

Methods: Nuclear magnetic resonance spectroscopy was used to analyze the metabolome pattern of human milk samples collected from 18 mothers. Twelve mothers collected 12 term milk samples (one for each mother) once between 4 and 7 d after delivery. Six mothers delivering prematurely (29–31 wk of gestational age) and collected three samples each, once a week after delivery until the third week after birth.

Results: Principal component analysis identified two distinct metabolite groups, one represented by the 18 preterm milk samples and the other by term milk samples. Metabolite profiling identified that lactose and oligosaccharide levels were significantly more represented in preterm than in milk term samples.

Conclusions: The preterm milk metabolome pattern undergoes maturation during the first 3 wk after birth, but at the end of the third week it still does not resemble the term milk pattern. The specific changes in mothers' milk metabolomic profiles according to their offspring might reflect the different nutritional requirement of each preterm infant. This knowledge is crucial to move from standardized nutritional protocols to tailored, individualized nutrition in preterm infants.

© 2019 Elsevier Inc. All rights reserved.

Introduction

Human milk is the best nourishment for the healthy growth and development of infants [1,2]. Milk is species specific and has a dynamic composition in terms of nutritive components and non-nutritive bioactive factors [3] that makes it uniquely suited for feeding infants [4]. The benefits of breastfeeding on children's health are numerous, but, according to European Society for Pediatric Gastroenterology Hepatology and Nutrition, the most documented effect is the reduced risk of infectious diarrhea and acute otitis media [2]. Besides these advantages, some studies have found

that breastfed infants have better short- and long-term outcomes compared with those of formula-fed infants [5,6]. Formula-fed infants have different growth patterns, nutritional status, and gut microbiota that may be associated with higher risk of obesity, diabetes, and cardiovascular diseases later in life. Even with all the current attempts to modify the composition of infant formulas to make them more similar to human milk, huge differences persist between these two kinds of milk [7–9].

Just as in the case of term infants, the first choice of nourishment for preterm newborns should be the mother's own milk, and when this is not available, either at all or in sufficient quantity, donor milk from lactating mothers represents the next best choice [10].

Recently we reported that there is a clear difference in the composition of milk collected from mothers delivering extremely preterm (23–25 wk of gestational age [GA]) compared with term births. Furthermore, we found a common temporal variation in the

This work was partially funded by the Europe Against Infant Brain Injury (EUR-AIBI) foundation.

* Corresponding author. Tel.: +39 0577 586542.

E-mail address: Saraspv@yahoo.it (S. Perrone).

first 3 wk of lactation of preterm deliveries. The milks tend to have very similar properties to that collected around the 30th week of postnatal age [8]. These data suggested that, after 3 wk of lactation, the milk composition seems to be more homogeneous as its “maturation” reaches completion.

The present study aims to evaluate whether the milk of mothers who delivered at a mean GA of 30 (more mature milk) had a similar composition to that collected from mothers with term delivery. In particular, we aimed to assess whether and when the metabolic profiles of milk samples from mothers of premature infants resembled the metabolic profiles of term mature milk, sampling weekly until the third week of postnatal age.

Materials and methods

Using nuclear magnetic resonance (NMR) spectroscopy, we analyzed 30 milk samples, obtained from 18 mothers delivering between the 29th and 41st wk of GA and collected directly from the milk bank of Santa Maria alle Scotte Hospital in Siena, Italy, from January 2013 to August 2015. Written informed consent was obtained from all the enrolled mothers.

Mothers were Caucasian, healthy, and had a similar home environment. They did not smoke or consume alcohol or drugs. All mothers had a similar lifestyle and followed a varied and balanced diet. Milk samples were collected from 18 mothers. Twelve mothers collected 12 term milk samples (one for each mother) once between 4 to 7 d after delivery. Six mothers who delivered prematurely (from 29th to 31st wk of GA) collected three samples each, once a week after delivery until the third week after birth. Milk samples were collected from mothers in the morning, before breast-feeding. Then, 1 mL of each milk sample was processed and stored in the milk bank until completing the recruitment. All milk samples were analyzed at the same time (Table 1).

Butylated hydroxytoluene (BHT) was added to each sample for its antioxidant properties [11], and the samples were frozen and stored at -80°C until the analysis was performed. Before analysis, we extracted and separated the water-soluble metabolites from fat-soluble ones. In each sample of milk, an 8 mL solution composed of chloroform, methanol, and water was added (1:1:0.3, respectively). Then these new samples were placed in glass beaker at 0°C for 15 min and subsequently centrifuged at a speed of 4000 rpm at 4°C for 20 min to separate the methanol and water phase from the chloroform phase. Finally, the methanol and water phase was transferred into small beakers and frozen, ready for the NMR analysis.

For NMR experiments, samples were lyophilized to eliminate water and methanol, then treated with deuterated water and trimethylsilyl-2,2,3,3-tetra-deuterio-propionic acid 20 mM to the calibration of the chemical shift at 0 ppm. A 600- μL sample was then transferred to NMR tubes and the pH was adjusted to 6 ± 0.02 for a first measurement. A second measurement was then carried out at a pH 2.50 ± 0.02 because the chemical shift of ionizable substances is highly pH dependent and, at a pH 2.50, all chemical shift values have a reproducibility of ± 0.01 ppm. NMR measurements were performed with a Bruker Avance DRX 600 MHz using a selective enhance inverse probe.

The spectra were acquired at a constant temperature of 298.0 ± 0.1 K using a 90° pulse. A delay of 10 s in the sequence was included to allow the complete relaxation of the spin system. We applied a line-broadening function of 0.3 Hz before Fourier transformation. The water signal, still present in the sample, was suppressed by a saturation pulse of the signal itself, with a duration of 2 s. For the acquisition of the spectra, 32 K data points per scan and the accumulation of 128 transients were used (Figs. 1 and 2).

For statistical analyses, we first performed principal component analysis with an unsupervised technique to find trajectories and clustering in our data [12,13]. Then partial least squares regression discriminant analysis was performed to identify differences in the metabolites profiles [14,15].

Table 1
Discriminating features of enrolled mothers and time of samples collection

Population Characteristics	Preterm (n = 6)	Term (n = 12)
Mean gestational age at the time of delivery (wk), SD	30 \pm 1	40 \pm 1
Mean mother's age at the time of delivery (y), SD	35 \pm 5	31 \pm 3
Mean time of milk collection (days after birth), SD		
First sample	5 \pm 2	6 \pm 1
Second sample	14 \pm 2	
Third sample	21 \pm 3	

Results

Partial least squares regression discriminant analysis performed on autoscaled data indicated a distinct grouping of milk samples and a significant difference between the metabolic profiles of milk from mothers with preterm delivery and the milk from mothers of term infants, at least until the fourth week (Fig. 3).

Profiling analysis led to the identification of the metabolite as lactose (4.46 ppm), which is present in higher concentration in preterm milk samples. Furthermore, milk samples from mothers of preterm infants had higher peaks of oligosaccharides, especially those fucosylated, such as fucose at a concentration of 1.23 ppm, N-acetyl-neuraminic acid at 2.06 ppm, and N-acetyl-glucosamine at 2.04 ppm (Fig. 4, Table 2).

Discussion

Human breast milk does not have a single composition but rather changes over time, from colostrum to transitional milk and finally to mature milk, during the course of lactation [2]. Understanding the composition of human milk therefore provides an important tool for planning infants' feeding, particularly for fragile, high-risk infants such as those born preterm [3]. Metabolomics should be considered as a key technology for this purpose. Measuring in a quantitative manner the complete set of low-molecular-weight metabolites that are the end products of gene expression, NMR metabolome is able to provide a molecular snapshot of biological fluids as human milk [16]. Using the NMR metabolome approach, we found that milk from mothers delivering preterm infants undergoes a sort of “maturation” in the first 3 wk of lactation, but at the end of the third week postpartum, it does not yet resemble the milk of mothers with term deliveries because of its higher level of lactose and oligosaccharides. The technique that we used intercepts small molecules but not amino acid. The disaccharide lactose is the principal sugar of human milk and the most common macronutrient, whereas oligosaccharides constitute the third most common solid component of human milk after lactose and lipids and they have important non-nutritional properties [17,18]. Synthesized by glycosyltransferases, human milk oligosaccharides range from 3 to 32 sugars in size and differ in composition from those of any other mammal [17] because they are species specific. They have lactose at the reducing end and fucose or sialic acid at the non-reducing end [19]. In contrast with bovine milk, most of human milk oligosaccharides are fucosylated, including 1 to 15 fucoses, and their production depends on enzymes encoded by the genes associated with the expression on the Lewis group system [18]. The presence of fucosylated oligosaccharides in human milk depends on a maternal secretory status that is determined by the presence of an α -1,2-fucosyltransferase gene (*FUT2*), and it is identified by NMR spectroscopy by the level of 2-fucosyllactose [20,21]. Higher levels of this metabolite in preterm milk than term samples were identified by Sundekilde et al. [21] in a metabolomics study. They also found differences between preterm and full-term milk in the level of lactose, citrate, and other metabolites [21].

It is now widely accepted that oligosaccharides are non-nutritive bioactive factors whose primary site of activity is the lumen of the gut. Oligosaccharides and their protein conjugates function as soluble “decoy” receptors for pathogens that inhibit the adhesion of pathogens to oligosaccharides receptors expressed on the gut surface. In addition, they have been described as exerting immunomodulatory effects [22]. In addition, oligosaccharides are prebiotic agents that selectively encourage the growth of commensal bacteria. The advantages for the gastrointestinal tract include dose-dependent protection against necrotizing enterocolitis and




Fig. 1. Spectrum representing the metabolomic profiling of human breast milk collected from mothers delivering at term.




Fig. 2. Spectrum representing the metabolomic profiling of human breast milk collected from mothers delivering prematurely.




Fig. 3. Discriminant plot partial least squares regression discriminant analysis. Two distinct metabolic groups are represented: The bottom left circle denotes preterm human milk samples collected during the 3 wk of lactation after birth; the upper right circle denotes full-term human milk collected between 4 to 7 d after parturition.

late-onset sepsis [23–26]. There is also a trophic effect on the gut in terms of more rapid maturation of motility and a more rapid decrease in permeability. This results in lower gastric residual volumes and facilitates early achievement of full feedings when prebiotics are used together with probiotics [23]. Prebiotics also promote growth of *Bifidobacteria* and decrease growth of pathogens in the gut [27].

It is also possible that an amount of oligosaccharides indirectly affects sites distant from the gut, perhaps through secondary metabolites from fermentation that traverse the bloodstream. In this way they protect infants from late-onset sepsis and, when eventually excreted into the urine, reduce the risks of urinary

tract infections [28]. For these reasons, human milk oligosaccharides content should be looked at as part of the innate immune system by which mothers protect their infants from disease by breastfeeding [18].

Finding that milk collected from mothers of preterm has higher bioactive properties than the milk collected from mothers delivering at term is crucial for tailored, individualized nutrition.

Although this study is limited by the small number of samples, the data come from a well-defined population, with no variability with respect to location, health, and lifestyle. The strength of the paper is the use of highly sophisticated method to evaluate the metabolomic profile of human preterm breast milk such as NMR spectroscopy.

The obtained data suggest that human milk is not only species specific but also gestational age specific.

In current clinical practice, if the mother has insufficient milk for preterm infants, milk from donor mothers delivering at term is considered an alternative [29]. Unfortunately, the differences in the metabolic profiles of breast milk of women who delivered preterm newborns compared with term might lead to some nutritional deficiencies in preterm newborns, affecting their proper growth [30,31]. The present study sheds a light on the need to use a more personalized nutrition in preterm newborns by adding oligosaccharides to breast milk from mothers delivering at term.

Conclusions

Our results contribute to understanding the peculiar nutritional needs of breastfed preterm newborns. The high content of oligosaccharides in preterm breast milk, assessed through metabolomic investigation, provides new insights about dietary modification to attain precise and specific nutritional characteristics of preterm breast milk. From the point of view of precision medicine, to promote proper growth, it is desirable that preterm infants receive




Fig. 4. Comparison of the spectra representing metabolite profiling identified that lactose (4.46 ppm) and oligosaccharides levels, especially those fucosylated (fucose, 1.23 ppm; N-acetyl-neuraminic acid, 2.06 ppm; N-acetylglucosamine, 2.04 ppm) were significantly higher in preterm milk samples compared with full-term milk.

Table 2

NMR identification of metabolites in breast milk

Metabolites	1 H chemical shift (ppm)
Lactose	4.46 ppm
Fucose	1.23 ppm
N-acetyl-neuraminic acid	2.06 ppm
N-acetyl-glucosaminic acid	2.04 ppm

H, hydrogen; NMR, nuclear magnetic resonance.

their own mother's milk or at least donor human milk from lactating mothers who delivered at the same gestational age.

Acknowledgments

The authors thank the EUROpe Against Infant Brain Injury (EUR-AIBI) foundation for its partial grant.

References

- [1] American Academy of Pediatrics, Section on Breastfeeding. Breastfeeding and the use of human milk. *Pediatrics* 2012;129(3):e827–41.
- [2] Agostoni C, Braegger C, Decsi T, Kolacek S, Koletzko B, Michaelsen KF, et al. Breastfeeding: a commentary by ESPGHAN committee on nutrition. *J Pediatr Gastroenterol Nutr* 2009;49(1):112–25.
- [3] Ballard O, Morrow AL. Human milk composition: nutrient and bioactive factors. *Pediatr Clin North Am* 2013;60(1):49–74.
- [4] Gartner LM, Morton J, Lawrence RA, Naylor AJ, O'Hare D, Schanler RJ, et al. Breastfeeding and the use of human milk. *Pediatrics* 2005;115(2):496–506.
- [5] Li R, Dee D, Li CM, Hoffman HJ, Grummer-Strawn LM. Breastfeeding and risk of infections at 6 years. *Pediatrics* 2014;134(Suppl.1):S13–20.
- [6] Ajetonmobi OM, Whyte B, Chalmers J, Tappin DM, Wolfson L, Fleming M, et al. Breastfeeding is associated with reduced childhood hospitalization: evidence from Scottish Birth Cohort (1997–2009). *J Pediatr* 2015;166(3):620–5.
- [7] Lonnerdal B. Infant formula and infant nutrition: Bioactive proteins of human milk and implications for composition of infant formulas. *Am J Clin Nutr* 2014;99(3):712S–75S.
- [8] Longini M, Tataranno ML, Proietti F, Tortoriello M, Belvisi E, Vivi A, et al. A metabolomic study of preterm human and formula milk by proton MRS analysis: preliminary results. *J Matern Fetal Neonatal Med* 2014;27(Suppl.2):27–33.
- [9] Marincola SC, Noto A, Caboni P, Reali A, Barberini L, Lussu M, et al. A metabolomic study of preterm human and formula milk by high resolution NMR and GC/MS analysis: preliminary results. *J Matern Fetal Neonatal Med* 2012;25(Suppl.5):62–7.
- [10] Moro GE, Arslanoglu S, Bertino E. Human milk in feeding premature infants: consensus statement. *J Pediatr Gastroenterol Nutr* 2015;61(Suppl.1):S16–9.
- [11] Yehye WA, Rahman NA, Ariffin A, Abd Hamid SB, Alhadi AA, Kadir FA, et al. Understanding the chemistry behind the antioxidant activities of butylated hydroxytoluene(BHT): a review. *Eur J Med Chem* 2015;101:295–312.
- [12] Jackson JE. A user's guide to principal components. New York(NY: John Wiley and Sons; 1991.
- [13] Jolliffe IT. Principal components analysis. New York(NY: Springer; 1986.
- [14] Wold H. In: Giani J. Perspectives in probability and statistics. Papers in honor of M. S. Bartlett—Soft modeling by latent variables: the nonlinear iterative partial least squares approach. London; 1975. p. 117–42.
- [15] Wold S, Sjostrom M, Eriksson L. PLS-regression: a basic tool for chemometrics. *Chem Intell Lab Syst* 2001;58(2):109–30.
- [16] Marincola CF, Dessì A, Corbu S, Reali A, Fanos V. Clinical impact of human breast milk metabolomics. *Clin Chim Acta* 2015;451(Pt A):103–6.
- [17] Bode L. Human milk oligosaccharides: Every baby needs a sugar mama. *Glycobiology* 2012;22(9):1147–62.
- [18] Newburg DS, Ruiz-Palacios GM, Morrow AL. Human milk glycans protect infants against enteric pathogens. *Annu Rev Nutr* 2005;25:37–58.
- [19] Morrow AL, Ruiz-Palacios GM, Jiang X, Newburg DS. Human milk glycans that inhibit pathogen binding protect breast-feeding infants against infectious diarrhea. *J Nutr* 2005;135(5):1304–7.
- [20] Praticò G, Capuani G, Tomassini A, Baldassarre ME, Delfini M, Micheli A, et al. Exploring human breast milk composition by NMR-based metabolomics. *Nat Prod Res* 2014;28:95–101.
- [21] Sundekilde UK, Downey E, O'Mahoney JA, O'Shea CA, Ryan CA, Kelly AL, et al. The effect of gestational and lactational age on the human milk metabolome. *Nutrients* 2016;8(5):304.
- [22] Plaza-Díaz J, Fontana L, Gil A. Human milk oligosaccharides and immune system development. *Nutrients* 2018;10:1038.
- [23] Ziegler EE. Advantages of human milk in feeding premature infants. *J Pediatr Gastroenterol Nutr* 2015;61(Suppl.1):S3.
- [24] Sullivan S, Schanler RJ, Kim JH, Patel AL, Trawöger R, Kiechl-Kohlendorfer U, et al. An exclusively human milk-based diet is associated with a lower rate of necrotizing enterocolitis than a diet of human milk and bovine milk-based products. *J Pediatr* 2010;156(4):562–7.
- [25] Neu J, Walker WA. Necrotizing enterocolitis. *N Engl J Med* 2011;364(3):255–64.
- [26] Jantscher-Krenn E, Marx C, Bode L. Human milk oligosaccharides are differentially metabolised in neonatal rats. *Br J Nutr* 2013;110(4):640–50.
- [27] Keunen K, van Elburg RM, van Bel F, Benders MJ. Impact of nutrition on brain development and its neuroprotective implications following preterm birth. *Pediatr Res* 2015;77(1-2):148–55.
- [28] Newburg DS, Grave G. Recent advances in human milk glycobiology. *Pediatr Res* 2014;75(5):675–9.
- [29] Schanler RJ. Mother's own milk, donor human milk, and preterm formulas in the feeding of extremely premature infants. *J Pediatr Gastroenterol Nutr* 2007;45(Suppl.3):S175–7.
- [30] Kreissl A, Zwiauer V, Repa A, Binder C, Thanhaeuser M, Jilma B, et al. Human milk analyzer shows that the lactation period affects protein levels in preterm breast milk. *Acta Paediatr* 2016;105(6):635–40.
- [31] Boyd CA, Quigley MA, Brocklehurst P. Donor breast milk versus infant formula for preterm infants: A systematic review and meta-analysis. *Arch Dis Child Fetal Neonatal Ed* 2007;92(3):F169–75.