

signalling pathways activated by INSL5 at RXFP4 is essential for understanding the biological roles of this novel gut hormone.

Reference

[1] Grosse, et al. Proc Natl Acad Sci U S A 2014;111:33–8.

<https://doi.org/10.1016/j.orcp.2016.10.055>

55

Insights into the trajectory of neuronal projections to brown adipose tissue derived from the use of novel “brainbow” neurotropic viruses

Brian Oldfield*, Nicole Weidmann, Aneta Stefanidis

Monash University, Clayton, VIC, Australia

The medullary raphe nuclei are regarded as the final common neural relay connecting higher brain centres to the spinal cord and sympathetic outflow to brown adipose tissue (BAT), an observation which is based on functional (electrophysiological) data. In order to define the trajectory of populations of neurons in the hypothalamus, synaptically linked to the medullary raphe nuclei and ultimately BAT, a novel approach was utilised which involves the injection of a modified form of pseudorabies virus (PRV) into the BAT. This changes the colour of its fluorescent reporter when it comes into contact with Cre recombinase. We hypothesise that contrary to the dogma stated at the outset there will be two populations of neurons projecting to BAT, one which passes through the midline raphe and the other that involves alternate premotor pathways in the brainstem.

An (AAV)-Cre recombinase construct was injected stereotaxically into the raphe nuclei of male Sprague Dawley rats weighing between 230 and 250g. This injection was followed 2 weeks later by injection of PRV Brainbow virus (PRV-263) into the interscapular BAT. After 4 days survival, rats were killed and their brains prepared for histological analyses.

PRV-263 which is replication competent was transported retrogradely from the BAT through chains of synaptically-connected neurons in the spinal cord, brainstem and hypothalamus including the midline raphe. After transport through neurons expressing Cre recombinase in the raphe there was recombination of the viral genome



at either paired lox2272 or loxP sites, resulting in the loss of the red reporter and expression of either cyan (mCerulean) or yellow (eYFP). Importantly there were distinct groups of neurons in the rostroventrolateral medulla, lateral hypothalamus and paraventricular nucleus that retained their red fluorescent reporter consistent with a trajectory other than through the raphe nuclei.

These data define the nature of descending neural projections to BAT.

<https://doi.org/10.1016/j.orcp.2016.10.056>

56

Oleylethanolamine and endocannabinoid responses to intraduodenal lipid infusion in humans: Relationships with BMI and energy intake

Tanya J. Little^{1,*}, Nada Cvijanovic¹, Donovan Argueta², Jaspreet Kaur², Christine Feinle-Bisset¹, Richard Young¹, Chris Rayner¹, Nicholas DiPatrizio²

¹ University of Adelaide, Adelaide, SA, Australia

² Biomedical Sciences, University of California Riverside, Riverside, USA

Introduction/aims: Fat stimulates intestinal secretion of oleylethanolamide (OEA) to reduce food intake, while fasting induces intestinal endocannabinoids with an orexigenic effect. Animal studies suggest that high-fat diet-induced obesity impairs intestinal control of endocannabinoid and OEA production, contributing to reduced satiety and weight gain. The aims of this study were to: (i) evaluate effects of intraduodenal (ID) lipid infusion on plasma levels of anandamide (AEA), 2-arachidonylethanolamide (2-AG) and OEA in humans, and to examine relationships with BMI and *ad libitum* energy and fat intakes, and (ii) to evaluate effects of ID lipid on duodenal concentrations of 2-AG, AEA and OEA.

Methods: 19 lean, 16 overweight and 17 obese participants underwent ID Intralipid[®] infusion (2 kcal/min) for 120 min during which blood samples were collected every 30 min. *Ad libitum* energy intake was assessed at a subsequent buffet meal. Endoscopic duodenal biopsies were collected from 4 lean participants, at baseline, and following 30 min ID Intralipid[®] infusion (2 kcal/min). Plasma and duodenal 2-AG, AEA and OEA concentrations were assessed by HPLC/tandem mass spectrometry

