

**Aim:** To assess the safety, efficacy, and mechanisms of the DJBS in 4 pilot patients with obesity and type 2 diabetes who had failed all previous conservative interventions.

**Methods:** The DJBS was placed endoscopically and left in situ for 48 weeks. Subjects received medical and dietetic support throughout. Metabolic, functional, psychological and dietary intake investigations were performed at baseline, and monthly or bi-monthly thereafter.

**Results:** All patients completed the 48 week period with devices in place. No device-related events were observed. The median weight loss was 27.85 kg (21.5–32.4), or 23.17% (19.76–26.51) of body weight. During the study period, mean daily energy intake was reduced as compared to baseline (baseline mean 6737 kJ (range 5156–7750) as compared to week 48: 3845 kJ (range 3442–5871)). Liver function tests substantially improved (median alanine transaminase baseline 37 U/L (15.5–73.5) as compared to week 48: 17 U/L (12.5–22.5); median aspartate transaminase baseline 25.5 U/L (21.5–51.5) as compared to week 48: 18 U/L (15–22)). In parallel, median glycated haemoglobin decreased from 6.8% (range 5.9–7.9) to 5.8% (range 5.7–6.0) at week 48. No significant decrease in circulating micronutrient concentrations was observed. Dietary quality did not change.

**Conclusion:** Weight loss during treatment with a DJBS in the setting of a multi professional team approach is clinically meaningful and appears to be largely explained by decreased energy intake. The small pilot study did not provide evidence for malabsorption of micronutrients.

<https://doi.org/10.1016/j.orcp.2016.10.082>

82

### Do making habits or breaking habits influence weight loss and weight loss maintenance? A randomised controlled trial



Gina Cleo<sup>1,\*</sup>, Rae Thomas<sup>2</sup>,  
Elisabeth Isenring<sup>1</sup>, Paul Glasziou<sup>2</sup>

<sup>1</sup> Bond University, Robina, QLD, Australia

<sup>2</sup> Faculty of Health Sciences and Medicine, Centre for Research in Evidence Based Practice, Robina, QLD, Australia

**Background:** Despite the significance placed on lifestyle interventions for obesity management, around 40% of weight loss is regained over the first year following treatment, and much of the rest

over the next three years. Two psychological concepts (habitual behaviour and automaticity) have been suggested as the most plausible explanation of this overwhelming lack of long-term weight loss success.

**Method:** We evaluated the efficacy of two interventions that explore these theories: Ten Top Tips (10TT) and Do Something Different (DSD). 10TT promotes automaticity; this is the ability to perform tasks without awareness or deliberation. Therefore, diet and exercise related behaviours become automatic or habitual. Conversely DSD promotes behavioural flexibility. This program disrupts daily routines by assigning an individual with unstructured tasks to perform. Behavioural flexibility therefore has an inverse relationship with automaticity and is defined as the measure of an individual's range of mindful behaviours. In previous studies, both interventions have achieved significant weight loss with results suggesting potential for maintenance of the weight lost. The research however is limited and long-term (12 month) results are yet to be explored. Men and women ( $n=75$ ), aged  $51+6$  (s.d.) years with body mass index  $34.5+4.1$  kg/m<sup>2</sup> were randomised to 12-week 10TT, DSD or no treatment control. Active intervention participants underwent 12 weeks of the program with 12-months follow-up.

**Results:** We collected data for weight, BMI, waist circumference as well as habitual behaviour and wellbeing. After 12 weeks intervention, weight loss averaged 4.6 kg in the 10TT group, 4.1 kg in the DSD group and 1.3 kg in the control group. There was significant improvement in wellbeing in the 10TT and DSD groups.

**Significance of research:** Results from this RCT have the potential to help in understanding the mechanisms relating to weight loss maintenance.

<https://doi.org/10.1016/j.orcp.2016.10.083>

83

### How many Australian women will be obese in twenty years' time?



Annette Dobson\*, Hsiu-Wen Chan,  
Richard Hockey, Gita Mishra

Australian Longitudinal Study on Women's Health, Herston, QLD, Australia

Successive generations of Australians are becoming fatter. The prediction of future trends in obesity is necessary in order to plan for future health service needs to manage the medical consequences of obesity, such as type II diabetes.