

316

### Adolescents with severe obesity attending a weight management clinic in Singapore



Wen Wei Christopher Ho<sup>2,1,\*</sup>, J.Y. Oh<sup>2,1</sup>, K.K. Rajasegaran<sup>2,1</sup>, Elaine C.S.E. Chew<sup>2,1</sup>

<sup>1</sup> KK Women's and Children's Hospital, Singapore

<sup>2</sup> Singhealth, Singapore, Singapore

**Background:** Adolescent obesity has led to chronic medical conditions such as type 2 diabetes mellitus, hypertension, and dyslipidemia, which used to be typically seen in adults. The earlier presentation of these comorbidities will have a significant impact for the future as this group of adolescents will present with complications of these chronic medical conditions at an earlier age and will have a significantly decreased life expectancy. American Academy of Pediatrics expert committee proposed the recognition of a BMI of above 35 kg/m<sup>2</sup>, which is above 99th percentile on the BMI chart, as extreme obesity.

**Objectives:** Objective is to look at the prevalence of adolescents with severe obesity attending a weight management clinic aged 13–17 years old and their co-morbidities.

**Methods:** This is a retrospective study of children aged 13 to 17 years old, who attended the weight management clinic, from January 2009 to December 2012. Patients' clinical, anthropometric, polysomnography and metabolic blood test results were collected.

**Results:** A total of 59 adolescents (64% males) with BMI above 35 kg/m<sup>2</sup> attended the weight management clinic. Mean age at presentation was 14.9 ± 0.97 years. 40 adolescents underwent oral glucose tolerance test and 7 adolescents underwent polysomnography. 4 adolescents had diabetes mellitus based on their oral glucose tolerance test. 4 adolescents had severe obstructive sleep apnea based on polysomnography results.

**Conclusion:** There is a significant proportion of adolescents with severe obesity attending the weight management clinic with obesity related co-morbidities. More studies should look into the outcomes of adolescents with extreme obesity and the role of surgical treatment.

<https://doi.org/10.1016/j.orcp.2018.11.204>

317

### Change in self-esteem following paediatric weight management interventions: a systematic review and meta-analysis



Melissa Su Yi Tee<sup>1</sup>, Natalie Lister<sup>2,3</sup>, Megan L. Gow<sup>2,3</sup>, Susan J. Paxton<sup>4</sup>, Katharine Aldwell<sup>3</sup>, Sarah Thomas<sup>3</sup>, Sarah Garnett<sup>2,3</sup>, Louise Baur<sup>2,3</sup>, Hiba Jebeile<sup>2,3,\*</sup>

<sup>1</sup> Faculty of Medicine and Health, The University of Sydney, Sydney, NSW, Australia

<sup>2</sup> Children's Hospital Westmead Clinical School, The University of Sydney, Sydney, NSW, Australia

<sup>3</sup> Institute of Endocrinology and Diabetes and Weight Management Services, The Children's Hospital at Westmead, Westmead, NSW, Australia

<sup>4</sup> School of Psychology and Public Health, La Trobe University, Melbourne, VIC, Australia

**Objective:** Children and adolescents with overweight or obesity are likely to have lower self-esteem compared to normal weight peers. Weight management interventions are successful at reducing BMI however, the effect on self-esteem is unclear. This systematic review investigates the impact of weight management

interventions in children and adolescents with overweight or obesity on self-esteem.

**Methods:** Four databases were searched in May 2017 to retrieve peer-reviewed articles that met the following eligibility criteria: (1) children and adolescents with overweight or obesity, aged < 18 years, (2) weight management intervention with a nutrition component, (3) case studies, pre-post studies and randomised controlled trials (RCTs), (4) pre- and post-intervention assessment of self-esteem using a validated tool.

**Results:** Of 2799 articles screened, 40 studies (9.3–15.6 years,  $n = 10131$ ) met the inclusion criteria. Meta-analyses showed a medium positive effect for self-esteem post-intervention (one week to 15 months) (standardised mean difference (SE) = 0.42 [0.05]  $p < 0.001$ ). Sixteen studies had follow-up measurements (2.5 months to 2.4 years post-intervention) which indicated the effect was maintained (0.36 [0.05],  $p < 0.001$ ). However, the effect is unlikely to be clinically significant. Analysis of RCTs (eight studies), supported these findings. Studies with a targeted self-esteem component had a larger effect (0.558 [0.13],  $p < 0.001$ ) than studies with no self-esteem component (0.373 [0.05],  $p < 0.001$ ). Studies which included parental involvement had a larger effect (0.417 [0.05],  $p < 0.001$ ) than studies with no parental involvement (0.365 [0.06],  $p < 0.001$ ). In terms of setting, there was no difference in effect between inpatient programs (0.477 [0.23],  $p = 0.038$ ), outpatient programs (0.452 [0.06],  $p < 0.001$ ) and summer camps (0.440 [0.06],  $p < 0.001$ ), with the smallest effect observed in a community setting (0.319 [0.05],  $p < 0.001$ ).

**Conclusion:** Engagement in weight management interventions was not associated with a decrease in self-esteem. The inclusion of targeted self-esteem components within weight management may lead to a clinically significant improvement.

<https://doi.org/10.1016/j.orcp.2018.11.205>

318

### Childhood obesity, weight management and depression: a systematic review and meta-analysis



Hiba Jebeile<sup>1,2,\*</sup>, Megan L. Gow<sup>1,2</sup>, Susan Paxton<sup>3</sup>, Katharine Aldwell<sup>2</sup>, Sarah Thomas<sup>2</sup>, Sarah P. Garnett<sup>1,2</sup>, Louise A. Baur<sup>1,2</sup>, Natalie B. Lister<sup>1,2</sup>

<sup>1</sup> Children's Hospital at Westmead Clinical School, The University of Sydney, Sydney, NSW, Australia

<sup>2</sup> Institute of Endocrinology and Diabetes, and Weight Management Services, The Children's Hospital at Westmead, Westmead, NSW, Australia

<sup>3</sup> School of Psychology and Public Health, La Trobe University, Melbourne, Victoria, Australia

**Introduction:** Child and adolescent overweight and obesity is associated with depression, and may contribute to increased depression into adulthood. The aim of this systematic review and meta-analysis was to investigate the impact of weight management interventions on depression in children and adolescents with overweight or obesity.

**Methods:** Four databases were searched in May 2017 to retrieve peer-reviewed articles that met the following eligibility criteria: (1) children and adolescents (<18 years) with overweight or obesity; (2) weight management intervention with a nutrition component; (3) case studies, pre-post studies, RCTs; (4) pre- and post-intervention assessment of depression using a validated tool.

**Results:** Of 2799 articles screened, 29 studies ( $n = 1542$ , 9.5–16.5 years) met the inclusion criteria. Meta-analyses showed reduced

levels of depression post-intervention (one week to 15 months, standardised mean difference, SMD [SE] =  $-0.47$  [0.077],  $p < 0.001$ ), which was maintained at follow-up (6–16 months, six studies,  $-0.528$  [0.21],  $p = 0.011$ ). However, the absolute reduction was small ( $-2.8$  points on a scale of 0–54) and may not be clinically significant. BMI z-score was reduced post-intervention (11 studies, mean difference [SE] =  $-0.183$  [0.014],  $p < 0.001$ ) and at follow-up (four studies,  $-0.261$  (0.022),  $p < 0.001$ ). Interestingly, change in depression did not differ between studies using an energy prescription (SMD [SE]  $-0.511$  [0.21],  $p = 0.015$ ) compared to healthy eating education ( $-0.452$  [0.068],  $p < 0.001$ ). Interventions with structured exercise classes had the highest effect size ( $-0.533$  [0.107],  $p < 0.001$ ), followed by interventions providing physical activity education ( $-0.388$  [0.092],  $p < 0.001$ ). Interventions which did not report incorporating or encouraging physical activity did not significantly change depression symptoms ( $-0.137$  [0.14],  $p = 0.335$ ).

**Conclusion:** Engagement in weight management interventions did not worsen depressive symptoms. In fact, weight management may lead to mild improvements and disrupt the progression of depression in children and adolescents with overweight or obesity. Inclusion of physical activity within treatment appears to be important.

<https://doi.org/10.1016/j.orcp.2018.11.206>

319

### Lifestyle interventions in postpartum women: a systematic review and meta-analysis of implementation characteristics

Siew S. Lim<sup>1,2,\*</sup>, Xinyu Liang<sup>2</sup>, Briony Hill<sup>2</sup>, Lisa Moran<sup>2</sup>, Sharleen O'Reilly<sup>3</sup>

<sup>1</sup> Monash University, Clayton, VIC, Australia

<sup>2</sup> Monash Centre for Health Research and Implementation, Monash University, Clayton, Australia

<sup>3</sup> University College Dublin, Dublin, Ireland

**Introduction:** Postpartum weight retention is a significant contributor to obesity in women of reproductive age. While a number of randomised controlled trials (RCTs) have looked at reducing weight retention in postpartum women, the impact of implementation characteristics on effectiveness is not known.

**Methods:** MEDLINE, CINAHL, EMBASE, PsycInfo, and EBM databases were searched to identify lifestyle intervention RCTs in postpartum women (within 2 years post-delivery) published up to January 2018. The primary outcome was weight change. Population penetration and participation rates were calculated. Subgroup analyses were performed for intervention provider, intervention duration, number of sessions, self-monitoring, use of technology, group or individual setting, attrition, fidelity and intervention types (diet/physical activity/combined).

**Results:** From 4512 studies, 34 studies were included in the systematic review and meta-analysis ( $n = 5000$  women, age 18–34 years). One study reported their penetration rate (2.5%) and five studies reported their participation rate (0.94% to 86%). Studies with self-monitoring had significantly greater weight loss than those without (mean difference  $-2.93$  kg, 95% confidence interval [4.12,  $-1.74$ ] vs  $-1.29$  kg [ $-2.35$ ,  $-0.23$ ],  $P = 0.04$  for subgroup differences). Diet and physical activity combined were significantly more effective for weight loss compared with physical activity only interventions ( $-3.29$  kg [ $-4.51$ ,  $-2.06$ ] vs  $-0.53$  kg [ $-1.52$ ,  $0.46$ ],  $P = 0.003$  for subgroup differences). Health professional delivered interventions were more effective for weight loss than those that were not ( $-4.45$  kg [ $-7.16$ ,  $-1.75$ ] vs  $-0.76$  kg [ $-1.41$ ,  $-0.12$ ],

$P = 0.009$  for subgroup differences). Intervention duration, number of sessions, individual or group setting were not associated with weight loss.

**Conclusion:** Despite the large number of RCTs conducted in postpartum women, a lack of information on the population penetration and participation in these interventions exists. Self-monitoring, delivery by health professionals and combining diet and physical activity increases the effectiveness of lifestyle interventions on weight loss in postpartum women.

<https://doi.org/10.1016/j.orcp.2018.11.207>

320

### Obesity and retinal microvasculature in mid-childhood and mid-life: The Longitudinal Study of Australian Children



Mengjiao Liu\*, Kate Lycett, David Burgner, Melissa Wake

Murdoch Children's Research Institute, The University of Melbourne, Parkville, VIC, Australia

**Objective:** To examine whether microvasculature at age 11–12 years and in mid-adulthood is predicted by: BMI at six time points across the preceding the decade, and decade-long growth patterns.

**Methods:** *Participants/Design:* 1288 children (51% girls) and 1264 parents (87% mothers) in the Longitudinal Study of Australian Children (LSAC), followed biennially since 2004. *Exposures:* Child BMI z-score (CDC) and parent BMI ( $\text{kg}/\text{m}^2$ ) at child ages 2–3, 4–5, 6–7, 8–9, 10–11 and 11–12 years. *Outcomes:* Arteriolar and venular calibre from retinal photographs at the Child Health CheckPoint, LSAC's biophysical assessment at age 11–12 years. *Analyses:* Mixed growth curve models identified BMI trajectories (five for children and four for adults). Adjusted linear regression models (age, sex and socioeconomic position) estimated associations of retinal vascular calibre with BMI at each time point/BMI trajectories.

**Results:** Higher BMI at six time points was modestly associated with adverse retinal arteriolar and venular calibre in both age groups. For children, associations strengthened with age, doubling from 2–3 (arteriolar standardised mean difference (SMD)  $-0.04$  (95%CI  $-0.09$  to  $0.00$ ) and venular SMD:  $0.04$ , 95%CI  $-0.01$  to  $0.09$ ) to 11–12 years of age (SMD:  $-0.14$ , 95%CI  $-0.18$  to  $-0.09$  and SMD:  $0.08$ , 95%CI  $0.03$  to  $0.13$ , respectively); yet associations were consistent over time in adults. Compared to the 'consistently average' BMI trajectory children: those in 'low to high' and 'consistently very high' trajectories had narrower arteriolar calibre (SMD  $-0.23$  (95%CI  $-0.43$  to  $-0.03$ ) and  $-0.36$  (95%CI  $-0.51$  to  $-0.20$ ), respectively). Compared to 'consistently healthy' BMI adults, those in the always overweight/obese/severely obese trajectories had narrower arteriolar (SMD  $-0.22$  to  $-0.37$ ) and wider venular ( $0.14$  to  $0.31$ ) calibre.

**Conclusions:** Adverse microvascular markers at age 11–12 years are predicted by higher BMI from age 2–3 years. Decade-long BMI trajectories show small, but consistent associations, with adverse retinal microvasculature in both age groups.

<https://doi.org/10.1016/j.orcp.2018.11.208>