

junctions and endosomes, suggesting a role in endocrine function. This was linked to changes in hepatic expression of several genes linked to glucose homeostasis and lipid metabolism. The primary effect of PKC ϵ on glucose homeostasis is, therefore, not exerted directly in the liver as currently posited. However, PKC ϵ activity in adipose tissue modulates glucose tolerance and is involved in crosstalk with the liver.

<https://doi.org/10.1016/j.orcp.2018.11.082>

78

Lipidomic profiling reveals early-stage metabolic dysfunction in overweight or obese humans



Aya Mousa^{1,*}, Negar Naderpoor¹, Natalie Mellett², Kirsty Wilson³, Peter J. Meikle⁴, Barbara De Courten¹, Magdalena Plebanski^{3,5}

¹ Monash Centre for Health Research and Implementation, School of Public Health and Preventive Medicine, School of Public Health and Preventive Medicine, Melbourne, VIC, Australia

² Metabolomics Laboratory, Baker Heart and Diabetes Institute, Melbourne, VIC, Australia

³ Department of Immunology and Pathology, Monash University, Melbourne, VIC, Australia

⁴ Metabolomics Laboratory, Baker Heart and Diabetes Institute, Melbourne, VIC, Australia

⁵ School of Health and Biomedical Science, RMIT University, Melbourne, VIC, Australia

Advances in mass spectrometry and lipidomics techniques are providing new insights into the role of lipid metabolism in obesity and its metabolic sequelae. However, human lipidomic studies have produced inconsistent results, owing in part to the use of indirect proxy measures of obesity and insulin resistance and the relatively limited coverage of the lipidome. Here, we explored the relationship between the plasma lipidome and metabolic profiles using direct gold-standard measures of adiposity, insulin sensitivity, and insulin secretion, in addition to comprehensive lipidomic profiling (>450 species) and measurement of inflammatory cytokines and adipokines. We present new evidence showing a strong and independent positive correlation between the lysophosphatidylinositol (LPI) lipid class and insulin secretion *in vivo* in humans, supporting the insulinotropic effects of LPI demonstrated in mouse islets. Dihydroceramide, a sphingolipid precursor, was independently and negatively correlated with insulin sensitivity, indicating a possible upregulation in sphingolipid synthesis in obese individuals. We also show that phosphatidylethanolamine and its vinyl ether-linked (plasmalogen) derivatives correlate negatively with body fat, while dihexosylceramide correlates positively with interleukin-10. Together, these lipid classes may signify early pathogenesis toward type 2 diabetes and could serve as novel therapeutic targets or biomarkers for identification of high-risk individuals.

<https://doi.org/10.1016/j.orcp.2018.11.083>

79

Insights into type 2 diabetes susceptibility and resilience: TOFI Asia study



Ivana R. Sequeira^{1,*}, Wilson Yip¹, Louise W.W. Lu¹, Lindsay Plank², Rinki Murphy³, Kieren G. Hollingsworth⁴, Jun Lu⁵, Sally D. Poppitt^{1,3}

¹ Human Nutrition Unit, School of Biological Sciences, University of Auckland, Auckland, New Zealand

² Body Composition Unit, Department of Surgery, University of Auckland, Auckland, New Zealand

³ Department of Medicine, University of Auckland, Auckland, New Zealand

⁴ Newcastle Magnetic Resonance Centre, Institute of Cellular Medicine, Newcastle University, Newcastle upon Tyne, United Kingdom

⁵ Faculty of Health and Environmental Sciences, AUT, Auckland, New Zealand

Predicting rate of progression and identifying those most at risk of adverse metabolic health is difficult and often associated with environmental, lifestyle, and dietary factors. Importantly site of lipid deposition, including ectopic 'overspill' into key organs, increases metabolic risk. We investigated susceptibility to type 2 diabetes (T2D) in healthy and prediabetic Asian Chinese ($N = 209$) and European Caucasian ($N = 156$) adults resident in Auckland, enrolled in the TOFI Asia Study; within the Peak Nutrition for Metabolic Health (PANMAH) program, one of four priority research platforms in the New Zealand National Science Challenges (NSC).

Phenotypic characterisation of the 2 ethnicity cohorts, using anthropometry, body composition (dual energy X-ray absorptiometry, DXA) and blood biochemistry revealed that although of similar age and body mass index (BMI), Chinese (Mean $\pm$ SEM, 42 ± 1 yrs; 27.4 ± 0.3 kg/m²) had significantly greater abdominal fat ($42.0 \pm 0.6\%$ vs $37.0 \pm 1.1\%$, $p < 0.01$) than Caucasian (43 ± 1 yrs; 26.9 ± 0.4 kg/m²) with correspondingly higher fasting plasma glucose (FPG) concentrations (5.4 ± 0.04 mmol/l vs 5.0 ± 0.04 mmol/l, $p < 0.001$) and glycated haemoglobin, HbA_{1c} (36.0 ± 0.3 mmol/mol vs 33.0 ± 0.3 mmol/mol, $p < 0.05$). In addition, pancreatic and liver fat were quantified, using magnetic resonance imaging (MRI) and spectroscopy (MRS), in a subset of 36 Chinese (41 ± 2 yrs; 26.9 ± 0.7 kg/m²) and 34 Caucasian (48 ± 3 yrs; 28.0 ± 0.7 kg/m²) women from the cohort. These overweight, younger Asian Chinese had significantly lower MRI-determined abdominal and subcutaneous fat ($p < 0.05$) but pancreas ($4.3 \pm 0.3\%$) and liver ($11.0 \pm 1.7\%$) fat matched that, or tended to be higher than, Caucasian (pancreas: $4.1 \pm 0.3\%$; liver: $8.1 \pm 1.7\%$). Importantly, fat in pancreas (Chinese: $R^2 = 0.10$, $p = 0.09$; Caucasian: $R^2 = 0.28$, $p = 0.001$) and liver (Chinese: $R^2 = 0.12$, $p = 0.04$; Caucasian: $R^2 = 0.11$, $p = 0.05$) were positively correlated with FPG. The Thin on the Outside Fat on the Inside 'TOFI' profile observed in this Asian Chinese cohort, may contribute to their greater risk of poor metabolic health compared to Caucasian counterparts even in individuals of the same BMI and younger age.

<https://doi.org/10.1016/j.orcp.2018.11.084>