



## Correction to: A Comparison of the Bariatric Procedures that Are Performed in the Treatment of Super Morbid Obesity

Kohei Uno<sup>1,2</sup> · Yosuke Seki<sup>1</sup> · Kazunori Kasama<sup>1</sup> · Kotaro Wakamatsu<sup>1</sup> · Akiko Umezawa<sup>1</sup> · Katsuhiko Yanaga<sup>2</sup> · Yoshimochi Kurokawa<sup>1</sup>

Published online: 3 December 2018  
© Springer Science+Business Media, LLC, part of Springer Nature 2018

### Correction to: OBES SURG (2017) 27:2537–2545

<https://doi.org/10.1007/s11695-017-2685-7>

On page 2539, in the section “Patient Characteristics”, in line 6 “LSG/DJB, n = 47, and LRYGB, n = 70” should be corrected to “LSG/DJB, n = 50, and LRYGB, n = 47”. On page 2539, in the section “The Perioperative Outcome (Table 3)” in line 8 “LRYGB, 165 ± 42.6” should be corrected to “LRYGB, 160 ± 42.6”.

---

The online version of the original article can be found at <https://doi.org/10.1007/s11695-017-2685-7>

---

✉ Yosuke Seki  
y-seki@mcube.jp

<sup>1</sup> Weight Loss and Metabolic Surgery Center, Yotsuya Medical Cube, 7-7 Nibancho, Chiyoda-ku, Tokyo 102-0084, Japan

<sup>2</sup> Department of Surgery, Jikei University School of Medicine, 3-19-18 Nishishinbashi, Minato-ku, Tokyo 105-8471, Japan