



Predictors of a Healthy Eating Disorder Examination-Questionnaire (EDE-Q) Score 1 Year After Bariatric Surgery

Daniel Gero¹ · Stefanos Tzafos¹ · Gabriella Milos² · Philipp A. Gerber³ · Diana Vetter¹ · Marco Bueter¹

Published online: 4 January 2019

© Springer Science+Business Media, LLC, part of Springer Nature 2019

Abstract

Background Bariatric surgery (BS) has been shown to ameliorate health-related quality of life and eating disorder symptoms. However, the correlation of these changes with weight loss is not uniform, suggesting that additional factors have an impact on postoperative outcomes.

Objective To assess the impact of BS on eating disorder symptoms at 1 year postoperatively and to generate predictive models for the achievement of optimal eating behavior.

Methods Retrospective cohort study on a prospectively collected database of all consecutive patients who underwent primary BS in our academic center between January 2015 and March 2017. Eating Disorder Examination-Questionnaire (EDE-Q) was used to measure eating psychopathology. Logistic regression was used to estimate the odds ratio of achieving “healthy” EDE-Q at 1 year. Missing data was handled by multiple imputations for the regression model.

Results Two-hundred thirty-four patients were included. A complete-case analysis in 135 cases showed a “healthy” EDE-Q in 27.4% at baseline and in 83.7% at 1 year (difference = 56.3%, $P = 0.018$). Only the baseline EDE-Q “healthy” status influenced significantly the odds of achieving “healthy” EDE-Q at 1 year (OR 6.7, 95% CI 1.18–38.14, $P = 0.04$).

Conclusion BS seems to promote successful treatment of self-reported eating disorder symptoms during the first postoperative year. The achievement of optimal results is independent of age, sex, weight loss, obesity-related comorbidity status, surgical technique, or 30-day surgical complications. Future studies, using validated questionnaires specifically designed to investigate eating behavior after BS and/or direct measurements of the eating behavior are needed to clarify the underlying neuropsychologic mechanisms that drive the observed postoperative changes.

Keywords Eating Disorder Examination Questionnaire · Eating behavior · Bariatric surgery · Excess weight loss · Predictors of outcome · Roux-en-Y gastric bypass

Daniel Gero and Stefanos Tzafos contributed equally to this work.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s11695-018-3596-y>) contains supplementary material, which is available to authorized users.

✉ Marco Bueter
marco.bueter@usz.ch

Daniel Gero
daniel.gero@usz.ch

Stefanos Tzafos
Stefanos.Tzafos@usz.ch

Gabriella Milos
Gabriella.Milos@usz.ch

Philipp A. Gerber
Philipp.Gerber@usz.ch

Diana Vetter
Diana.Vetter@usz.ch

¹ Department of Surgery and Transplantation, University Hospital Zurich, Rämistrasse 100, 8091 Zürich, Switzerland

² Department of Psychiatry and Psychotherapy, University Hospital of Zurich, CH-8091 Zürich, Switzerland

³ Division of Endocrinology, Diabetes and Clinical Nutrition, University Hospital Zurich, Zürich, Switzerland

Introduction

Most studies using models to predict outcomes of bariatric surgery (BS) strived to identify predictors of successful weight loss [1–5]. However, modern concepts in BS relativize the role of weight loss as single criterion of postbariatric success and advocate offering BS preferably to patients with complications of obesity that would be expected to be resolved postoperatively [6, 7]. Screening tools to identify individuals who may benefit the most from BS need to be improved accordingly. Along these lines, Wood and Ogden suggested that recognition of factors that may predict the normalization of pathologic eating behaviors could represent another basis of improving patient selection for BS [8].

Patients tend to report lower level of baseline hunger, less frequent emotional eating, and faster intra-meal satiation following BS [9]. In addition, BS seems to induce changes in food preferences, with a shift from high-fat high-sugar nutrients to a less calorie-dense diet [10, 11], which has been recently hypothesized to beneficially influence postbariatric eating behavior [12]. Overall, BS has a beneficial effect on health-related quality of life and eating disorder symptoms; however, the correlation of these improvements with postoperative weight loss is not uniform [9, 11, 13, 14]. This suggests that additional factors may have an impact on the postbariatric amelioration of eating disorders.

Therefore, this study aimed to assess the prevalence of self-reported eating disorders symptoms before and at 1 year after BS by using the Eating Disorder Examination-Questionnaire (EDE-Q), and to generate predictive models for the achievement of optimal eating behavior in an explorative way, including demographic and health-related variables.

Methods

Participants

A retrospective cohort study was performed on a prospectively collected database of all consecutive patients who had sufficient German language skills and underwent primary laparoscopic BS (Roux-en-Y gastric bypass [RYGB] or sleeve gastrectomy [SG]) in our tertiary referral center from January 2015 to March 2017. Data analysis was performed in September 2018. According to the national guidelines of the Swiss Society for the Study of Morbid Obesity and Metabolic Disorders (SMOB), patients were eligible for surgery if they were aged over 18, had a BMI ≥ 40 kg/m² or BMI ≥ 35 kg/m² with obesity-related comorbidities, and had a 2-year history of failed conservative attempts to achieve sustained weight loss [15]. All patients were evaluated and prepared by a multidisciplinary team specialized in the

management of severely obese individuals. Data are reported in accordance with the STROBE guidelines [16].

Measurements

Data collection was prospective, took place at the outpatient visits, and included age, sex, objective BMI measurement, percentage of excess BMI loss (%EBMIL, with BMI = 25 considered ideal) [17], hip and waist measurements, comorbidities, and surgical complications according to the Clavien-Dindo classification [18].

The German translation of the EDE-Q [19] was used to measure eating disorder psychopathology as part of the regular preoperative evaluation. Patients filled in the EDE-Q alone at the outpatient clinic, with a physician at disposal in case of need. The questionnaire was administered again at the 1-year follow-up visit; however, patients were then allowed to bring the questionnaire home and send it back by mail. EDE-Q has well-established psychometric properties and has been validated in German cohorts [20]. The EDE-Q focuses on the previous 28 days and measures key eating disorder behavior and cognitive symptoms. The questions are rated on a 7-point Likert scale from 0 to 6, where a higher score indicates increased frequency of eating disorder symptoms. The EDE-Q consists of one total score and four subscales (restraint, eating concern, weight concern, and shape concern). Cutoff is defined in normal weight populations as mean total EDE-Q score plus one standard deviation, which is approximately 2.5–2.8, depending on different normative samples [21, 22]. In the current study, a threshold of < 2.5 was used to define absence of eating disorders and to dichotomize total EDE-Q outcomes to “healthy” and “unhealthy” subgroups.

Statistics

Demographic and clinical characteristics were presented as means and standard deviations (SD) or frequencies and percentages. Effect size for continuous variables was expressed as Cohen’s *d*, and for binary variables as odds ratios [23]. Cronbach’s alpha was used to measure internal consistency of the EDE-Q. Statistical testing was performed by (a) paired *t* test to assess changes from baseline to 1 year in continuous variables, (b) by independent sample *t* test for subgroup outcomes for continuous variables, and (c) by Wilcoxon rank sum tests with continuity correction for changes in categorical variables. Baseline characteristics of patients with and without available 1-year EDE-Q scores were compared to identify specific traits that might be overrepresented in patients lost to follow-up. A complete-case analysis was used for descriptive statistics. Logistic regression was used to estimate the odds ratio with a 95% confidence interval of different parameters that might influence the postoperative total EDE-Q score, dichotomized to “healthy” or “unhealthy” outcomes.

Based on recent studies, BMI, %EBMIL (grouped into four categories $\leq 50\%$, 50–75%, 75–100%, and $> 100\%$) [24], existence and persistence of comorbidities [25], age (grouped into three categories ≤ 30 , 30–50 and > 50 years), and sex were considered potential predictors of EDE-Q and were included in the analysis [26]. Other possible causal variables included in the calculation were as follows: type of procedure (RYGB or SG), presence or absence of any surgical complication during the 1st month, and baseline total EDE-Q score (dichotomized to “healthy”/“unhealthy”). Assuming that missing data in the 1-year EDE-Q scores occurred at random, five series of multiple imputations, using baseline and postbaseline variables selected for the regression model, were performed with the R package “mice” [27, 28]. The $m = 5$ imputed datasets were combined for the logistic regression model according to Rubin’s formula [29].

Statistical tests were two-tailed and $P < 0.05$ was considered significant. Analyses were performed using R software version 3.5.1 (The R foundation for Statistical Computing, Vienna, Austria).

Results

Study Population

The institutional database contained 234 cases of primary BS with available EDE-Q scores at baseline. Among those patients, 202 had a 1-year follow-up visit and 135 completed the EDE-Q at 1-year. There was no available information regarding the reasons of dropouts at 1-year. There were no differences in baseline characteristics of patients with and without 1-year EDE-Q follow-up (Supplemental Digital Content 1). Table 1 summarizes the characteristics of the study cohort available for complete-case analysis. Females were overrepresented in the study population (71.8%), the mean age was 40.3 ± 12.1 years, and the mean BMI 43.97 ± 6.5 kg/m². Significant decreases in body weight and BMI were recorded. The 1-year %BMI loss was $32.4 \pm 9\%$ (range 9.1–50.9) and the %EBMIL was $79.9 \pm 23.6\%$ (range 19.6–149.8). The 30-day surgical complications according to the Clavien-Dindo classification are presented in Table 2. Overall, 28.15% of patients presented a complication (defined as any deviation from the normal postoperative course [18]), although only 2.2% required a re-operation due to a complication and the 30-day mortality was null. Of note, between the 2nd and the 12th month, six patients (4.4%) underwent re-operation for internal hernia.

Self-Reported Eating Pathology Symptoms

The EDE-Q had a high internal consistency both at baseline and at 1 year (Cronbach’s $\alpha = 0.86$ and 0.85 ,

respectively). Overall, EDE-Q scores decreased significantly in total (Fig. 1) and for each subscale (Table 3). The total EDE-Q score of the participants was “healthy” in 27.4% of cases at baseline and in 83.9% at 1 year (difference = 56.3%, $P = 0.018$). There was a highly significant improvement in each individual EDE-Q question, except for EDE-Q-16 ($P = 0.837$) and EDE-Q-18 ($P = 0.197$). These questions consisted of the frequency of self-induced vomiting and compulsive exercise for weight/shape control. Most patients never engaged in such behavior at baseline (95.6% and 74.5%, respectively), which gave almost no room for significant improvement by 1 year postoperatively.

Subgroup analyses based on sex, age groups, and %EBMIL showed no clinically relevant differences in baseline or 1-year EDE-Q total and subscale outcomes (Supplemental Digital Content 2). Patients who achieved $> 75\%$ EBMIL at 1 year had a lower EDE-Q total score (1.25 vs. 1.64, 95% CI of difference in means: 0.065 to 0.714, $P = 0.019$); however, the mean value of both subgroups (%EBMIL \leq or $> 75\%$) was in the healthy range.

Correlations of Eating Disorders with BMI, EBMIL, and Age

The correlations between EDE-Q scores and BMI, %EBMIL, and age are illustrated in Supplemental Digital Content 3. Baseline total EDE-Q correlated negatively with baseline BMI ($r = -0.2$, $P = 0.022$), whereas at 1 year, the direction of the correlation tended to be positive ($r = 0.15$, $P = 0.082$). EDE-Q did not correlate significantly with age and correlated negatively only at the threshold of significance with %EBMIL. Nevertheless, delta EDE-Q correlated positively with 1-year %EBMIL ($r = 0.21$, $P = 0.016$).

Prediction Model of Healthy EDE-Q Scores at 1 Year

Prior to performing logistic regression on the database of the entire patient cohort ($n = 234$), the following missing data were accounted for by multiple imputations: healthy or unhealthy EDE-Q score at 1 year ($n = 99$), BMI at 1 year ($n = 32$), %EBMIL at 1 year ($n = 32$), age ($n = 4$), and presence or lack of any postoperative complications ($n = 1$). Results of the logistic regression on predictors of “healthy” 1-year total EDE-Q outcome are listed in Supplemental Digital Content 4. Only baseline EDE-Q “healthy” status was found to significantly predict a healthy EDE-Q score at 1-year after BS (OR 6.7, 95% CI 1.18–38.14, $P = 0.04$). All other parameters of our prediction model had no statistically significant predictive value for the achievement of a healthy EDE-Q score 1 year after BS.

Table 1 Characteristics of the study population with available 1-year follow-up ($n = 135$)

	Baseline	1 year	Mean of differences	Effect size* (95% CI)	<i>P</i>
Age at baseline, M \pm SD	40.29 \pm 12.07				
Female/male (<i>n</i>)	97 / 38				
Operation: RYGB/SG (<i>n</i>)	120 / 15				
BMI (kg/m ²), M \pm SD	43.97 \pm 6.53	26.69 \pm 5.4	14.29	3.4 (3–3.7)	< 0.001
Weight (kg), M \pm SD	124.23 \pm 26.01	83.82 \pm 19.41	40.33	2.9 (2.6–3.3)	< 0.001
Hip (cm), M \pm SD	122.4 \pm 44.5	114.6 \pm 89.76	7.05	0.07 (–0.17–0.31)	0.41
Waist (cm), M \pm SD	112.37 \pm 41.36	95.09 \pm 16.77	16.77	0.45 (0.2–0.7)	< 0.001
Hip/waist ratio, M \pm SD	1.097 \pm 0.13	1.218 \pm 0.93	0.132	–0.13 (–0.4–0.12)	0.152
Absence of obesity-related comorbidity (%)	12.59	44.44	31.85	5.5 (3–10)	< 0.001
Type 2 diabetes mellitus (%)	22.22	6.66	15.56	4 (1.82–8.8)	< 0.001
Arterial hypertension (%)	36.3	14.81	21.48	3.27 (1.81–5.91)	< 0.001
Sleep apnea syndrome (%)	30.37	12.59	17.77	3 (1.6–5.7)	< 0.001
Dyslipidemia (%)	24.44	8.89	15.55	3.3 (1.6–6.7)	< 0.001
Ischemic heart disease (%)	7.41	5.93	1.48	1.27 (0.5–3.3)	0.662
Joint pain (%)	65.18	28.89	36.3	4.6 (2.7–7.7)	< 0.001
Gastro-esophageal reflux disease (%)	40	11.11	28.89	5.3 (2.8–10.1)	< 0.001
Depression (%)	23.7	19.26	4.44	1.3 (0.7–2.3)	0.392

M mean, SD standard deviation, CI confidence interval, RYGB Roux-en-Y gastric bypass, SG sleeve gastrectomy, BMI body mass index

*Effect size for continuous variables: Cohen's *d*; effect size for proportions based on binary outcomes: odds ratio

Discussion

Pathologic eating symptoms, as measured by the EDE-Q, were very frequent among BS candidates. However, in most cases, pathologic eating symptoms normalized during the first year after BS. The achievement of a “healthy” eating behavior could not be predicted by baseline parameters (age, sex, BMI, comorbidity status or type of BS [RYGB or SG]), or by postoperative outcomes, such as 1-year %EBMIL or presence of surgical complications during the first postoperative month. The only significant predictor of “healthy” postoperative EDE-Q was a “healthy” EDE-Q score at baseline. Thus, in BS candidates with an unhealthy EDE-Q, we were unable to identify any factors predicting the normalization of their EDE-

Q scores at 1 year. However, as there was a significant positive correlation between %EBMIL and delta EDE-Q at 1 year, we cannot rule out that postsurgical weight loss at least partly contributed to the amelioration of eating disorders symptoms. Nutritional counseling might also have assisted in the observed improvement of EDE-Q scores after BS [30]. However, every patient in Switzerland has to undergo 2 years of unsuccessful conservative treatment including intense nutritional counseling before formally qualifying for BS [15]. Thus, the high incidence of pathological EDE-Q scores at baseline may indicate a high level of unresponsiveness to nutritional counseling in the study population.

The results of the present study are in line with previous studies investigating associated changes in prevalence

Table 2 Thirty-day surgical complications according to the Clavien-Dindo classification in the study population with available 1-year follow-up ($n = 135$) [18]

Grade	<i>n</i>	%	Description
I	7	5.19	Hypocalcemia, hypoesthesia of the thigh, minor bleeding
II	11	8.15	Perigastric hematoma, wound infection, pneumonia, urinary tract infection, bleeding, peripheral edema
IIIa	17	12.59	Endoscopic dilation of anastomotic stenosis, endoscopic treatment of bleeding at gastrojejunostomy, CT-guided pleural puncture
IIIb	3	2.22	Liver hematoma post-liver biopsy, intra-abdominal bleeding
IVa	0	0	
IVb	0	0	
V	0	0	
Total	38	28.15	




Fig. 1 Eating Disorder Examination-Questionnaire total score at baseline and at 1 year after bariatric surgery in patients with available 1-year follow-up ($n = 135$). Red dots: individual cases, green line: cutoff for healthy < 2.5 . The P value shows the result of paired t test statistic

of binge eating and weight loss following gastric banding [8]. Our findings are in further agreement with Conceição et al., who reported that preoperative EDE-Q scores did not correlate with weight loss at 2 years after gastric banding and RYGB. The same authors further found that problematic eating in the postoperative phase predicted worse weight loss trajectories [31]. Nasirzadeh et al. also found that the presence of eating disorders, especially binge eating, at 1 year after BS predicted future weight regain [32].

Together with the available literature mentioned above, our data emphasizes the need for systematic assessment of eating behaviors during routine bariatric follow-up. In this

context, additional live or internet-based behavioral and/or psychological interventions targeting patients with pathologic eating patterns have been recommended [31].

Bariatric patients' quality of life is known to be influenced by BMI [33], the extent of weight loss [34] and weight regain [35], and by the persistence of obesity-related diseases [33]. We therefore hypothesized that these factors along with age, sex, type of procedure, and surgical complications might interfere with the postoperative improvement of eating disorders. Surprisingly, none of these parameters influenced significantly the odds of achieving a healthy postoperative EDE-Q outcome, suggesting that BS has a uniform beneficial effect on eating behavior across very different patient subgroups. Examination of individual EDE-Q question scores showed some marked differences between subgroups; however, these specific questions were rather connected to weight-related anxiety and body shape perception than to the eating behavior itself.

The present study has a number of limitations that have to be considered when interpreting the results. Firstly, eating behavior psychopathology was measured indirectly, by a self-reported questionnaire, which is particularly prone to recall bias and potentially also to the biasing effect of body shame of obese patients. Nevertheless, the EDE-Q has been validated in different languages, both in paper and online format [36, 37]. Ideally, pre- and postoperative eating behavior should be measured *directly*, for example by a wearable system that detects eating episodes or by video recordings of a body camera worn by the patients over weeks [38]. At the price of ethical and methodological obstacles, videotaping would have enabled the measurement of the frequency and type of food intake, as well as how and when the nutrients were consumed (snacking, binge eating, night eating, vomiting, "picky" eating, etc.). Secondly, the EDE-Q was designed for the assessment of general eating disorders [39], and BS-specific eating pathologies such as dumping,

Table 3 Eating Disorder Examination-Questionnaire (EDE-Q) scores in total and in subscales at baseline and at 1 year after bariatric surgery in patients with available 1-year follow-up ($n = 135$)

		Mean \pm SD	Mean of differences	Cohen's d (95% CI)	P
Total	Baseline	3.24 \pm 1.3	1.80	1.23 (0.96–1.49)	< 0.001
	1 year	1.44 \pm 0.95			
Restrain	Baseline	2.71 \pm 1.5	1.25	0.68 (0.43–0.93)	< 0.001
	1 year	1.46 \pm 0.95			
Eating concern	Baseline	1.57 \pm 1.38	0.89	0.6 (0.35–0.84)	< 0.001
	1 year	0.68 \pm 1.02			
Weight concern	Baseline	3.73 \pm 1.22	2.02	1.2 (0.94–1.46)	< 0.001
	1 year	1.71 \pm 1.41			
Shape concern	Baseline	4.28 \pm 1.3	2.07	1.22 (0.96–1.48)	< 0.001
	1 year	2.2 \pm 1.45			

P values show the result of paired t tests

SD standard deviation, CI confidence interval

rumination, spitting out food, food “plugging,” or night-eating [12, 40–42] do not fall under its scope and thus were not assessed. This might have resulted in an underestimation of the frequency of postoperative de novo eating disturbances. Thirdly, the 1-year follow-up might have been too short. Several studies reported long-term recidivism in eating pathologies, notably in those patients who experience weight regain [32, 43]. Fourthly, the majority of patients in our cohort were female and underwent RYGB, therefore the overall outcomes (EDE-Q scores, comorbidity resolution, surgical complications) are less generalizable to males and to patients who undergo SG. Fifthly, although we failed to identify baseline differences between patients with and without available follow-up, our analyses may have been biased by the “missing not at random” effect [44].

Despite its limitations, this study has several important strengths, particularly the complete 1-year follow-up in a relatively large BS patient cohort that received standardized single-institutional perioperative care. The pathways that link the postbariatric neurophysiologic changes and the improvements in eating behavior are currently not entirely understood [36, 37, 39, 44, 45]. The presented findings might generate further hypotheses on the amelioration of eating behavior after BS and may foster the design of future studies, including assessment of “signature” postbariatric eating disorders [46] and *direct* measurement of pre- and postoperative food intake [47].

Conclusions

BS seems to promote a significant improvement of self-reported eating behavior disorders during the first postoperative year. The achievement of these positive results is independent of age, sex, weight loss, obesity-related comorbidity status, surgical technique, or surgical complications. The only predictor of healthy EDE-Q score at 1 year is a healthy EDE-Q score at baseline. Future studies are needed to clarify the underlying neuropsychologic mechanisms that drive the changes in eating behavior following BS, including validated BS-specific pathological eating behavior questionnaires and direct measurement of eating behavior.

Acknowledgements We are grateful to Samuel Aemisegger, clinical male nurse, to Mira Giama, and to the resident doctors of the Department of Surgery, University Hospital Zurich, for their valuable help in the data collection process. We also acknowledge Lukas Frick, MD, for his input in data analysis with the R software and Amy Taheri, PhD candidate, for her help in proofreading.

Funding The study was entirely funded by the assistant-professorship research grant awarded by the University of Zurich to Prof. Dr. med. Marco Bueter, PhD.

Compliance with Ethical Standards

Conflict of Interest The authors declare that they have no conflict of interest.

Ethical Approval and Informed Consent All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and cantonal research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. The Cantonal Ethics Committee of Zurich approved the study (nr: 2016-00022). For this type of study, formal consent is not required, as it was a retrospective study.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- Bordignon S, Aparicio MJG, Bertoletti J, et al. Personality characteristics and bariatric surgery outcomes: a systematic review. *Trends Psychiatry Psychother.* 2017;39:124–34.
- Livhits M, Mercado C, Yermilov I, et al. Preoperative predictors of weight loss following bariatric surgery: systematic review. *Obes Surg.* 2012;22:70–89.
- Krumpuri RD, Yokley JM, Seeholzer EL, et al. Qualifying for bariatric surgery: is preoperative weight loss a reliable predictor of postoperative weight loss? *Surg Obes Relat Dis.* 2018;14:60–4.
- Holsen LM, Davidson P, Cerit H, et al. Neural predictors of 12-month weight loss outcomes following bariatric surgery. *Int J Obes.* 42(4):785–793. <https://doi.org/10.1038/ijo.2017.190>
- Herpertz S, Kielmann R, Wolf AM, et al. Do psychosocial variables predict weight loss or mental health after obesity surgery? A systematic review. *Obes Res.* 2004;12:1554–69.
- Livingston EH. Pitfalls in using BMI as a selection criterion for bariatric surgery. *Curr Opin Endocrinol Diabetes Obes.* 2012;19:347–51.
- Cordero P, Li J, Oben JA. Bariatric surgery as a treatment for metabolic syndrome. *J R Coll Physicians Edinb.* 2017;47:364–8.
- Wood KV, Ogden J. Explaining the role of binge eating behaviour in weight loss post bariatric surgery. *Appetite.* 2012;59:177–80.
- Devlin MJ, King WC, Kalarchian MA, et al. Eating pathology and experience and weight loss in a prospective study of bariatric surgery patients: 3-year follow-up. *Int J Eat Disord.* 2016;49:1058–67.
- Gero D, Steinert RE, le Roux CW, et al. Do food preferences change after bariatric surgery? *Curr Atheroscler Rep.* 2017;19:38.
- Rieber N, Giel KE, Meile T, et al. Psychological dimensions after laparoscopic sleeve gastrectomy: reduced mental burden, improved eating behavior, and ongoing need for cognitive eating control. *Surg Obes Relat Dis.* 2013;9:569–73.
- Bartsch M, Langenberg S, Gruner-Labitzke K, et al. Physical activity, decision-making abilities, and eating disturbances in pre- and postbariatric surgery patients. *Obes Surg.* 2016;26:2913–22.
- Dawes AJ, Maggard-Gibbons M, Maher AR, et al. Mental health conditions among patients seeking and undergoing bariatric surgery: a meta-analysis. *JAMA.* 2016;315:150–63.
- Alfonsson S, Weinland-Strandskov S, Sundbom M. Self-reported hedonism predicts 12-month weight loss after Roux-en-Y gastric bypass. *Obes Surg.* 2017;27:2073–8.
- Swiss Study Group for Morbid Obesity and Metabolic Disorders. Directives médicale pour le traitement chirurgical de l'obésité. 1 January 2018. <http://www.smob.ch/fr/component/jdownloads/send/1-root/64-smob-directives-medicales-2018-01-01>

16. von Elm E, Altman DG, Egger M, et al. The strengthening of reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Int J Surg.* 2014;12:1495–9.
17. Hatoum IJ, Kaplan LM. Advantages of percent weight loss as a method of reporting weight loss after Roux-en-Y gastric bypass. *Obesity (Silver Spring).* 2013;21:1519–25.
18. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg.* 2004;240:205–13.
19. Anja Hilbert BT-C. Eating disorder Examination Questionnaire - Deutschsprachige Übersetzung. Münster: Verlag für Psychotherapie, PAG Institut für Psychologie AG; 2006.
20. Berg KC, Peterson CB, Frazier P, et al. Psychometric evaluation of the eating disorder examination and eating disorder examination-questionnaire: a systematic review of the literature. *Int J Eat Disord.* 2012;45:428–38.
21. Morseth MS, Hanvold SE, Ro O, et al. Self-reported eating disorder symptoms before and after gastric bypass and duodenal switch for super obesity—a 5-year follow-up study. *Obes Surg.* 2016;26:588–94.
22. Mond JM, Hay PJ, Rodgers B, et al. Eating Disorder Examination Questionnaire (EDE-Q): norms for young adult women. *Behav Res Ther.* 2006;44:53–62.
23. Sullivan GM, Feinn R. Using effect size-or why the P value is not enough. *J Grad Med Educ.* 2012;4:279–82.
24. Sarwer DB, Lavery M, Spitzer JC. A review of the relationships between extreme obesity, quality of life, and sexual function. *Obes Surg.* 2012;22:668–76.
25. Julia C, Ciangura C, Capuron L, et al. Quality of life after Roux-en-Y gastric bypass and changes in body mass index and obesity-related comorbidities. *Diabetes Metab.* 2013;39:148–54.
26. Peterhansel C, Nagl M, Wagner B, et al. Predictors of changes in health-related quality of life 6 and 12 months after a bariatric procedure. *Obes Surg.* 2017;27:2120–8.
27. Jakobsen JC, Gluud C, Wetterslev J, et al. When and how should multiple imputation be used for handling missing data in randomised clinical trials - a practical guide with flowcharts. *BMC Med Res Methodol.* 2017;17:162.
28. Young-Saver DF, Gornbein J, Starkman S, et al. Handling of missing outcome data in acute stroke trials: advantages of multiple imputation using baseline and postbaseline variables. *J Stroke Cerebrovasc Dis.* 2018;27(12):3662–3669.
29. Rubin DB. Multiple imputation for nonresponse in surveys. New York: Wiley; 1987.
30. Spitznagel MB, Garcia S, Miller LA, et al. Cognitive function predicts weight loss after bariatric surgery. *Surg Obes Relat Dis.* 2013;9:453–9.
31. Conceicao EM, Mitchell JE, Pinto-Bastos A, et al. Stability of problematic eating behaviors and weight loss trajectories after bariatric surgery: a longitudinal observational study. *Surg Obes Relat Dis.* 2017;13:1063–70.
32. Nasirzadeh Y, Kantarovich K, Wnuk S, et al. Binge eating, loss of control over eating, emotional eating, and night eating after bariatric surgery: results from the Toronto Bari-PSYCH Cohort Study. *Obes Surg.* 2018;28:2032–9.
33. Versteegden DPA, Van Himbeeck MJJ, Nienhuijs SW. Improvement in quality of life after bariatric surgery: sleeve versus bypass. *Surg Obes Relat Dis.* 2018;14:170–4.
34. Reynolds CL, Byrne SM, Hamdorf JM. Treatment success: investigating clinically significant change in quality of life following bariatric surgery. *Obes Surg.* 2017;27:1842–8.
35. Jirapinyo P, Abu Dayyeh BK, Thompson CC. Weight regain after Roux-en-Y gastric bypass has a large negative impact on the bariatric quality of life index. *BMJ Open Gastroenterol.* 2017;4:e000153.
36. Chan CW, Leung SF. Validation of the eating disorder examination questionnaire: an online version. *J Hum Nutr Diet.* 2015;28:659–65.
37. Conradt M, Dierk JM, Schlumberger P, et al. A consultation with genetic information about obesity decreases self-blame about eating and leads to realistic weight loss goals in obese individuals. *J Psychosom Res.* 2009;66:287–95.
38. Bedri A, Li R, Haynes M, et al. EarBit: using wearable sensors to detect eating episodes in unconstrained environments. *Proc ACM Interact Mob Wearable Ubiquitous Technol.* 2017;1:1–20.
39. Fairburn CG, Beglin SJ. Assessment of eating disorders: interview or self-report questionnaire? *Int J Eat Disord.* 1994;16:363–70.
40. de Zwaan M, Hilbert A, Swan-Kremeier L, et al. Comprehensive interview assessment of eating behavior 18–35 months after gastric bypass surgery for morbid obesity. *Surg Obes Relat Dis.* 2010;6:79–85.
41. Conceicao E, Mitchell JE, Vaz AR, et al. The presence of maladaptive eating behaviors after bariatric surgery in a cross sectional study: importance of picking or nibbling on weight regain. *Eat Behav.* 2014;15:558–62.
42. Sarwer DB, Dilks RJ, West-Smith L. Dietary intake and eating behavior after bariatric surgery: threats to weight loss maintenance and strategies for success. *Surg Obes Relat Dis.* 2011;7:644–51.
43. Vangoitsenhoven R, Frederiks P, Gijbels B, et al. Long-term effects of gastric bypass surgery on psychosocial well-being and eating behavior: not all that glitters is gold. *Acta Clin Belg.* 2016;71:395–402.
44. Bell ML, Kenward MG, Fairclough DL, et al. Differential dropout and bias in randomised controlled trials: when it matters and when it may not. *BMJ.* 2013;346:e8668.
45. Makaronidis JM, Batterham RL. Obesity, body weight regulation and the brain: insights from fMRI. *Br J Radiol.* 2018;91(1089):20170910.
46. Grilo CM, Henderson KE, Bell RL, et al. Eating disorder examination-questionnaire factor structure and construct validity in bariatric surgery candidates. *Obes Surg.* 2013;23:657–62.
47. Nielsen MS, Christensen BJ, Ritz C, et al. Roux-En-Y gastric bypass and sleeve gastrectomy does not affect food preferences when assessed by an ad libitum buffet meal. *Obes Surg.* 2017;27:2599–605.