



Correction to: Preoperative Weight Loss and Operative Outcome After Laparoscopic Sleeve Gastrectomy

Atsushi Watanabe^{1,2}  · Yosuke Seki² · Hidenori Haruta^{2,3} · Eri Kikkawa² · Kazunori Kasama²

Published online: 11 February 2019
© Springer Science+Business Media, LLC, part of Springer Nature 2019

Correction to: OBES SURG (2017) 27:2515–2521
<https://doi.org/10.1007/s11695-017-2697-3>

In Table 2 on p. 2517 the heading for Group 3 should read as follows:

Group 3
3.1–5% TWL ($n = 61$)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s11695-017-2697-3>

✉ Atsushi Watanabe
atsuwata555@gmail.com

¹ Department of Surgery, Jikei University School of Medicine, 3-25-8 Nishi-shinbashi, Minato-ku, Tokyo 105-8461, Japan

² Weight Loss and Metabolic Surgery Center, Yotsuya Medical Cube, 7-7 Nibancho, Chiyoda-ku, Tokyo 102-0084, Japan

³ Clinical Institute of Digestive Disease Surgical Branch, Jichi Medical University, 3311-1 Yakushiji, Shimotsukeshi, Tochigi 329-0498, Japan