



# TGR5 Protects Against Colitis in Mice, but Vertical Sleeve Gastrectomy Increases Colitis Severity

Darline Garibay<sup>1</sup> · Karolina E. Zaborska<sup>1</sup> · Michael Shanahan<sup>1</sup> · Qiaonan Zheng<sup>1</sup> · Katie M. Kelly<sup>1</sup> · David C. Montrose<sup>2</sup> · Andrew J. Dannenberg<sup>2</sup> · Andrew D. Miller<sup>1</sup> · Praveen Sethupathy<sup>1</sup> · Bethany P. Cummings<sup>1</sup> 

Published online: 8 January 2019

© Springer Science+Business Media, LLC, part of Springer Nature 2019

## Abstract

**Background and Aims** Bariatric surgery, such as vertical sleeve gastrectomy (VSG), is the most effective long-term treatment for obesity. However, there are conflicting reports on the effect of bariatric surgery on inflammatory bowel disease (IBD). Bariatric surgery increases bile acid concentrations, which can decrease inflammation by signaling through the bile acid receptor, TGR5. TGR5 signaling protects against chemically induced colitis in mice. VSG increases circulating bile acid concentrations to increase TGR5 signaling, which contributes to improved metabolic regulation after VSG. Therefore, we investigated the effect of VSG on chemically induced colitis development and the role of TGR5 in this context.

**Methods** VSG or sham surgery was performed in high fat diet-fed male *Tgr5*<sup>+/+</sup> and *Tgr5*<sup>-/-</sup> littermates. Sham-operated mice were food restricted to match their body weight to VSG-operated mice. Colitis was induced with 2.5% dextran sodium sulfate (DSS) in water post-operatively. Body weight, energy intake, fecal scoring, colon histopathology, colonic markers of inflammation, goblet cell counts, and colonic microRNA-21 levels were assessed.

**Results** VSG decreased body weight independently of genotype. Consistent with previous work, genetic ablation of TGR5 increased the severity of DSS-induced colitis. Notably, despite the effect of VSG to decrease body weight and increase TGR5 signaling, VSG increased the severity of DSS-induced colitis. VSG-induced increases in colitis were associated with increased colonic expression of TNF- $\alpha$ , IL-6, MCP-1, and microRNA-21.

**Conclusions** While our data demonstrate that TGR5 protects against colitis, they also demonstrate that VSG potentiates chemically induced colitis in mice. These data suggest that individuals undergoing VSG may be at increased risk for developing colitis; however, further study is needed.

**Keywords** VSG · Colitis · TGR5

---

✉ Bethany P. Cummings  
bpc68@cornell.edu

Darline Garibay  
dg599@cornell.edu

Karolina E. Zaborska  
kz64@cornell.edu

Michael Shanahan  
mts253@cornell.edu

Qiaonan Zheng  
qcz2@cornell.edu

Katie M. Kelly  
kkb65@cornell.edu

David C. Montrose  
dam2040@med.cornell.edu

Andrew J. Dannenberg  
ajdann@med.cornell.edu

Andrew D. Miller  
andrew.miller@cornell.edu

Praveen Sethupathy  
pr46@cornell.edu

<sup>1</sup> Department of Biomedical Sciences, College of Veterinary Medicine, Cornell University, Ithaca, NY 14853, USA

<sup>2</sup> Department of Medicine, Weill Cornell Medical College, New York, NY 10021, USA

## Introduction

Inflammatory bowel disease (IBD) is a chronic inflammatory condition encompassing both Crohn's disease and ulcerative colitis [1]. The mechanisms underlying IBD are complex and involve both genetic and environmental factors [2]. Recent work demonstrates that obesity is positively correlated with IBD, suggesting that obesity may contribute to IBD pathogenesis [3–5]. Obesity is known to induce chronic low-grade inflammation, and it is possible that obesity-related inflammation may promote IBD development and progression [6–8]. While bariatric surgery, such as vertical sleeve gastrectomy (VSG), is arguably the most effective long-term treatment for obesity and produces numerous cardiometabolic benefits [9–11], there is limited and conflicting literature on the effects of bariatric surgery on IBD.

Several small case reports describe the development of inflammatory bowel disease (IBD) after bariatric surgery [12–14]. While some small case studies suggest that bariatric surgery is safe in patients with pre-existing IBD [15, 16], and may even improve IBD-related symptoms [17, 18], the largest case series reported to date demonstrate that bariatric surgery increases the risk of developing IBD [19, 20]. IBD can predispose patients to the development of colorectal cancer [21, 22]. Consistent with this, bariatric surgery has been reported to increase the risk of developing colorectal cancer [23–25]. Clinical studies have analyzed IBD risk in cohorts of patients undergoing a variety of different bariatric surgery types; therefore, the relative impact of different bariatric surgery types on IBD risk is unknown. VSG is the most commonly performed bariatric procedure in the USA [26]. Therefore, we focused our study on the effect of VSG on colitis development and progression in a mouse model of chemically induced colitis.

Bariatric surgery, including VSG, increases circulating bile acid concentrations in humans and rodent models [27–29]. While bile acids are best known as steroid molecules that help digest dietary cholesterol and lipids, they also play an important immunomodulatory role. For example, increased signaling through the bile acid receptor, TGR5, on immune cells reduces inflammatory cytokine secretion [30, 31]. Furthermore, TGR5 has been shown to protect against chemically induced colitis in mice [32]. We and others have previously demonstrated that increased signaling through the bile acid receptor, TGR5, contributes to improved glucose regulation after VSG [27, 33]. Therefore, we investigated the role of TGR5 on the effect of VSG in a chemically induced mouse model of colitis.

## Materials and Methods

### Animals and Diets

Heterozygous *Tgr5*<sup>+/-</sup> breeding pairs generated on a C57BL/6J background were obtained from the UC Davis Knockout

Mouse Project Repository and a breeding colony established. *Tgr5*<sup>+/+</sup> and *Tgr5*<sup>-/-</sup> male littermates were placed on 60% high fat diet at 2 months of age (D12492, Dyets, Bethlehem, PA) for 2 months to produce an obese and insulin resistant phenotype. At 4 months of age, mice underwent sham surgery or VSG, as previously described [27, 34]. The following groups were studied: sham-operated TGR5 wild-type (Sham WT), sham-operated TGR5 knockout (Sham KO), VSG operated TGR5 wild-type (VSG WT), and VSG operated TGR5 knockout (VSG KO). Groups were matched for body weight at the time of surgery. Sham-operated mice were food restricted to match body weight to their respective VSG-operated groups to control for body weight. Food intake and body weight were measured twice per week. 21 days after surgery mice received 2.5% dextran sodium sulfate (DSS) in water for 5 days (TdB Consultancy; Upsala, Sweden). We chose a short low-dose DSS dosing paradigm to study a mouse model with a mild colitis phenotype to ensure that the potential differential effects of VSG and TGR5 were not overwhelmed by the severity of disease. Six-hour fasted mice were euthanized 25 days after surgery by an overdose of pentobarbital (200 mg/kg i.p.). The experimental protocols were approved by the Cornell University Institutional Animal Care and Use Committee.

### Colitis Evaluation

During DSS treatment, fecal scores were performed daily to assess disease severity in vivo. Specifically, diarrhea score was assessed by measuring the softness and appearance of the stool on a scale of 0–3. Bleeding score was measured by detection of heme in stool using the Hemocult Sensa test (Beckman Coulter; Fullerton, CA) or evidence of gross bleeding on a scale from 0 to 3.

At sacrifice, colons were removed, weighed, and length measured. A small full-thickness section of colon was removed from the proximal colon and frozen in liquid nitrogen for immunoblotting analysis. Remaining tissue samples were prepared as spiral “Swiss rolls” and fixed in 4% formalin for 24 h followed by paraffin embedding. Paraffin blocks were sectioned on a microtome at 5 μm and slides stained with hematoxylin and eosin (H&E). Stained sections were analyzed for the presence and severity of colitis in a blinded fashion by a board-certified veterinary pathologist. Briefly, a pathological score from 1 to 4 was given to each sample: (1) none, (2) mild, (3) moderate, and (4) severe. Scoring was based on the presence and severity of colitis, inflammatory infiltration, the presence of erosion/ulcers, and epithelial changes.

### Histologic Quantification of Crypt Density, Crypt Depth, and Goblet Cell Numbers

H&E stained slides were used for histomorphometric measurements of colonic crypt density and crypt depth. For both

measurements, multiple non-overlapping fields were evaluated per slide at 200x magnification. Fields involving at least five adjacent, clearly discernible, individual crypts were enumerated over a measured linear distance for determining crypt density. Crypt depth was assessed by measuring the linear distance of clearly delineated crypts spanning the crypt base and the epithelial surface junction. Linear measurements were made using ImageJ software. Alcian blue stained sections were used to enumerate the number of goblet cells per colonic crypt. Distinguishable goblet cells from clearly delineated crypts from multiple fields at  $\times 200$  magnification were numbered.

### Immunoblotting

For immunoblotting, radio immunoprecipitation assay (RIPA) buffer (10 mM Tris-HCL, pH 7.4, 150 mM NaCL, 0.1% sodium dodecyl sulfate (SDS), 1% Triton-X-100, 1% sodium deoxycholate, 5 mM EDTA, 1 mM NaF, 1 mM sodium orthovanadate and protease inhibitors) was used to lyse tissues in liquid nitrogen. Lysates were separated by centrifugation at 10,000 rpm for 10 min and protein concentrations determined by bicinchoninic acid protein assay (Pierce Chemical, IL). Proteins (40  $\mu$ g) were resolved by SDS-PAGE (10%) and transferred to PVDF membranes. Immunoblots were performed with relevant antibodies and proteins visualized with Luminata Forte (Millipore; Billeorica, MA). The actin antibody (0.002  $\mu$ g/ml) was from Santa Cruz Biotechnology (Dallas, TX), the antibodies for TNF- $\alpha$  (1  $\mu$ g/ml) and MCP-1 (1  $\mu$ g/ml) from Abcam (Cambridge, MA), and the antibody for IL-6 (1  $\mu$ g/ml) from Cell Signaling (Danvers, MA). Horseradish peroxidase (HRP)-conjugated mouse secondary antibody (0.04  $\mu$ g/ml) was from Santa Cruz Biotechnology (Dallas, TX) and horseradish peroxidase (HRP)-conjugated rabbit secondary antibody (0.04  $\mu$ g/ml) was from Bio-Rad Laboratories (Hercules, CA). Chemiluminescence was detected using the Chemidoc MP imaging system and protein expression quantified using ImageLab 5.1 software.

### miR-21 Quantification

Total RNA was extracted from 5-mm sections of frozen proximal colon samples with the Total RNA Purification Kit (Norgen Biotek Corp., Thorold, ON, Canada) according to the manufacturer's instructions. To remove potentially contaminating DSS, purified RNA samples were precipitated with 2.5 M lithium chloride (Thermo Fisher Scientific, Waltham, MA), as described previously [35]. Reverse transcription and real time-PCR were performed with specific TaqMan primers for miR-21 (#000397, Thermo Fisher Scientific, Waltham, MA) and U6 snRNA (#001973, Thermo Fisher Scientific, Waltham, MA). Fold change of expression was calculated based on the  $\Delta\Delta C_t$  method.

### Statistics and Data Analysis

Data are presented as mean  $\pm$  SEM. All statistical analysis was performed with GraphPad Prism 6.00 for windows (GraphPad Software, San Diego, CA). Data were analyzed by two-factor ANOVA with Bonferroni's post-test unless otherwise indicated. Differences were indicated by an  $\alpha$  of 0.05.

## Results


### VSG Decreases Body Weight Independently of TGR5 in DSS-Treated Mice

To test the impact of TGR5 and VSG on colitis development, we treated sham and VSG-operated *Tgr5*<sup>+/+</sup> and *Tgr5*<sup>-/-</sup> mice with 2.5% DSS for 5 days, starting 21 days after surgery. Sham-operated mice were food restricted to match body weight to VSG-operated mice to control for this variable. Therefore, cumulative energy intake, body weight, and adiposity did not differ between sham and VSG-operated groups throughout study (Fig. 1a–c). VSG lowered body weight relative to baseline in both VSG WT and VSG KO groups ( $P < 0.05$ ; Fig. 1b). Energy intake, body weight, and adiposity did not differ between genotype in VSG or sham-operated mice throughout study (Fig. 1a–c). This is consistent with previous work showing that VSG decreases body weight independently of TGR5 in non-DSS treated mice [27].

### Genetic Ablation of TGR5 and VSG Potentiates Histologic Injury Following DSS-Induced Colitis

To assess disease severity in vivo, diarrhea and colitis-associated bleeding were scored daily during DSS administration. Diarrhea score did not differ between Sham WT and VSG WT (Fig. 2a). However, diarrhea score was elevated 2.5-fold in VSG KO compared with Sham KO on day 4 of DSS treatment ( $P < 0.05$ ; Fig. 2a). Bleeding score, colon length, and colon weight did not differ between groups (Fig. 2b–d).

Histologic examination of H&E stained colons revealed a trend for a higher incidence of erosion, hyperplasia, lymphocytic infiltration, and colitis in VSG WT compared with Sham WT and Sham KO compared with Sham WT (Fig. 3a; Table 1). Specifically, 63% of Sham WT colons were positive for histopathological changes consistent with colitis, compared with 86–90% in all other groups (Table 1). Interestingly, the incidence of neutrophilic infiltration was significantly elevated in VSG KO compared with Sham KO, but not in VSG WT compared with Sham WT ( $P < 0.05$ ; Table 1). To gain more detailed insight into the histologic changes induced by VSG and genetic ablation of TGR5, we quantified colonic crypt depth, crypt




**Fig. 1** VSG decreases body weight independently of TGR5. Cumulative energy intake (a) and body weight (b). Adipose depot weight (Subcutaneous adipose tissue (Sub), mesenteric adipose tissue (Mes), epididymal adipose tissue (Epi), retroperitoneal adipose tissue (RP), and

brown adipose tissue (BAT)) (c). Data are presented as mean  $\pm$  SEM.  $n = 7$ –10 per group. \* $P < 0.05$  for all groups compared with baseline body weight by two-factor ANOVA

density, and goblet cell number per crypt. Crypt density and crypt depth did not differ between groups (Fig. 3b, c). There was a trend for goblet cell number per crypt to be elevated in VSG compared with Sham in both  $Tgr5^{+/+}$  and  $Tgr5^{-/-}$  mice; however, this did not reach significance (Fig. 3d). If we set a threshold of average goblet cell count per crypt at 10 (based on the Sham WT data), only 50% of mice exceeded this threshold in the sham-operated group whereas 75% of mice exceeded this threshold in the VSG WT groups. Histopathologic scoring revealed that colitis severity was higher in VSG KO compared with Sham WT ( $P < 0.05$ ; Fig. 3e). Direct comparison of VSG WT with Sham WT revealed a 45% increase in colitis severity score in VSG WT compared with Sham WT ( $P < 0.05$ ; Fig. 3e). Direct comparison of Sham WT with Sham KO revealed that genetic ablation of TGR5 increases colitis severity score in sham-operated mice ( $P < 0.05$ ; Fig. 3e). Colitis severity score did not differ between VSG WT and VSG KO (Fig. 3e). Overall, these data suggest that genetic ablation of TGR5 and VSG exacerbate histologic evidence of colitis and its severity.

### Genetic Ablation of TGR5 and VSG Enhance Colonic Inflammation

To quantify the degree of colonic inflammation, immunoblotting was performed on colon tissue for the following inflammatory markers: monocyte chemoattractant protein 1 (MCP-1), tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) and interleukin-6 (IL-6). MCP-1, TNF- $\alpha$ , and IL-6 expression were higher in VSG WT compared with Sham WT, demonstrating that VSG increases colonic inflammation in DSS-treated mice ( $P < 0.05$ ; Fig. 4a–d). MCP-1 was elevated in VSG KO compared with Sham KO ( $P < 0.05$ ; Fig. 4a, b). Furthermore, MCP-1, TNF- $\alpha$ , and IL-6 expression were elevated in Sham KO compared with Sham WT, demonstrating that TGR5 ablation augments DSS-induced colonic inflammation in mice ( $P < 0.05$ ; Fig. 4a–d). Finally, IL-6 and MCP-1 expression were elevated in VSG KO compared with VSG WT, suggesting that TGR5 ablation aggravates DSS-induced colonic inflammation after VSG ( $P < 0.05$ ; Fig. 4a, b, d). Thus, these data are consistent with the histologic finding that Sham KO and VSG-operated mice had higher colitis severity. Unlike our



**Fig. 2** In vivo assessment of DSS-induced disease severity. Diarrhea score (a) and fecal blood score (b). Colon weight (c) and colon length (d). Data are presented as mean  $\pm$  SEM.  $n = 7$ –10 per group. \* $P < 0.05$  for VSG KO compared with Sham KO by two-factor ANOVA

histologic analysis, these data suggest that VSG in combination with genetic ablation of TGR5 has an additive effect on indices of colonic inflammation. Together, these data demonstrate that genetic ablation of TGR5 and VSG increases colonic inflammation.

### VSG Enhances Colonic microRNA-21 Expression


MicroRNAs (miRNAs) are critical regulators of gastrointestinal health through the post-transcriptional regulation of gene expression. In particular, colonic miR-21 expression has been shown to be upregulated in patients with Crohn's disease and ulcerative colitis [36–39]. Furthermore, recent studies show that genetic ablation of miR-21 protects against DSS-induced colitis [38, 40]. Therefore, we hypothesized that miR-21 would be aberrantly elevated in the context of TGR5 ablation and/or VSG, both of which aggravate DSS-induced colitis according to our data. To test this hypothesis, we carried out real-time quantitative PCR and found that VSG significantly increased colonic expression of miR-21 (3-fold,  $P < 0.05$ ; Fig. 4e). On the other hand, genetic ablation of TGR5 did not significantly increase miR-21 (Fig. 4e). Therefore, increased colonic miR-21 expression may contribute to the effect of VSG to promote DSS-induced colitis but

does not appear to be related to the effect of genetic ablation of TGR5 to increase colitis severity.

### Conclusion

Herein, we report that VSG is associated with worse DSS-induced colitis in mice. We have previously reported that increased TGR5 signaling contributes to improved glucose regulation after VSG, and our findings herein demonstrate that TGR5 protects against colitis. This is consistent with previous reports that genetic ablation of TGR5 promotes DSS-induced colitis in mice [30, 41]. Nevertheless, our data demonstrate that VSG promotes DSS-induced colitis in mice despite increased TGR5 signaling. Our findings are consistent with data in human patients reporting worsening of colitis after bariatric surgery and may help explain findings of de novo IBD in patients that undergo bariatric surgery [12].

There is conflicting information in the literature regarding the impact of bariatric surgery on IBD. There are reports that bariatric surgery is effective in IBD patients and can improve their IBD status [16, 17]. However, IBD increases the complexity and post-operative care of patients after bariatric surgery [42]. For example, chronic steroid administration, such



**Fig. 3** VSG and genetic ablation of TGR5 potentiate histomorphologic signs of DSS-induced colitis. Representative H&E stained colon sections (a). Quantification of colonic crypt depth (b), crypt density (c), number of goblet cells per crypt (d), and colitis severity score (e). Data are presented

as mean  $\pm$  SEM.  $n = 7$ – $10$  per group. \* $P < 0.05$  Sham WT compared with VSG WT and Sham KO by Student's  $t$  test. <sup>+</sup> $P < 0.05$  Sham WT compared with VSG KO by two-factor ANOVA

as that used for IBD treatment, is associated with increased morbidity and mortality after bariatric surgery [43]. In addition, recent reports suggest that bariatric surgery can lead to the development of de novo IBD. One retrospective study reports 44 cases of de novo IBD from electronic medical records from two institutions over the past 20 years, with an incidence rate of 26.7 cases per 100,000 persons per year [19]. This study demonstrates a higher incidence of IBD development in patients with bariatric surgery compared with the normal population. Our data suggest that VSG worsens the severity of experimental colitis which may explain the

increased incidence of IBD development observed in human patients after bariatric surgery.


It is likely that the impact of bariatric surgery on colitis varies between surgery types. One previous study has investigated the effect of colitis in a rodent model of bariatric surgery. Interestingly, they found that duodenojejunal bypass decreased colitis severity compared to both sham and VSG-operated mice [44]. Our data also demonstrate that VSG exacerbates colitis in a DSS-induced colitis model. Thus, it appears that there may be differences between surgery types on the impact on colitis development and/or progression and that

**Table 1** Effect of VSG and genetic ablation of TGR5 on incidence of colitis, inflammatory infiltrates, and hyperplasia

% of mice with	Sham WT (%)	VSG WT (%)	Sham KO (%)	VSG KO (%)
Colitis	63	89	90	86
Erosion/ulcer	25	56	50	57
Neutrophilic infiltration	38	44	30	86*
Lymphocytic infiltration	63	89	90	86
Hyperplasia	13	44	30	57

Percent of mice with colitis, erosion/ulcer, neutrophilic infiltration, lymphocytic infiltration, and hyperplasia.  $n = 7$ – $10$  per group

\* $P < 0.05$  VSG KO compared with Sham KO by chi-square test



**Fig. 4** VSG and genetic ablation of TGR5 increase colonic inflammation. Representative immunoblots for monocyte chemoattractant protein 1 (MCP-1), tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), interleukin 6 (IL-6), and  $\beta$ -actin (a). MCP-1 relative to  $\beta$ -actin (b), TNF- $\alpha$  relative to  $\beta$ -actin (c), and IL-6 relative to  $\beta$ -actin (d). Results were quantified in densitometric units.  $n = 6$

per group. Colonic miR-21 expression (e).  $n = 4-7$  per group. Data are presented as mean  $\pm$  SEM. \* $P < 0.05$  Sham WT compared with VSG WT, # $P < 0.05$  Sham KO compared with VSG KO, & $P < 0.05$  Sham KO compared with Sham WT, + $P < 0.05$  VSG KO compared with VSG WT by two-factor ANOVA

VSG may have a greater impact on IBD than other bariatric surgery types.

One potential contributor to the effect of VSG to promote the development of colitis is through changes in gut microbial composition. Bariatric surgery results in dramatic alterations in gut microbial composition in both rodent models and human patients [45–49]. Surprisingly, these gut microbial shifts are remarkably similar between vastly different bariatric surgery types [45–47, 49, 50], demonstrating that the relevant microbial changes occur irrespective of differences in the specific gastrointestinal anatomic alterations. These shifts in the gut microbiome have been suggested to play an important role in the metabolic benefits of bariatric surgery. For example, transplantation of gut microbiota derived from RYGB-operated humans or mice to germ-free mice produces lower adiposity than control transplants [47, 50]. However, the changes in gut microbial composition include increased relative abundance of bacterial genera considered pathogenic and associated with the development of colitis. Indeed, these gut microbial shifts are remarkably similar to those observed in human IBD patients. One of the most consistently reported gut microbial changes after various types of bariatric surgery in both humans and rodents is an increase in the class, Gammaproteobacteria, and its associated family, Enterobacteriaceae [45, 47–50]. We have previously reported that our mouse VSG model exhibits a significant increase

in the relative abundance of Gammaproteobacteria and Enterobacteriaceae up to 6.5 months after surgery [27]. Enterobacteriaceae are increased in patients with IBD and in mouse models of IBD [51]. In particular, *Escherichia* and *Shigella*, subdivisions of Enterobacteriaceae, are increased in patients with IBD and treatment with anti-inflammatory drugs decreases the relative abundance of these genera [52, 53]. *Escherichia* and *Shigella* are elevated in human patients and rodents, including our rodent model, after bariatric surgery [45, 47–50]. Therefore, while metabolically beneficial, these gut microbial changes may contribute to increased colitis risk after bariatric surgery.

Our data also reveal that increases in colonic miR-21 expression may contribute to the effect of VSG to promote colitis development. MiR-21 is among the most significantly upregulated miRNAs in the colon of adult and pediatric patients with Crohn's disease [37, 38]. Other studies have demonstrated that colonic miR-21 levels are elevated in ulcerative colitis and that this may increase risk for colitis-associated colon cancer [36, 39, 54]. Moreover, it has been reported that the levels of miR-21 in colonic CD3+ T cells are rescued during remission of ulcerative colitis [40]. Very recent studies show that genetic ablation of miR-21 induces a healthy shift in the gut microbiome and protects against DSS-induced colitis [38, 40]. Therefore, it is possible that post-operative increases

in colonic miR-21 expression contribute to pathogenic shifts in gut microbial composition to promote colitis development; however, further work is needed to evaluate this connection.

Our work demonstrates that VSG potentiates DSS-induced colitis in mice. The current results are consistent with prior reports in humans demonstrating worsening of IBD symptoms in patients after surgery and may help explain de novo colitis development in patients who undergo bariatric surgery. Further work is needed to confirm these findings in human patients and differentiate the relative impact of different bariatric surgery types on IBD risk in humans. Overall, these data suggest that caution should be exercised in performing VSG on patients with increased risk of colitis development.

**Acknowledgements** We thank the Animal Health and Diagnostic Center Histopathology Core and Martin Slade for preparation of samples for H&E.

**Funding** This research was supported by NIH grant R21CA195002.

### Compliance with Ethical Standards

The experimental protocols were approved by the Cornell University Institutional Animal Care and Use Committee.

**Conflict of Interest** The authors declare that they have no conflict of interest.

**Ethical Approval** All experimental protocols were approved by the Cornell University Institutional Animal Care and Use Committee, and all applicable institutional and/or national guidelines for the care and use of animals were followed.

**Informed Consent** Does not apply.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

### References

- Wehkamp J, Götz M, Herrlinger K, et al. Inflammatory bowel disease: Crohn's disease and ulcerative colitis. *Dtsch Arztebl Int*. 2016;113(5):72–82.
- Biasi F, Leonarduzzi G, Oteiza PI, et al. Inflammatory bowel disease: mechanisms, redox considerations, and therapeutic targets. *Antioxid Redox Signal*. 2013;19(14):1711–47.
- Singh S, Dulai PS, Zarrinpar A, et al. Obesity in IBD: epidemiology, pathogenesis, disease course and treatment outcomes. *Nat Rev Gastroenterol Hepatol*. 2017;14(2):110–21.
- Harper JW, Zisman TL. Interaction of obesity and inflammatory bowel disease. *World J Gastroenterol*. 2016;22(35):7868–81.
- Nic Suibhne T, Raftery TC, McMahon O, et al. High prevalence of overweight and obesity in adults with Crohn's disease: associations with disease and lifestyle factors. *J Crohn's Colitis*. 2013;7(7):e241–8.
- Winer DA, Luck H, Tsai S, et al. The intestinal immune system in obesity and insulin resistance. *Cell Metab*. 2016;23(3):413–26.
- Pereira SS, Alvarez-Leite JI. Low-grade inflammation, obesity, and diabetes. *Curr Obes Rep*. 2014;3(4):422–31.
- Monteiro R, Azevedo I. Chronic inflammation in obesity and the metabolic syndrome. *Mediat Inflamm*. 2010;2010(289645):1–10.
- Buchwald H, Avidor Y, Braunwald E, et al. Bariatric surgery: a systematic review and meta-analysis. *JAMA*. 2004;292(14):1724–37.
- Pories WJ, Swanson MS, MacDonald KG, et al. Who would have thought it? An operation proves to be the most effective therapy for adult-onset diabetes mellitus. *Ann Surg*. 1995;222(3):339–50. discussion 50–2
- Flores L, Vidal J, Canivell S, et al. Hypertension remission 1 year after bariatric surgery: predictive factors. *Surg Obes Relat Dis*. 2014;10(4):661–5.
- Ahn LB, Huang CS, Forse RA, et al. Crohn's disease after gastric bypass surgery for morbid obesity: is there an association? *Inflamm Bowel Dis*. 2005;11(6):622–4.
- Dodell GB, Albu JB, Attia L, et al. The bariatric surgery patient: lost to follow-up; from morbid obesity to severe malnutrition. *Endocr Pract*. 2012;18(2):e21–5.
- Janczewska I, Nekzada Q, Kapraali M. Crohn's disease after gastric bypass surgery. *BMJ Case Rep*. 2011;2011:bcr0720103168.
- Aelfers S, Janssen IMC, Aarts EO, et al. Inflammatory bowel disease is not a contraindication for bariatric surgery. *Obes Surg*. 2018;28(6):1681–7.
- Keidar A, Hazan D, Sadot E, et al. The role of bariatric surgery in morbidly obese patients with inflammatory bowel disease. *Surg Obes Relat Dis*. 2015;11(1):132–6.
- Aminian A, Andalib A, Ver MR, et al. Outcomes of bariatric surgery in patients with inflammatory bowel disease. *Obes Surg*. 2016;26(6):1186–90.
- Colombo F, Rizzi A, Ferrari C, et al. Bariatric surgery in patients with inflammatory bowel disease: an accessible path? Report of a case series and review of the literature. *J Crohn's Colitis*. 2015;9(2):185–90.
- Braga Neto MB, Gregory M, Ramos GP, et al. De-novo inflammatory bowel disease after bariatric surgery: a large case series. *J Crohn's Colitis*. 2018;12(4):452–7.
- Ungaro R, Fausel R, Chang HL, et al. Bariatric surgery is associated with increased risk of new-onset inflammatory bowel disease: case series and national database study. *Aliment Pharmacol Ther*. 2018;47(8):1126–34.
- Beaugerie L, Itzkowitz SH. Cancers complicating inflammatory bowel disease. *N Engl J Med*. 2015;372(15):1441–52.
- Fornaro R, Caratto M, Caratto E, et al. Colorectal cancer in patients with inflammatory bowel disease: the need for a real surveillance program. *Clin Colorectal Cancer*. 2016;15(3):204–12.
- Derogar M, Hull MA, Kant P, et al. Increased risk of colorectal cancer after obesity surgery. *Ann Surg*. 2013;258(6):983–8.
- Kant P, Sainsbury A, Reed KR, et al. Rectal epithelial cell mitosis and expression of macrophage migration inhibitory factor are increased 3 years after Roux-en-Y gastric bypass (RYGB) for morbid obesity: implications for long-term neoplastic risk following RYGB. *Gut*. 2011;60(7):893–901.
- Tao W, Konings P, Hull MA, et al. Colorectal cancer prognosis following obesity surgery in a population-based cohort study. *Obes Surg*. 2017;27(5):1233–9.
- Esteban Varela J, Nguyen NT. Laparoscopic sleeve gastrectomy leads the U.S. utilization of bariatric surgery at academic medical centers. *Surg Obes Relat Dis*. 2015;11(5):987–90.
- McGavigan AK, Garibay D, Henseler ZM, et al. TGR5 contributes to glucoregulatory improvements after vertical sleeve gastrectomy in mice. *Gut*. 2017;66:226–34.
- Patti ME, Houten SM, Bianco AC, et al. Serum bile acids are higher in humans with prior gastric bypass: potential contribution to

- improved glucose and lipid metabolism. *Obesity* (Silver Spring). 2009;17(9):1671–7.
29. Steinert RE, Peterli R, Keller S, et al. Bile acids and gut peptide secretion after bariatric surgery: a 1-year prospective randomized pilot trial. *Obesity* (Silver Spring). 2013;21(12):E660–8.
  30. Guo C, Qi H, Yu Y, et al. The G-protein-coupled bile acid receptor Gpbar1 (TGR5) inhibits gastric inflammation through antagonizing NF- $\kappa$ B signaling pathway. *Front Pharmacol*. 2015;6:287.
  31. Fiorucci S, Cipriani S, Mencarelli A, et al. Counter-regulatory role of bile acid activated receptors in immunity and inflammation. *Curr Mol Med*. 2010;10(6):579–95.
  32. Cipriani S, Mencarelli A, Chini MG, Distrutti E, Renga B, Bifulco G, Baldelli F, Donini A, Fiorucci S. The bile acid receptor GPBAR-1 (TGR5) modulates integrity of intestinal barrier and immune response to experimental colitis. *PLoS One*. 2011;6(10):e25637.
  33. Ding L, Sousa KM, Jin L, et al. Vertical sleeve gastrectomy activates GPBAR-1/TGR5 to sustain weight loss, improve fatty liver, and remit insulin resistance in mice. *Hepatology*. 2016;64(3):760–73.
  34. Garibay D, McGavigan AK, Lee SA, et al. Beta-cell glucagon-like peptide-1 receptor contributes to improved glucose tolerance after vertical sleeve gastrectomy. *Endocrinology*. 2016;157(9):3405–9.
  35. Viennois E, Chen F, Laroui H, et al. Dextran sodium sulfate inhibits the activities of both polymerase and reverse transcriptase: lithium chloride purification, a rapid and efficient technique to purify RNA. *BMC Res Notes*. 2013;6:360.
  36. Ludwig K, Fassan M, Mescoli C, et al. PDCD4/miR-21 dysregulation in inflammatory bowel disease-associated carcinogenesis. *Virchows Arch*. 2013;462(1):57–63.
  37. Peck BC, Weiser M, Lee SE, et al. MicroRNAs classify different disease behavior phenotypes of Crohn's disease and may have prognostic utility. *Inflamm Bowel Dis*. 2015;21(9):2178–87.
  38. Shi C, Liang Y, Yang J, et al. MicroRNA-21 knockout improve the survival rate in DSS induced fatal colitis through protecting against inflammation and tissue injury. *PLoS One*. 2013;8(6):e66814.
  39. Yang Y, Ma Y, Shi C, et al. Overexpression of miR-21 in patients with ulcerative colitis impairs intestinal epithelial barrier function through targeting the Rho GTPase RhoB. *Biochem Biophys Res Commun*. 2013;434(4):746–52.
  40. Ando Y, Mazzurana L, Forkel M, et al. Downregulation of MicroRNA-21 in colonic CD3+ T cells in UC remission. *Inflamm Bowel Dis*. 2016;22(12):2788–93.
  41. Biagioli M, Carino A. The bile acid receptor GPBAR1 regulates the M1/M2 phenotype of intestinal macrophages and activation of GPBAR1 rescues mice from murine colitis. *J Immunol*. 2017;199(2):718–33.
  42. Boutros M, Maron D. Inflammatory bowel disease in the obese patient. *Clin Colon Rectal Surg*. 2011;24(4):244–52.
  43. Kaplan GG. The global burden of IBD: from 2015 to 2025. *Nat Rev Gastroenterol Hepatol*. 2015;12(12):720–7.
  44. Li S, Vinci A, Behnsen J, et al. Bariatric surgery attenuates colitis in an obese murine model. *Surg Obes Relat Dis*. 2017;13(4):661–8.
  45. Cummings BP, Bettaieb A, Graham JL, et al. Bile-acid-mediated decrease in endoplasmic reticulum stress: a potential contributor to the metabolic benefits of ileal interposition surgery in UCD-T2DM rats. *Dis Model Mech*. 2013;6(2):443–56.
  46. Li JV, Ashrafian H, Bueter M, et al. Metabolic surgery profoundly influences gut microbial-host metabolic cross-talk. *Gut*. 2011;60(9):1214–23.
  47. Liou AP, Paziuk M, Luevano Jr JM, et al. Conserved shifts in the gut microbiota due to gastric bypass reduce host weight and adiposity. *Sci Transl Med*. 2013;5(178):178ra41.
  48. Ryan KK, Tremaroli V, Clemmensen C, et al. FXR is a molecular target for the effects of vertical sleeve gastrectomy. *Nature*. 2014;509:183–8.
  49. Zhang H, DiBaise JK, Zuccolo A, et al. Human gut microbiota in obesity and after gastric bypass. *Proc Natl Acad Sci U S A*. 2009;106(7):2365–70.
  50. Tremaroli V, Karlsson F, Werling M, et al. Roux-en-Y gastric bypass and vertical banded gastroplasty induce long-term changes on the human gut microbiome contributing to fat mass regulation. *Cell Metab*. 2015;22(2):228–38.
  51. Lupp C, Robertson ML, Wickham ME, et al. Host-mediated inflammation disrupts the intestinal microbiota and promotes the overgrowth of Enterobacteriaceae. *Cell Host Microbe*. 2007;2(3):204.
  52. Benjamin JL, Hedin CR, Koutsoumpas A, et al. Smokers with active Crohn's disease have a clinically relevant dysbiosis of the gastrointestinal microbiota. *Inflamm Bowel Dis*. 2012;18(6):1092–100.
  53. Frank DN, St Amand AL, Feldman RA, et al. Molecular-phylogenetic characterization of microbial community imbalances in human inflammatory bowel diseases. *Proc Natl Acad Sci U S A*. 2007;104(34):13780–5.
  54. Shi C, Yang Y, Xia Y, et al. Novel evidence for an oncogenic role of microRNA-21 in colitis-associated colorectal cancer. *Gut*. 2016;65(9):1470–81.