

Background: Maxillary anterior teeth play an important role in aesthetics, phonetics, and mastication. Because of the aesthetic and functional significance, it is important to evaluate their position in association with surrounding structures in 3-D before initiating orthodontic treatment. With the advent of cone beam computed tomography (CBCT) in orthodontic diagnosis, it is now possible to identify these structures. One such structure is the incisive canal. Uncertainty in locating the canal can cause root resorption postorthodontic retraction of anterior maxillary teeth.

Objective(s): The aim of this study was to calculate the relative distance between the incisive canal and maxillary central incisor using CBCT and utilize the results for orthodontic treatment planning.

Study Design: All quantitative measurements between the incisive canal and both maxillary central incisors were performed using CBCT on 61 patients. The anteroposterior measurements were taken on both sides, and the average of both values was considered for the statistical analysis. The Steiner cephalometric analysis was performed to select patients with skeletal and dental Class I relationship. All linear measurements were performed on the axial plane at 3 different vertical reference points located on the sagittal plane: (1) the palatal opening of the incisive canal (opening level, P1), (2) midlevel between the opening level and the root apex of the maxillary central incisors (midlevel, P2), and (3) the root apex of the maxillary central incisors (root apex level, P3).

Results: The average anteroposterior distance between the maxillary central incisor roots and the incisive canal was approximately 56 mm.

Discussion/Conclusions: The results of our study could be clinically helpful to plan orthodontic treatment that requires significant retraction of maxillary incisors. The authors support the idea of a pretreatment CBCT in cases that require maximum anterior retraction in the maxilla.

Acknowledgments

I would like to acknowledge Ms. Shuying Jiang (Statistician at RSDM), who assisted with the statistics.

References

- Zachrisson BU. Esthetic factors involved in anterior tooth display and the smile; vertical dimension. *J Clin Orthod*. 1998;32:432-445.
- Riedel RA. Esthetics and its relation to orthodontic therapy. *Angle Orthod*. 1950;20:168-178.
- Friedrich RE, Laumann F, Zrnc T, Assaf AT. Nasopalatine canal in adults on cone beam computed tomograms: a clinical study and review of the literature. *In Vivo*. 2015;29:467-486.
- Chung CJ, Choi YJ, Kimc K-H. Approximation and contact of the maxillary central incisor roots with the incisive canal after maximum retraction with temporary anchorage devices: report of 2 patients. *Am J Orthodont Dentofac Orthoped*. 2016;149:9-10.

CYCLICAL CHANGE IN THE APPEARANCE OF FLORID CEMENTO-OSSEOUS DYSPLASIA OVER 15 YEARS: A CASE REPORT. R.C.

GILLIES^{A,B}, C.P. WOO^{A,B}, L. LEE^{A,B}, E.W.N. LAM^{A,B}. ^A UNIVERSITY OF TORONTO, TORONTO, ON, CANADA, ^B BLOOR WEST DENTAL GROUP, TORONTO, ON; CANADA

Background: Florid cemento-osseous dysplasia is a dynamic dysplastic bone abnormality where normal bone is replaced with fibrous tissue and various patterns of nonmature bone. Over time, these changes can be appreciated on imaging with the appearance of an early radiolucent entity becoming a mixed radiopaque/radiolucent entity and finally a predominantly radiopaque entity with a radiolucent rim of variable width.

Discussion/Conclusions: We present a case of florid cemento-osseous dysplasia, which did not follow a typical pattern, with imaging spanning 15 years in a single patient. The patient, a 41-year-old female, first presented to our clinic in 2008. Previous images from 2002 demonstrated typical patterns of florid cemento-osseous dysplasia in the mandible; radiolucent foci; and mixed radiolucent/radiopaque foci with epicenters located inferior to the apices of the teeth. In 2006, the pattern then changed to a wholly radiolucent one, and we interpreted this as being a simple bone cyst developing within the mixed radiopaque/radiolucent focus in the left mandible. Later images in 2013 showed peripheral new bone deposition; an appearance that we interpreted as being consistent with healing. Similar changes in the foci in the right mandible were also seen between 2013 and 2016. The left mandible showed fluctuations between both bone resorption and deposition over the years of imaging. External root resorption of the mandibular teeth also occurred, and these eventually involved the apical one-third of the affected teeth—a process that is rare, but occasionally seen.

This case illustrates the dynamic clinical behavior that can occur with florid cemento-osseous dysplasia. It is important to recognize the range of variations in the presentation of this dysplastic bone entity and to correlate imaging findings with a clinical history to avoid misdiagnosis and mismanagement. Monitoring in certain cases may be necessary to confirm an initial interpretation of florid cemento-osseous dysplasia.

Reference

- White SC, Pharoah MJ. *Oral Radiology: Principles and Interpretation*. 7th ed. St. Louis, MO: Elsevier Mosby; 2014:408-414.

DENTAL IMAGING DILEMMAS IN KLIPPEL-FEIL ANOMALY. KENNETH ABRAMO-VITCH^{A,B}, B. PASS^{A,B}, D.D. RICE^{A,B}. ^A LOMA

LINDA UNIVERSITY, LOMA LINDA, CA, ^B HOWARD UNIVERSITY, WASHINGTON, DC

Background: Because of the broad range of findings, Klippel-Feil syndrome has also been referred to as *Klippel-Feil sequence* and *Klippel-Feil anomaly*. It is classified into 3 types that differ in degree and location of vertebral fusion. Another contemporary classification also includes patterns of inheritance.

Case Report: A 19-year-old male needed panoramic imaging to evaluate his third molars. However, because of Klippel-Feil disfigurement, his head and neck posture did not fit into various panoramic imaging units. With an altered oblique head positioning in a cone beam computed tomography (CBCT) scanner (NewTom VGi), his jaws were scanned. The CBCT volumetric data were then realigned and reconstructed in CBCT viewing software (InVivoDental 6.0). These reconstructions were appropriate to assess the indicated dental status and to update the assessment of the cervical vertebral development—a limiting condition of Klippel-Feil syndrome.