

lead to disfigurement and loss or altered mandibular function, if not treated.

References

- 1 AFIP Archives. CME. Radiographics. 2000;20:5.
- 2 Miller TT. Bone tumors and tumorlike conditions: analysis with conventional radiography. Radiology. 2008;246(3):662-674.
- 3 Woertler K. Benign bone tumors and tumor-like lesions: value of cross-sectional imaging. Eur Radiol. 2003;13:1820-1835.
- 4 Costelloe CM, Churang HH, Madewell J. FDG PET-CT of primary bone tumors. Am J Roentgenol. 2014;202:6.

THE BEATEN-COPPER PATTERN ON CONE BEAM COMPUTED TOMOGRAPHY. P.M. DE MOURA, D.J. FLINT, M.K. NAIR, H. LIANG. TEXAS A&M UNIVERSITY, DALLAS, TX

Background: Convolutional markings are inner-table indentations that conform to the cerebral surface of growing brain in infants. If they are pronounced over the more anterior parts of the skull, this is referred to as a *beaten-copper skull* (BCS). The significance of BCS has been debated, and is generally considered a normal finding in children. However, a diffuse beaten-copper pattern has been shown to be associated with raised intracranial pressure (ICP).

Objective(s): The aim of this study was to increase awareness of the features of BCS on cone beam computed tomography (CBCT).

Results: A CBCT study of a 4-year-old male was referred for interpretation of radiolucent spots in his head and skull. Findings included diffuse scalloping of the inner table of the anterior and posterior cranial compartments, with localized thinning of the diploe in over 50% of the visualized portions of the frontal, parietal, and occipital bones. No sellar erosion or suture diastases were present. The sagittal suture was not included in the field of view; however, coronal, lambdoidal, frontosphenoidal, frontonasal, sphenosquamous, sphenoparietal, occipitomastoid sutures, and the spheno-occipital synchondrosis were visualized and perceived as not fused.

Discussion/Conclusions: The appearance of BCS is age dependent in both normal children and those with craniosynostoses. Clinically, children with craniosynostoses should be managed by a multispecialty team providing interdisciplinary care; they have unique oral health and craniofacial growth problems and may require CBCT. The wormian bones are considered abnormal or clinically significant when radiolucencies are greater than 10 in number, measuring over 6 × 4 mm, and presenting with a general mosaic pattern. The majority of patients with craniosynostoses who have elevated ICP have no related symptoms. The associated finding of BCS using CBCT may be incidental; however, physician referral to further evaluate patients for ICP may be warranted in addition to follow-up over a period of time.

References

- 1 Tuite GF, Evanson J, Chong WK, et al. The beaten copper cranium: a correlation between intracranial pressure, cranial radiographs, and computed tomographic scans in children with craniosynostosis. Neurosurgery. 1996;39:691-699.

- 2 Glass RB, Fernbach SK, Norton KI, Choi PS, Naidich TP. The infant skull: a vault of information. Radiographics. 2004;24:507-522.
- 3 van der Meulen J, van der Vlugt J, Okkerse J, Hofman B. Early beaten-copper pattern: its long-term effect on intelligence quotients in 95 children with craniosynostosis. J Neurosurg Pediatr. 2008;1:25-30.
- 4 Vargervik K, Rubin MS, Grayson BH, et al. Parameters of care for craniosynostosis: dental and orthodontic perspectives. Am J Orthod Dentofacial Orthop. 2012;141:S68-S73.
- 5 Cremin B, Goodman H, Spranger J, et al. Wormian bones in osteogenesis imperfecta and other disorders. Skeletal Radiol. 1982;8:35-38.

THE ROLE OF MONITORS IN THE VISUALIZATION AND ASSESSMENT OF THE INFERIOR ALVEOLAR CANAL. J. ORGILL, S. ANAMALI, S. VIJAYAN, V. ALLAREDDY. THE UNIVERSITY OF IOWA. IOWA CITY, IA

Background: The inferior alveolar canal (IAC) is a familiar landmark for dentists. Clear visualization of the IAC and its relationship with developing or impacted mandibular third molars is especially important. Cone beam computed tomography (CBCT) has improved the ability for more accurate assessment of the IAC.

Objective(s): The aim of this study was to determine if there is a difference in the ability to appropriately assess the third molar-IAC relationship between 3 different monitor types.

Study Design: In all, 105 scans were randomized and evaluated by 2 calibrated and masked evaluators. Evaluation was performed on 3 different monitors: BARCO 3 MP medical-grade monitor, a prototype BARCO 2 MP monitor, and DELL ultra-sharp monitor. Evaluations were completed in a dimly lit area. The luminance and ambient light were measured using a light meter. All 3 monitors were placed in same position for the evaluators and were adjusted such that the luminance was the same. The gold standard was established by 2 board-certified oral and maxillofacial radiologists and 1 oral and maxillofacial radiology resident, who assessed the data sets after the evaluation was completed and reached a consensus on the location of the IAC.

Results: The medical 3 MP monitor demonstrated the best interrater reliability with a percent agreement of 87% and a kappa value of 0.83. Accuracy was significantly greater with the medical-grade 3 MP compared with the consumer-grade display monitor, with an average increase in accuracy of 10.1%. A significantly higher accuracy (7.2%) was also obtained for the medical-grade 3 MP in comparison with the medical-grade 2 MP by 1 observer.

Discussion/Conclusions: This study found that the IAC can be well visualized and with a higher degree of accuracy on medical-grade display (3 MP) monitors compared with consumer-grade display monitors. The 2 MP prototype medical monitor showed a higher degree of accuracy compared with the consumer-grade monitor, although the differences were not statistically significant.

References

- 1 Sandstedt P, Sørensen S. Neurosensory disturbances of the trigeminal nerve: a long-term follow-up of traumatic injuries. J Oral Maxillofac Surg. 1995;53:498-505.