



Original article

Transtibial pullout repair of medial meniscus posterior root tear restores physiological rotation of the tibia in the knee-flexed position



Yuki Okazaki, Takayuki Furumatsu*, Yuya Kodama, Tomohito Hino, Yusuke Kamatsuki, Yoshiki Okazaki, Shin Masuda, Shinichi Miyazawa, Hirotsuke Endo, Tomonori Tetsunaga, Kazuki Yamada, Toshifumi Ozaki

Department of Orthopaedic Surgery, Okayama University Graduate School of Medicine, Dentistry, and Pharmaceutical Sciences, 2-5-1 Shikatacho, Kitaku, 700-8558 Okayama, Japan

ARTICLE INFO

Article history:

Received 13 July 2018

Accepted 23 October 2018

Keywords:

Medial meniscus

Posterior root tear

Magnetic resonance imaging

Tibial rotation

Transtibial pullout repair

ABSTRACT

Background: Medial meniscus posterior root tear (MMPRT) results in joint overloading and degenerative changes in the knee. Favorable clinical outcomes have been reported after transtibial pullout repair of MMPRT. To date, however, in vivo tibial rotational changes before and after root repair remain poorly understood. The purpose of this study was to investigate postoperative changes in tibial rotation following MMPRT pullout repair.

Hypothesis: Pathological external rotation of the tibia in the knee-flexed position is caused by MMPRT and is reduced after transtibial pullout repair.

Patients and methods: Fifteen patients who underwent MMPRT pullout repair and 7 healthy volunteers were included. Magnetic resonance imaging examinations were performed in the 10° and 90° knee-flexed positions. The angles between the surgical epicondylar axis and a line between the medial border of the patellar tendon and the apex of the medial tibial spine were measured. Baseline was defined as a line lying at a right angle to the other, and a value was positive and negative when the tibia rotated internally and externally, respectively.

Results: In the volunteer's normal knees, tibial internal rotation was $+1.00^\circ \pm 3.27^\circ$ at 10° flexion and $+4.14^\circ \pm 3.46^\circ$ at 90° flexion. In the MMPRT preoperative knees, tibial internal rotation was $+1.07^\circ \pm 3.01^\circ$ at 10° flexion and $+1.27^\circ \pm 2.96^\circ$ at 90° flexion. In the postoperative knees, tibial internal rotation was $+1.60^\circ \pm 2.85^\circ$ at 10° flexion and $+4.33^\circ \pm 2.89^\circ$ at 90° flexion.

Discussion: This study demonstrates discontinuity of the MM posterior root may induce a pathological external rotation of the tibia during knee flexion and that MMPRT pullout repair reduces the pathological external rotation of the tibia in the knee-flexed position.

Level of evidence: III, comparative retrospective study.

© 2018 Elsevier Masson SAS. All rights reserved.

1. Introduction

Medial meniscus posterior root tear (MMPRT) decreases the contact area and increases the contact pressure between the femoral condyle and tibial plateau [1], resulting in joint overloading and rapid knee degeneration. Although favorable clinical outcomes are reported after transtibial pullout repair of MMPRT [2,3], unfavorable outcomes are reported for non-operative treatment in patients with medial meniscus (MM) posterior horn/root tears [4] and partial meniscectomy of complete MMPRT [5,6]. MM

extrusion progresses shortly after MMPRT [7] and is correlated with injury duration in patients with knee osteoarthritis [8,9]. Patients with decreased meniscus extrusion at 1 year postoperatively had more favorable clinical scores and radiographic findings at midterm follow-up than did those with increased extrusion [10]. Therefore, prompt diagnosis of MMPRT on the basis of clinical signs [11] or a single painful popping event is necessary [12], and MMPRT repair is recommended to prevent subsequent cartilage degeneration or meniscus extrusion [13].

A robotic arm study reported that tibial external rotation is increased in association with MMPRT, and rotation is restored by posterior root repair [14]. However, to our knowledge, no previous study has reported changes in in vivo tibial rotation before and after MMPRT pullout repair. Although magnetic resonance imaging

* Corresponding author.

E-mail address: matino@md.okayama-u.ac.jp (T. Furumatsu).

(MRI) can confirm reduction of the MM posterior root to the original footprint of the tibial surface on the basis of disappearance of the giraffe neck sign [15] or cleft/truncation sign after pullout repair, in vivo tibial rotational change after root repair is poorly understood.

The purpose of this study was to investigate postoperative change in tibial rotation following MMPRT pullout repair. We hypothesized that MMPRT pullout repair could reduce pathological external rotation of the tibia in the knee-flexed position.

2. Patients and methods

2.1. Patients

The study was approved by our Institutional Review Board, and all patients and volunteers provided informed consent. This retrospective study analyzed the changes in tibial rotation preoperatively and at 3 months after MMPRT pullout repair in comparison with healthy knees. Forty-two patients who underwent MMPRT pullout repair between March 1, 2016, and October 31, 2017, and 7 healthy volunteers were included. Patients were treated with a FasT-Fix[®] (Smith & Nephew)-dependent modified Mason–Allen suture using the all-inside technique after creating the tibial bone tunnel with an MMPRT guide, as previously described [16–20]. Twenty-seven patients were excluded because their postoperative MRI (axial views) data were unavailable. The final sample size ($n=15$) demonstrated adequate power (>0.90) to detect a

significant difference between postoperative tibial rotation at 10° and 90° flexion and between pre- and postoperative tibial rotation at 90° flexion. The Kellgren–Lawrence grade was determined using plain radiographs and MMPRT type, through careful arthroscopic examinations [13]. MRI examinations were performed preoperatively and at 3 months postoperatively (Figs. 1 and 2).

2.2. MRI evaluation

Open MRI scanning was performed preoperatively and at 3 months postoperatively using an OASIS 1.2 T device (Hitachi Medical, Chiba, Japan) with a coil at 10° and 90° flexion under a non-weight-bearing condition. To standardize the position of the knee in each patient, lower leg and foot were held using sandbags and formed styrene in 0° of dorsal flexion of the ankle and keeping the second metatarsal bone horizontally. Standard OASIS sequences included an axial T2-weighted multi-echo sequence (repetition time/echo time of 4438 ms/84 ms) with a 90° flip angle. The slice thickness was 4 mm with a 0-mm gap. The field of view was 16 cm with an acquisition matrix size of 352 × 320.

We used a picture archiving and communication system (FUJIFILM Holdings Corporation, Tokyo, Japan) to measure the tibial rotation. Axial views were taken parallel to the medial tibial plateau, and after assessing all axial slices for each knee, measurements were taken from a view in which we could identify the intercondylar tibial ridge. Then, we measured tibial rotation on the basis of the angle formed between the surgical epicondylar axis




Fig. 1. Magnetic resonance images of a normal left knee. A. Surgical epicondylar axis (SEA, solid line) in 10° knee-flexed position. B. Internal rotation (+1°) in 10° knee-flexed position. An angle is formed by a solid line perpendicular to the SEA and a dotted line linking the medial border of the patellar tendon (circle) and the apex of the medial tibial spine (filled circle). C. SEA (solid line) in 90° knee-flexed position. D. Internal rotation (+5°) in 90° knee-flexed position.




Fig. 2. Magnetic resonance images of a knee with MMPRT and a repaired left knee. A. Preoperative image. External rotation (-1°) in 10° knee-flexed position. An angle is formed by the solid line perpendicular to the SEA and the dotted line linking the medial border of the patellar tendon (circle) and the apex of the medial tibial spine (filled circle). B. Preoperative image. External rotation (-1°) in 90° knee-flexed position. C. Postoperative image. External rotation (-1°) in 10° knee-flexed position. D. Postoperative image. Internal rotation ($+2^\circ$) in 90° knee-flexed position.

(SEA) and a line between the medial border of the patellar tendon and the apex of the medial tibial spine (PTMS) (Figs. 1 and 2), which is reportedly a good reference line for MM rotation [21]. Baseline was defined as a line lying at a right angle to the other, and the value was positive and negative when the tibia rotated internally and externally, respectively.

2.3. Statistical analysis

Data are reported as mean \pm standard deviation. Statistical analysis and sample size/power calculation were performed using EZR software (Saitama Medical Center Jichi Medical University, Tochigi, Japan). The *t*-test and Wilcoxon signed rank test were used to compare values. Statistical significance was set at $p < 0.05$. Two orthopedic surgeons independently measured the angle formed between SEA and PTMS. Each observer performed each measurement twice, with a 2-week interval. Inter-observer and intra-observer reliabilities were assessed using the intra-class correlation coefficient (ICC). The inter-observer and intra-observer reliabilities for the measurements were satisfactory with the mean ICC values of 0.91 and 0.95, respectively.

3. Results

3.1. Demographic data and MRI findings

Table 1 reports the patients' demographic and clinical characteristics. The mean age of the volunteers was 27.6 (range, 22–33) years.

Table 1
Demographic and clinical characteristics.

	MMPRT group	Volunteer group	<i>P</i> value
Number of patients	15	7	
Age (years)	59.8 \pm 9.7	27.6 \pm 4.65	<0.001 ^a
Men/women	2/13	3/4	
Height (m)	1.56 \pm 0.06	1.65 \pm 0.09	0.009 ^a
Weight (kg)	65.9 \pm 13.7	55.7 \pm 12.3	0.11
BMI (kg/m ²)	27.0 \pm 4.62	20.3 \pm 2.64	0.002 ^a
Root tear classification type 1/2/3/4/5	0/13/0/2/0	N/A	
Kellgren–Lawrence grade I/II	10/5	N/A	
Duration from injury to surgery (weeks)	15.5 \pm 13.9	N/A	

Data are presented as mean \pm standard deviation. MMPRT: medial meniscus posterior root tear; BMI: body mass index; NA: not available.

^a Statistically significant.

The mean age at the time of surgery was 59.8 (range, 35–72) years, and the mean duration from injury to surgery was 15.5 (range, 1–42) weeks. Giraffe neck sign was observed in 14/15 patients (93.3%), and cleft/truncation signs were observed in 13/15 patients (86.7%) before surgery. Those signs each became negative in all patients and in 12/13 patients (92.3%) after surgery. Restoration of medial extrusion over 0.50 mm was identified in only one case.




Fig. 3. Magnetic resonance imaging-based tibial internal rotation in 10° and 90° knee-flexed positions. A. Normal knee. * $p < 0.05$. B. Preoperative knee. C. Postoperative knee. ** $p < 0.001$.

3.2. Tibial rotation on serial MRI

In the volunteers' normal knees, tibial internal rotation was $+1.00^\circ \pm 3.27^\circ$ at 10° flexion and $+4.14^\circ \pm 3.46^\circ$ at 90° flexion. In the MMPRT preoperative knees, tibial internal rotation was $+1.07^\circ \pm 3.01^\circ$ at 10° flexion and $+1.27^\circ \pm 2.96^\circ$ at 90° flexion. In the postoperative knees, tibial internal rotation was $+1.60^\circ \pm 2.85^\circ$ at 10° flexion and $+4.33^\circ \pm 2.89^\circ$ at 90° flexion.

In the normal and postoperative knees, significantly greater tibial internal rotation was observed at 90° flexion than at 10° ($p < 0.05$), whereas in the preoperative knees, no significant difference in tibial rotation was observed between 90° and 10° flexion (Fig. 3).

At 90° flexion, tibial internal rotation was significantly restored in the postoperative knees compared with the preoperative knees ($p < 0.001$), whereas no difference was observed between the postoperative and normal knees (Fig. 4). At 10° flexion, no significant differences in tibial rotation were observed among the groups.

4. Discussion

The most important finding of this study was that intrinsic internal rotation of the tibia during knee flexion was not observed in patients with MMPRTs, and MMPRT pullout repair restored physiological tibial internal rotation of middle-aged patients in the 90° knee-flexed position. Our hypothesis is confirmed.

The tibia rotates internally from 0° to 120° in normal knee, the so-called medial pivot, and shows a bilateral femoral condyle roll back from 120° to full flexion [22]. A cadaveric study with a 100-N axial load using a robotic arm reported that tibial external rotation increased in association with MMPRT in 30°, 60°, and 90° knee-flexed positions; no significant difference was found in the knee-extended position, and external rotation in the knee-flexed position was restored by posterior root repair [23]. However, extrapolating these data to the clinical population is difficult because muscle contraction, proprioception, and healing were not considered. Furthermore, arthroscopic repair was not performed. Our results are consistent with the results of this cadaveric study reporting increased tibial external rotation with MMPRT, and restoration of internal rotation by posterior root repair. Intrinsic internal rotation of the tibia during knee flexion was not observed in patients with MMPRTs.

The MM extrudes posteriorly with MMPRT in the knee-flexed position [24]; thus, given the disruption of the original MM mechanism as a secondary stabilizer, posterior translation of the medial femoral condyle cannot be prevented by the MM posterior segment, and MMPRT may induce pathological tibial external rotation during knee flexion via discontinuity of the MM posterior root. The dynamic stability structures of the knee posteromedial corner, including the MM posterior root, are important [25]; we consider that MMPRT pullout repair restores MM function as a joint stabilizer against tibial external rotation, resulting in internal rotation.




Fig. 4. Magnetic resonance imaging-based tibial internal rotation in 90° knee-flexed position. A. Normal and preoperative knees. * $p < 0.05$. B. Preoperative and postoperative knees. ** $p < 0.001$.

This study has several limitations. First, given its retrospective design, relatively small number of patients, and insufficient results of second-look arthroscopy due to short-term follow-up, the duration from injury to surgery was not investigated in the study, although a longer duration from injury to surgery may lead to the progression of degenerative change and tibial external rotation. However, the sample size had acceptable statistical power for the endpoint. Second, the non-weight-bearing condition and evaluation only from the 10° and 90° knee-flexed positions might not be precise enough to detect subtle tibial rotational changes. Third, the initial tension of suture fixation was not considered. Pullout repair is essential for MMPRT, because the end-to-end suture technique is not suitable, and we performed tibial fixation using the double-spike plate and screw, with the knee flexed in neutral rotation at 45° and a 20-N initial tension. It is possible that the fixed tension or rotation would affect postoperative tibial rotation. Additional MRI investigations involving long-term follow-up and larger sample sizes are required.

5. Conclusions

MMPRT may induce pathological external rotation of the tibia during knee flexion because of the discontinuity of the MM posterior root. Pullout repair may restore physiological internal rotation of the tibia during knee flexion.

Disclosure of interest

The authors declare that they have no competing interest.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Contributions of authors

Takayuki Furumatsu designed the study. Yoshiaki Okazaki and Shin Masuda contributed to the analysis and interpretation of data. All authors have contributed to data collection and interpretation and critically reviewed the manuscript. All authors approved the final version of the manuscript, and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Acknowledgment

We thank Editage (<http://www.editage.jp>) for English language editing.

References

- [1] Marzo JM, Gurske-DePerio J. Effects of medial meniscus posterior horn avulsion and repair on tibiofemoral contact area and peak contact pressure with clinical implication. *Am J Sports Med* 2009;37:124–9.
- [2] Moatshe G, Chahla J, Slette E, Engebretsen L, LaPrade RF. Posterior meniscal root injuries: a comprehensive review from anatomy to surgical treatment. *Acta Orthop* 2016;87:452–8.
- [3] Kim SB, Ha JK, Lee SW, Kim DW, Shim JC, Kim JG, et al. Medial meniscus root tear refixation: comparison of clinical, radiologic, and arthroscopic findings with medial meniscectomy. *Arthroscopy* 2011;27:346–54.
- [4] Krych AJ, Reardon PJ, Johnson NR, Mohan R, Peter L, Levy BA, et al. Non-operative management of medial meniscus posterior horn root tears is associated with worsening arthritis and poor clinical outcome at 5-year follow-up. *Knee Surg Sports Traumatol Arthrosc* 2017;25:383–9.
- [5] Han SB, Shetty GM, Lee DH, Chae DJ, Seo SS, Wang KH, et al. Unfavorable results of partial meniscectomy for complete posterior medial meniscus root tear with early osteoarthritis: a 5- to 8-year follow-up study. *Arthroscopy* 2010;26:1326–32.
- [6] Ozkoc G, Circi E, Gonc U, Irgit K, Pourbagher A, Tandogan RN. Radial tears in the root of the posterior horn of the medial meniscus. *Knee Surg Sports Traumatol Arthrosc* 2008;16:849–54.
- [7] Furumatsu T, Kodama Y, Kamatsuki Y, Hino T, Okazaki Y, Ozaki T. Meniscal extrusion progresses shortly after the medial meniscus posterior root tear. *Knee Surg Relat Res* 2017;29:295.
- [8] Furumatsu T, Kamatsuki Y, Fujii M, Kodama Y, Okazaki Y, Masuda S, et al. Medial meniscus extrusion correlates with disease duration of the sudden symptomatic medial meniscus posterior root tear. *Orthop Traumatol Surg Res* 2017;103:1179–82.
- [9] Lee DW, Moon SG, Kim NR, Chang MS, Kim JG. Medial knee osteoarthritis precedes medial meniscal posterior root tear with an event of painful popping. *Orthop Traumatol Surg Res* 2018;104:1009–15.
- [10] Chung KS, Ha JK, Ra HJ, Nam GW, Kim JG. Pullout fixation of posterior medial meniscus root tears: correlation between meniscus extrusion and midterm clinical results. *Am J Sports Med* 2017;45:42–9.
- [11] Seil R, Duck K, Pape D. A clinical sign to detect root avulsions of the posterior horn of the medial meniscus. *Knee Surg Sports Traumatol Arthrosc* 2011;19:2072–5.
- [12] Bae JH, Paik NH, Park GW, Yoon JR, Chae DJ, Kwon JH, et al. Predictive value of painful popping for a posterior root tear of the medial meniscus in middle-aged to older Asian patients. *Arthroscopy* 2013;29:545–9.
- [13] Chung KS, Ha JK, Ra HJ, Kim JG. A meta-analysis of clinical and radiographic outcomes of posterior horn medial meniscus root repairs. *Knee Surg Sports Traumatol Arthrosc* 2016;24:1455–68.
- [14] Allaire R, Muriuki M, Gilbertson L, Harner CD. Biomechanical consequences of a tear of the posterior root of the medial meniscus similar to total meniscectomy. *JBJS* 2008;90:1922–31.
- [15] Furumatsu T, Fujii M, Kodama Y, Ozaki T. A giraffe neck sign of the medial meniscus: a characteristic finding of the medial meniscus posterior root tear on magnetic resonance imaging. *J Orthop Sci* 2017;22:731–6.
- [16] Furumatsu T, Kodama Y, Fujii M, Tanaka T, Hino T, Kamatsuki Y, et al. A new aiming guide can create the tibial tunnel at favorable position in transtibial pullout repair for the medial meniscus posterior root tear. *Orthop Traumatol Surg Res* 2017;103:367–71.
- [17] Fujii M, Furumatsu T, Kodama Y, Miyazawa S, Hino T, Kamatsuki Y, et al. A novel suture technique using the Fast-Fix combined with Ultrabraid for pullout repair of the medial meniscus posterior root tear. *Eur J Orthop Surg Traumatol* 2017;27:559–62.
- [18] Kodama Y, Furumatsu T, Fujii M, Tanaka T, Miyazawa S, Ozaki T. Pullout repair of a medial meniscus posterior root tear using a Fast-Fix all-inside suture technique. *Orthop Traumatol Surg Res* 2016;102:951–4.
- [19] Fujii M, Furumatsu T, Xue H, Miyazawa S, Kodama Y, Hino T, et al. Tensile strength of the pullout repair technique for the medial meniscus posterior root tear: a porcine study. *Int Orthop* 2017;41:2113–8.
- [20] Fujii M, Furumatsu T, Miyazawa S, Kodama Y, Hino T, Kamatsuki Y, et al. Bony landmark between the attachment of the medial meniscus posterior root and the posterior cruciate ligament: CT and MR imaging assessment. *Skeletal Radiol* 2017;46:1041–5.
- [21] Lee BK, Lee YS, Oh WS, Kim KH. Restoration of the anatomic position during a meniscal allograft transplantation using pre-existing landmarks. *Arch Orthop Trauma Surg* 2015;135:393–9.
- [22] Freeman MA, Inskero V. The movement of the normal tibio-femoral joint. *J Biomech* 2005;38:197–208.
- [23] Allaire R, Muriuki M, Gilbertson L, Harner CD. Biomechanical consequences of a tear of the posterior root of the medial meniscus. Similar to total meniscectomy. *JBJS Am* 2008;90:1922–31.
- [24] Masuda S, Furumatsu T, Okazaki Y, Kodama Y, Hino T, Kamatsuki Y, et al. Medial meniscus posterior root tear induces pathological posterior extrusion of the meniscus in the knee-flexed position: an open magnetic resonance imaging analysis. *Orthop Traumatol Surg Res* 2018;104:485–9.
- [25] Tang H, Bai L. Anatomy and biomechanical research progress of knee postero-medial corner. *Chinese J Repar Construct Surg* 2009;23:1058–61.