



Original article

Total knee replacement on more than 20° valgus: A case control study

Sophie Putman^{a,*}, Matthieu Ehlinger^{b,c}, Bruno Tillie^d, Benjamin Puliero^b,
Nassima Ramdane^e, Franck Remy^f, Gilles Pasquier^a, the French Hip,
Knee Society (Société Française de la hanche et du genou)^g

^a Département universitaire de chirurgie orthopédique et de traumatologie, hôpital Salengro, CHRU de Lille, rue Emile-Laine, 59037 Lille, France

^b Service de chirurgie orthopédique et de traumatologie, hôpital de Haute pierre, hôpitaux universitaires de Strasbourg, 1, avenue Molière, 67098 Strasbourg cedex, France

^c Laboratoire ICube, CNRS UMR 7357, 30, boulevard Sébastien-Brant, 67400 Illkirch, France

^d Hôpital privé Les Bonnettes, 2, rue du Dr Forgeois, BP 990, 62012 Arras, France

^e University Lille, CHU Lille, EA 2694 - santé publique: épidémiologie et qualité des soins, 59000 Lille, France

^f Clinique chirurgicale de Saint-Omer, 71, rue Ambroise-Paré, 62575 Blendecques, France

^g Société française de chirurgie de la hanche et du genou (SFHG), 56, rue Boissonnade, 75014 Paris cedex, France



ARTICLE INFO

Article history:

Received 27 June 2018

Accepted 10 December 2018

Keywords:

Knee prosthesis

Valgus knee

Complication

Constrained prosthesis

ABSTRACT

Introduction: Lower-limb valgus deformity exceeding 20° is a particular case, with few publications assessing the impact of the severity of the valgus. The present retrospective case control study compared a series of >20° valgus versus a series of 10–20° valgus, assessing (1) operative data [approach, type of total knee replacement (TKR)], (2) complications and implant survival, and (3) clinical and radiological results.

Hypothesis: Severe valgus deformity requires TKR with greater constraint, incurring a higher rate of complications and poorer implant survival.

Material and method: A multicenter retrospective study for the period January 2006 to December 2010 included 53 patients, with a mean age of 72 ± 10 years, presenting >20° valgus. The study series was matched for age and gender with a series of 53 cases of 10–20° valgus. Convexity laxity was greater in the >20° group ($p = 0.004$).

Results: There was no significant inter-group difference in approach ($p = 0.13$). Greater constraint was more frequent in the >20° group (7/53 versus 1/53; $p = 0.03$), independently of convexity laxity or Krackow grade ($p = 0.14$). There were 7 complications (13.2%) in the >20° group and 7 in the 10–20° group (NS). Eight-year survivorship was 95.12% in the >20° group and 94.9% in the 10–20° group ($p = 0.63$). There were no significant differences in Oxford score ($p = 0.30$) or HKA angle ($p = 0.78$) at last follow-up.

Conclusion: The study hypothesis was partially confirmed: greater constraint was more frequent in >20° valgus. The number of complications was low, and survival was identical to that of a control group with less severe deformity.

Level of evidence: III, retrospective case control study.

© 2019 Elsevier Masson SAS. All rights reserved.

1. Introduction

Total knee replacement (TKR) is a reliable and reproducible procedure, showing >90% 10-year survivorship [1–9]. Deformities are sometimes severe, and it is consensual that TKR in severe valgus is complicated, but that results are satisfactory and similar to those in minor deformity [10,11].

Few reports have focused on severe deformity, whether in varus [10–15] or valgus [10,11], and the latter combined both valgus and

varus, hindering analysis and interpretation of the results. Deformities exceeding 20° are a special case, rarely encountered, for which high-constraint implants are often recommended [15]. The surgical approach is also a matter of discussion, and can entail skin problems.

Few series concerned >20° valgus with a control group not of varus or normal alignment but of less severe valgus. The present retrospective case-control study comparing >20° valgus versus 10–20° valgus aimed to assess:

- operative data (approach, implant constraint);
- complications and implant survival;
- clinical and radiological data.

* Corresponding author.

E-mail address: sophie.putman@chru-lille.fr (S. Putman).

Table 1
Preoperative data for the two series.

Preoperative data	Valgus > 20°	Valgus 10–20°	p-value
<i>Devane activity grade, n (%) [17]</i>			0.20
Grade 1	8 (15.1%)	4 (7.6%)	
Grade 2	26 (49.1%)	23 (43.4%)	
Grade 3	13 (24.5%)	20 (37.7%)	
Grade 4	5 (9.4%)	6 (11.3%)	
Grade 5	1 (1.9%)	0 (0.0%)	
<i>Mean ± SD [range]</i>			
IKS pain [18]	10.2 ± 8.9 [0–30]	13.3 ± 11.3 [0–45]	0.10
IKS knee [18]	22.5 ± 15.9 [0–63]	33.4 ± 17.9 [0–80]	0.0012
IKS function [18]	40.4 ± 21.6 [0–90]	48.7 ± 20.1 [0–90]	0.0342
Flexion	105.9 ± 16.9 [50–30]	114.4 ± 16.3 [60–140]	0.0046
HKA angle	203.6 ± 4.0 [201–215]	191.8 ± 1.2 [190–194]	–
Femoral angle	98.0 ± 4.0 [91–111]	96.0 ± 3.3 [89–103]	0.0107
Tibial angle	96.5 ± 5.7 [90–116]	91.5 ± 3.8 [79–105]	<0.0001
Tibial slope	8.3 ± 5.7 [–10–20]	6.7 ± 3.7 [–3–18]	0.08
<i>Krackow classification n (%)</i>			0.001
Type 1	15 (28.3%)	45 (85%)	
Type 2	32 (60.4%)	4 (7.5%)	
Type 3	6 (11.4%)	4 (7.5%)	
<i>Convexity laxity n (%)</i>			0.004
zero	9 (17%)	23 (43.4%)	
<5°	16 (30.2%)	21 (39.6%)	
5–10°	25 (47.2%)	7 (13.2%)	
>10°	3 (5.6%)	2 (3.8%)	

The study hypothesis was that severe valgus deformity requires greater implant constraint, entailing a greater rate of complications and lower survivorship.

2. Materials and Methods

2.1. Study design

A multicenter retrospective study was conducted in 17 public and private sector hospitals under the aegis of the French Society of Hip and Knee Surgery (SFHG). The inclusion period was January 2006 to December 2010, allowing a minimum 5 years' and maximum 10 years' follow-up so as to limit incomplete records.

2.2. Patients

A total of 557 patients with valgus knee exceeding 10° received TKR.

Fifty-three patients (8 male, 45 female; mean age, 72 ± 10 years; mean body-mass index (BMI), 28 ± 4.5 [range, 18–37]) had > 20° valgus, and were matched for age, gender and BMI to a control group of 53 patients with 10–20° valgus.

The two groups were comparable for age, gender, BMI, Devane activity score and IKS pain score. They differed on IKS knee and function scores, Krackow classification [16], and convexity laxity (Table 1).

2.3. Assessment methods

Operative data comprised: medial or lateral approach, any ligament release, associated extensor system procedures, type of implant, and complications. TKR was classified as: PCL-retaining (CR), ultracongruent (UC), posterior stabilized (PS), constrained condylar knee (CCK) or hinged.

Data at last follow-up comprised: preoperative scores, and Oxford-12 score [19]. Radiographic assessment on full long-leg view was identical to the preoperative assessment. Any progressive radiolucency exceeding 2 mm, osteolysis or wear were noted.

Post-operative complications were noted as early (first 3 months) or late (> 3 months). Survival was defined as TKR present at follow-up without revision of any component.

2.4. Statistical analysis

Data were analyzed on SAS software, version 9.4 (SAS Institute Inc., Cary, NC, USA). Qualitative variables were reported as number and percentage and quantitative variables as mean, standard deviation and range. Normal distribution was checked graphically and on Shapiro-Wilk test.

To avoid exclusion of cases from matched analysis due to missing data, simple imputation was applied using regression switching (chained equation with $m = 1$), using all preoperative parameters. This enabled case-control matching on gender, age (to within 5 years) and BMI (to within 3 points). Quantitative variables were compared between groups on Student test for matched variables or Wilcoxon signed ranks test, and qualitative variables on McNemar test. Pre- to post-operative progression within groups was analyzed on Student test for matched variables or Wilcoxon signed ranks test. Survivorship was estimated following Kaplan-Meier. Associations between implant survival and qualitative variables were analyzed

Table 2
Approach and TKR type.

Operative results	Valgus > 20°	Valgus 10–20°	p-value
<i>Approach n (%)</i>			0.43
Lateral	32 (60.4%)	24 (45.3%)	
With ATT osteotomy	21 (39.6%)	9 (17%)	
Medial	21 (39.6%)	29 (54.7%)	
Mid-vastus	8 (15.1%)	6 (11.4%)	
Transquadriceps	13 (24.5%)	22 (41.5%)	
Sub-vastus	0 (0%)	1 (1.8%)	
<i>Type of TKR n (%)</i>			0.03
Posterior-stabilized	21 (39.6%)	30 (56.6%)	
Ultra-congruent	13 (24.5%)	16 (30.2%)	
Cruciate ligament-sparing	8 (15.1%)	6 (11.4%)	
CCK	4 (7.6%)	0 (0%)	
Hinged	7 (13.2%)	1 (1.8%)	

on log-rank test, and by Cox model for quantitative variables. The significance threshold was set at 5%.

3. Results

3.1. Operative data

3.1.1. Approach

There was no significant inter-group difference in approach ($p=0.43$). With the numbers available, no difference in complications according to associated anterior tibial tubercle (ATT) osteotomy was found (3 complications for 30 osteotomies).

3.1.2. Type of TKR

Higher constraint was more frequent in the $>20^\circ$ group ($p=0.03$). The rate of high-constraint implants did not differ according to whether convexity laxity was zero ($n=3$), $<5^\circ$ ($n=3$), $5-10^\circ$ ($n=3$) or $>10^\circ$ ($n=3$), or according to Krackow classification ($p=0.14$) (Table 2).

3.2. Complications and survival

Four patients in each group had died by last follow-up, of causes unrelated to TKR.

There were 7 complications (1 early, 6 late) in the $>20^\circ$ group and 7 (3 early, 4 late) in the $10-20^\circ$ group (Appendix A). Relative risk of complications was identical in both groups: HR, 0.460 [0.134–1.583].

Eight-year survivorship was 95.12% (95% CI, 81.4–98.8) in the $>20^\circ$ group, and 94.9% in the $10-20^\circ$ group ($p=0.63$).

3.3. Clinical and radiological results

IKS scores improved significantly after TKR (Table 3). The $>20^\circ$ group showed poorer clinical results than the $10-20^\circ$ group on IKS knee score ($p=0.024$) and function score ($p=0.012$) and flexion (0.0046), but these values had been poorer preoperatively. There was no significant difference in Oxford score at last follow-up ($p=0.30$).

HKA angle showed improvement over preoperative values in both groups: $180.8^\circ \pm 2.8^\circ$ [range, $172-189^\circ$] at last follow-up in the $>20^\circ$ group, and $180.6^\circ \pm 2.3^\circ$ [$176-186^\circ$] in the $10-20^\circ$ group, with no significant difference ($p=0.78$).

Table 3

Post-operative results in the case and control groups.

Post-operative results	Valgus $>20^\circ$	Valgus $10-20^\circ$	p-value
Devane activity grade n (%) [17]			0.77
Grade 1	7 (13.5%)	6 (12.8%)	
Grade 2	21 (40.4%)	17 (36.2%)	
Grade 3	15 (28.9%)	14 (29.8%)	
Grade 4	9 (17.3%)	10 (21.3%)	
Mean \pm SD [range]			
IKS pain [18]	43.4 \pm 10.2 [10–50]	44.3 \pm 11.3 [10–50]	0.23
IKS knee [18]	83.3 \pm 19.7 [0–100]	87.5 \pm 21.0 [10–100]	0.0244
IKS function [18]	60.1 \pm 27.5 [0–100]	73.4 \pm 24.8 [20–100]	0.0126
Flexion	112.9 \pm 16.7 [40–135]	120.6 \pm 11.4 [90–140]	0.0024
HKA angle	180.8 \pm 2.8 [172–189]	180.6 \pm 2.3 [176–186]	0.78
Femoral angle	90.6 \pm 1.4 [88–94]	90.6 \pm 1.6 [86–94]	0.80
Tibial angle	90.3 \pm 1.6 [86–94]	89.9 \pm 1.8 [85–96]	0.12
Tibial slope	2.4 \pm 4.1 [–6–12]	4.7 \pm 3.7 [–3–13]	0.0094
Oxford score [20]	33.6 \pm 10.4 [8–45]	37.9 \pm 10.5 [9–48]	0.30

Table 4

Main literature series on severe deformity.

Series	Number	Type of deformity	Comparative study?	Survival	Follow-up (years)	Results
Teeny et al. [12]	27	Varus $>20^\circ$	Yes (vs. “non-significant deformity”)	ND	4.9	Knee Evaluation Score = 89 points Significantly poorer than in controls
Karachalios et al. [10]	51	Varus and valgus $>20^\circ$	Yes (vs. $<5^\circ$ deformity)	ND	5.5	Bristol knee score = 81 points Equivalent clinical results
Ritter et al. [11]	82	Varus and valgus $>20^\circ$	Yes (vs. $<5^\circ$ varus/valgus)	98%	6.5	IKS knee = 87.8 points No significant difference between $>20^\circ$ and $<5^\circ$ groups
Mullaji et al. [13]	173	Varus $>20^\circ$	No	96%	2.6	IKS global = 163.2 points IKS knee = 91.1 points IKS function = 72.1 points
Kim et al. [14]	50	Varus $>20^\circ$	Yes (vs. $<20^\circ$ varus)	100%	1	IKS global = 174.3 points IKS knee = 93.3 points IKS function = 81 points No difference in clinical scores or alignment
Czekaj et al. [28]	170	Varus >10 and valgus $>10^\circ$	No	98%	5–12	10-year survival $>98\%$ with non-constrained implants
Present series	53	Valgus $>20^\circ$	No	95.12%	8	

ND: no data.

4. Discussion

The present study found that valgus deformity exceeding 20° required more frequent high-constraint TKR, but did not incur more complications or affect implant survival as compared to a control group with 10–20° valgus. The study hypothesis was thus partially confirmed.

Various approaches have been described for valgus deformity [20–24], and notably a lateral approach in severe valgus, often associated to tibial tubercle osteotomy. In the present series, the approach was medial or lateral, whatever the implant, and proved unrelated to the rate of complications, as already suggested by Gunst et al. [25]. Tibial tubercle osteotomy is often thought to increase the risk of complications [26], but did not alter the present results.

On the other hand, >20° valgus was independently associated with more frequent high constraint; >20° valgus was indeed also associated with greater convexity laxity, but this factor was not itself associated with choice of implant. According to Girard et al. [15], in valgus exceeding 5°, valgus laxity was the only factor associated with higher constraint on multivariate analysis. It thus seems difficult to clearly identify risk factors.

There were few complications in the present series. Patellar complications are commonly reported in valgus deformity [27]. In the present series, however, there was only 1 case of late patellar instability, in the >20° group. The literature on implant survival in severe deformity is sparse (Table 4). Ritter et al. [11] compared cases of severe varus and valgus versus a control group with milder deformity. Ten-year survivorship in series of TKR for all deformities is greater than 90%: Rand et al. [4] reported 91%, Argenson et al. [9] 92% and Sundberg et al. [7] 97% in the Swedish register. The present series of severe valgus likewise showed more than 90% 8-year survivorship, suggesting that severity does not impact implant survival.

Ritter et al. [11] reported no significant difference in clinical scores between >20° and <5° deformity, suggesting that the severity of initial deformity does not impact results, in contrast to the findings of Teeny et al. [10] for >20° varus. In the present series, certain scores were poorer in the >20° group, but these patients had lower baseline scores and actually showed significant improvement.

The present study had certain limitations:

- the design was retrospective; however, such severe deformity is too rare to allow prospective assessment;
- the design was also multicenter, with a variety of TKR techniques; this, however, corresponds to average clinical activity;
- the participating surgeons varied in their experience; this, however, only strengthens the external validity of the conclusions;
- the series was not continuous and, despite all effort, some cases were doubtless not included, as it is difficult to identify these severe deformities without a dedicated data-base; however, the inclusion period was limited and follow-up was not too long, so that implant survival rates were not biased by patient mortality.

A minimum 5 years' follow-up is, moreover, considered sufficient to screen for risk factors for failure.

5. Conclusion

High-constraint implants were used in 20% of cases, although no associated factors could be identified. The choice of surgical approach was at the surgeon's discretion, and did not affect the rate of complications. The complications rate in >20° valgus was

low and identical to that in less severe deformity, with implant survival exceeding 95%.

Disclosure of interest

Outside the study, SP is a consultant for Corin; ME is a consultant for Depuy-Synthes, Newclip and Lepine and associate editor for the SoFCOT Instructional Courses; FR is a consultant for Zimmer/Biomet; GP is a consultant for Zimmer/Biomet.

The other authors declare that they have no competing interest.

Authors' contributions

SP: article writing and data collection; ME: article writing and data collection; BP: data collection; BT: data collection; NR: statistical analysis; FR: data collection; GP: article writing and data collection.

Acknowledgments

We thank all the centers that contributed data: Amiens, Arras, Berck-Calot, Clermont-Ferrand, Grenoble, Lille, Marseille, Montélimar, Rennes, Rouen, Strasbourg, St Étienne, Arras, St Omer, Toulouse, Antilles and Limoges, and the SFHG.

We thank personally the symposium members and their teams for involvement in the study: J.L. Rouvillain, C. Mabit, B. Boyer, M. Ehlinger, D. Hutten, B. Rubens-Duval, S. Descamps, A. Cazenave, A. Gabrion, T. Gaillard, T. Jouanin, R. Bouchet, P. Bizzozéro, B. Boyer, H. Common, J. Zalahoui, J. Taviaux, J. Murgier, R. Pailhé, and J.L. Briard.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.otsr.2018.12.014>.

References

- [1] Ritter MA. The Anatomical Graduated Component total knee replacement: a long-term evaluation with 20-year survival analysis. *J Bone Joint Surg Br* 2009;91:745–9.
- [2] Fernandez-Fernandez R, Rodriguez-Merchan EC. Better survival of total knee replacement in patients older than 70 years: a prospective study with 8 to 12 years follow-up. *Arch Bone Jt Surg* 2015;3:22–8.
- [3] Rand JA, Ilstrup DM. Survivorship analysis of total knee arthroplasty: Cumulative rates of survival of 9200 total knee arthroplasties. *J Bone Joint Surg Am* 1991;73:397–409.
- [4] Rand JA, Trousdale RT, Ilstrup DM, Harmsen WS. Factors affecting the durability of primary total knee prostheses. *J Bone Joint Surg Am* 2003;85:259–65.
- [5] Furnes O, Espehaug B, Lie SA, Vollset SE, Engesaeter LB, Havelin LI. Early failures among 7,174 primary total knee replacements: a follow-up study from the Norwegian Arthroplasty Register 1994–2000. *Acta Orthop Scand* 2002;73:117–29.
- [6] Vollset SE, Furnes O. Failure of total knee arthroplasty with or without patella resurfacing. *Acta Orthop* 2011;82:282–92.
- [7] Sundberg M, Lidgren L, W-Dahl A, Robertsson O. Swedish Knee Arthroplasty Register; Lund University. <http://www.myknee.se/pdf/SVK>.
- [8] Gøthesen O, Espehaug B, Havelin L, Petursson G, Lygre S, Ellison P, et al. Survival rates and causes of revision in cemented primary total knee replacement: a report from the Norwegian Arthroplasty Register 1994–2009. *Bone Joint J* 2013;95:636–42.
- [9] Argenson JN, Boisgard S, Parratte S, Descamps S, Bercovy M, et al. Survival analysis of total knee arthroplasty at a minimum 10 years' follow-up: a multicenter French nationwide study including 846 cases. *Orthop Traumatol Surg Res* 2013;99:385–90.
- [10] Karachalios T, Sarangi PP, Newman JH. Severe varus and valgus deformities treated by total knee arthroplasty. *J Bone Joint Surg Br* 1994;76:938–42.
- [11] Ritter MA, Faris GW, Faris PM, Davis KE. Total knee arthroplasty in patients with angular varus or valgus deformities of > or = 20 degrees. *J Arthroplasty* 2004;19:862–6.
- [12] Teeny SM, Krackow KA, Hungerford DS, Jones M. Primary total knee arthroplasty in patients with severe varus deformity. A comparative study. *Clin Orthop Relat Res* 1991;273:19–31.
- [13] Mullaji AB, Padmanabhan V, Jindal G. Total knee arthroplasty for profound varus deformity: technique and radiological results in 173 knees with varus of more than 20 degrees. *J Arthroplasty* 2005;20:550–61.

- [14] Kim MS, Koh IJ, Choi YJ, Kim YD. In Y. Correcting severe varus deformity using trial components during total knee arthroplasty. *J Arthroplasty* 2017;32:1488–95.
- [15] Girard J, Amzallag M, Pasquier G, Mulliez A, Brosset T, et al. Total knee arthroplasty in valgus knees: predictive preoperative parameters influencing a constrained design selection. *Orthop Traumatol Surg Res* 2009;95:260–6.
- [16] Krackow KA, Jones MM, Teeny SM, Hungerford DS. Primary total knee arthroplasty in patients with fixed valgus deformity. *Clin Orthop Relat Res* 1991;273:9–18.
- [17] Devane PA, Robinson EJ, Bourne RB, Rorabeck CH, Nayak NN, Horne JG. Measurement of polyethylene wear in acetabular components inserted with and without cement. A randomized trial. *J Bone Joint Surg Am* 1997;79:682–9.
- [18] Insall JN, Dorr LD, Scott RD, Scott WN. Rationale of the Knee Society clinical rating system. *Clin Orthop Relat Res* 1989;248:13–4.
- [19] Jenny JY, Diesinger Y. Validation of a French version of the Oxford knee questionnaire. *Orthop Traumatol Surg Res* 2011;97:267–71.
- [20] Mertl P, Jarde O, Blejwas D, Vives P. Lateral approach of the knee with tibial tubercle osteotomy for prosthetic surgery. *Rev Chir Orthop* 1992;78:264–7.
- [21] Lange J, Haas SB. Correcting severe valgus deformity: taking out the knock. *Bone Joint J* 2017;99:60–6.
- [22] Apostolopoulos AP, Nikolopoulos DD, Polyzois I, Nakos A, Liarokapis S, et al. Total knee arthroplasty in severe valgus deformity: interest of combining a lateral approach with a tibial tubercle osteotomy. *Orthop Traumatol Surg Res* 2010;96:777–84.
- [23] Chalidis BE, Ye K, Sachinis NP, Hawdon G, McMahon S. Lateral parapatellar approach with tibial tubercle osteotomy for the treatment of non-correctable valgus knee osteoarthritis: a retrospective clinical study. *Knee* 2014;21:204–8.
- [24] Ranawat AS, Ranawat CS, Elkus M, Rasquinha VJ, Rossi R, Babhulkar S. Total knee arthroplasty for severe valgus deformity. *J Bone Joint Surg Am* 2005;87:271–84.
- [25] Gunst S, Villa V, Magnussen R, Servien E, Lustig S, Neyret P. Equivalent results of medial and lateral parapatellar approach for total knee arthroplasty in mild valgus deformities. *Int Orthop* 2016;40:945–51.
- [26] Young CF, Bourne RB, Rorabeck CH. Tibial tubercle osteotomy in total knee arthroplasty surgery. *J Arthroplasty* 2008;23:371–5.
- [27] Meding JB, Fish MD, Berend ME, Ritter MA, Keating EM. Predicting patellar failure after total knee arthroplasty. *Clin Orthop Relat Res* 2008;466:2769–74.
- [28] Czekaj J, Fary C, Gaillard T, Lustig S. Does low-constraint mobile bearing knee prosthesis give satisfactory results for severe coronal deformities? A five to twelve year follow up study. *Int Orthop* 2017;41:1369–77.