



Review article

Incidence and risk factors for cyclops syndrome after anterior cruciate ligament reconstruction: A systematic literature review



Thibaut Noailles^a, Antoine Chalopin^b, Mathieu Boissard^c, Ronny Lopes^d,
Nicolas Bouguennec^e, Alexandre Hardy^{f,*}

^a Département de Chirurgie Orthopédique, Polyclinique de Bordeaux Nord, 15, rue Claude-Boucher, 33000 Bordeaux, France

^b Département de Chirurgie Orthopédique Infantile, Hôpital Universitaire Hôtel-Dieu, 1, place Alexis-Ricordeau, 44000 Nantes, France

^c Département de Chirurgie Orthopédique, Polyclinique de l'Atlantique, avenue Claude-Bernard - BP 40419, 44819 Saint Herblain Cedex, France

^d Département de Chirurgie Orthopédique, Clinique Bretéché, 3, rue de la Béraudière, 44000 Nantes, France

^e Département de Chirurgie Orthopédique, Clinique du Sport, 4, rue Georges-Negrevergne, 33700 Mérignac, France

^f Département de Chirurgie Orthopédique, Hôpital Ambroise-Paré, 9, avenue Charles de Gaulle, 92100 Boulogne-Billancourt, France

ARTICLE INFO

Article history:

Received 17 November 2018

Accepted 18 July 2019

Keywords:

Anterior cruciate ligament

Arthrofibrosis

Cyclops syndrome

Extension lag

Stiffness

Systematic literature review

ABSTRACT

Background: Cyclops syndrome after anterior cruciate ligament (ACL) reconstruction is due to a fibrous nodule that develops in the anterior part of the intercondylar notch and prevents full knee extension. The primary objective of this systematic literature review was to evaluate the incidence of symptomatic cyclops lesion after ACL reconstruction. The secondary objective was to identify risk factors for cyclops syndrome.

Hypothesis: Cyclops syndrome is common after ACL reconstruction and has several risk factors reported in the literature.

Methods: A systematic literature review was performed by searching the PubMed, Medline, CINAHL, Cochrane, and Embase databases with the key terms 'cyclops' and 'ACL reconstruction'. The data thus retrieved were evaluated independently by two investigators. All articles in English or French that reported the incidence and risk factors of cyclops syndrome after ACL reconstruction were included.

Results: The search retrieved the titles and abstracts of 79 articles, of which 20 were selected to be read in full; among these, 10 were included in the study. The incidence of symptomatic cyclops lesion ranged from 1.9% to 10.9%. Identified risk factors were as follows: pre-operatively, knee inflammation and/or motion restriction at the time of ACL reconstruction; intra-operatively, narrow intercondylar notch and excessively anterior position of the tibial tunnel; and post-operatively, persistent hamstring muscle spasm.

Discussion: Development of a cyclops lesion is common after ACL reconstruction, occurs early, and may require further surgery. The knowledge of the risk factors provided by this study may improve the ability to devise effective preventive measures.

Level of evidence: II, systematic literature review.

© 2019 Elsevier Masson SAS. All rights reserved.

1. Introduction

Reconstruction of the anterior cruciate ligament (ACL) is a commonly performed procedure that produces reliable and reproducible outcomes [1–3]. Although the post-operative complication rate is low, loss of knee extension may require revision surgery [4–7]. Cyclops syndrome was first described in 1990 by Jackson and

Schaefer as loss of full knee extension due to the development of a fibrous nodule at the base of the ACL [8–10]. Early impingement of the nodule on the roof of the intercondylar notch limits knee extension [9]. Repeated microtrauma perpetuates the fibrotic process responsible for nodule formation, resulting in painful extension loss [9–11]. Early revision surgery is required to prevent the development of degenerative lesions in the short term [12,13]. Cyclops syndrome has been reported in patients who did not undergo ACL reconstruction [10,14,15]. Fibrous tissue at the tibial attachment of the reconstructed ACL is often visible by magnetic resonance imaging (MRI) but is not associated with clinical symptoms [16,17].

* Corresponding author.

E-mail address: alexandre.hardy@me.com (A. Hardy).




Fig. 1. PRISMA flow diagram.

The diagnosis of cyclops syndrome rests on the post-operative clinical and MRI findings [18,19]. Early revision surgery is effective in restoring motion range in the absence of osteoarthritis [12].

The primary objective of this systematic literature review was to evaluate the incidence of symptomatic cyclops lesion after ACL reconstruction. The secondary objective was to identify risk factors for cyclops syndrome. The working hypothesis was that cyclops syndrome is common after ACL reconstruction and has several risk factors reported in the literature.

2. Material and Methods

The study followed recommendations for systematic literature reviews and meta-analyses [20,21]. The study objectives, methodology, and criteria for including and excluding articles were determined before data collection. The PubMed, Medline, CINAHL, Cochrane, and Embase databases were searched in July and August 2018 using two MeSH indexing terms, 'cyclops' and 'ACL reconstruction'. As cyclops syndrome was first described in 1990 [7], no major studies on the topic were published before this date. Consequently, indexing terms such as 'arthrofibrosis' and 'stiffness' did not retrieve additional articles.

Articles were included if they supplied information on the incidence and risk factors of cyclops syndrome after ACL reconstruction. Information was recorded on the diagnostic tools used to determine the incidence of cyclops syndrome (e.g., physical examination, imaging studies, arthroscopy, or several tools combined). The other inclusion criteria were as follows: article in English or French, abstract available online, patients aged 18 years or older, autograft or allograft ACL reconstruction, and clinical trial or cohort study. No time limits were applied. Studies were excluded if they involved patients with injuries to multiple ligaments or concomitant bone lesions, more than one ACL reconstruction procedure, or procedures on the bone in addition to the ACL reconstruction. In addition, cadaver studies and case reports were excluded.

Two of us (TN and AH), working independently of each other, selected articles based on the titles and abstracts. Disagreements

were resolved by consensus. After an extensive review of the articles, a second selection procedure was applied to ensure that no major articles had been missed. The following data were extracted from each selected article: demographics, incidence of cyclops syndrome, risk factors for cyclops syndrome, time to symptom onset, and level of evidence.

The methodological quality of the articles was assessed by the same two investigators (TN, AH), who independently applied the Cochrane risk-of-bias tool based on the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA [20]) statement. Heterogeneity was evaluated qualitatively by comparing study designs, study populations, types of ACL reconstruction, outcomes, and blinding. Homogeneity testing was not performed.

3. Results

3.1. Article selection

Of the 79 initially retrieved articles (Fig. 1), 20 were selected based on the title and abstract. Of these, 9 were included in the study, together with an additional study identified by manually checking the reference lists of selected articles. Table 1 lists the 10 included articles. Publication dates ranged from 1995 to 2017. There was no overlap in study populations. The level of evidence was II for two articles, III for two articles, and IV for six articles. The diagnosis of cyclops syndrome was established by arthroscopy only in 3 studies, physical examination only in 3 studies, imaging studies only in 1 study, physical examination and arthroscopy combined in 2 studies, and physical examination and imaging studies combined in 1 study.

The results of the study quality assessment according to PRISMA are reported in the [electronic appendix](#).

3.2. Incidence

The incidence of symptomatic cyclops syndrome ranged from 1.9% to 10.2% after single-bundle reconstruction [18,21–23] and

Table 1
Selected studies and incidence of cyclops syndrome diagnosed by physical examination, arthroscopy, and/or magnetic resonance imaging.

Authors	Year of publication	Study design	Primary objective	Patients, <i>n</i>	Follow-up (months)	Level of evidence	Incidence of cyclops syndrome		
							Clinical	Arthroscopic	MRI
Delincé [26]	1998	Descriptive	Characteristics of the lesions	65	14-35	IV	-	35.4%	-
Muellner [30]	1999	Prospective	Outcomes of ACL reconstruction	119	5.9-9.5	II	-	17.6%	-
Ahn [5]	2007	Retrospective	Identification of good prognostic factors	208	21.2	IV	-	21.5%	-
Wang [17]	2009	Therapeutic case series	Histological study	311	13.3	IV	1.9%	14.5%	-
Sonnery-Cottet [24]	2010	Therapeutic case series	Characteristics of double-bundle ACL reconstruction	387	12	IV	3.6%	-	-
Gohil [18]	2014	RCT	Clinical and MRI evaluation of the lesions	48	12	II	10.2%	-	46.8%
Fujii [25]	2015	Case-control	Lesions and intercondylar notch size	55	45.6	IV	10.9%	27.3%	-
Pinto [28]	2017	Case-control	Cyclops syndrome and hamstring spasm	45	6	III	2.1%	-	-
Sanders [22]	2017	Historical cohort	Incidence, risk factors, and outcomes	1841	123.6	III	2%	-	-
Facchetti [8]	2017	Descriptive	MRI findings	113	24	IV	-	-	33% (after 2 years)

ACL: anterior cruciate ligament; MRI: magnetic resonance imaging; RCT: randomised controlled trial.

from 3.6% to 10.9% after double-bundle reconstruction [24,25]. The incidence of cyclops lesions visible arthroscopically as a fibrous nodule in the intercondylar notch ranged from 15.0% to 35.0% [5,24,26] and the incidence of cyclops lesions seen by MRI from 33.0% to 46.8% [8,18].

3.3. Risk factors

3.3.1. Intrinsic risk factors

A 2.6-fold increase in the risk of cyclops syndrome in young females was found in two studies [22,25]. In contrast, another study found no risk increase in young females but noted a higher risk of radiological cyclops lesions in patients with low body mass index values ($P=0.002$) [8]. The rate of surgical revision to treat cyclops syndrome was 6.3% in patients whose ACL reconstruction was performed within 48 h after the injury compared to 2.5% when the procedure was done after resolution of the inflammatory response; however, this difference was not statistically significant [27].

No associations with the development of cyclops syndrome were found for age [7,22,25,28], level of sports activities [28], or presence of bone bruising [28].

3.3.2. Surgical factors

The risk of cyclops syndrome was 5.3 times higher after surgical ACL reconstruction than after non-operative treatment [22]. Among intra-operative factors, a narrow intercondylar notch was associated with a higher risk of cyclops syndrome ($P=0.01$) [25]. Watanabe et al. [29] suggested that anterior positioning of the tibial tunnel might increase the risk, but this possibility was not supported by studies by Fujii et al. [25] and Muellner et al. [30]. When double-bundle reconstruction was performed, the use of a quadriceps tendon autograft was associated with a higher risk compared to the risk of a hamstring tendon autograft [24]. None of the studies compared the incidence of cyclops syndrome between patients managed by single-bundle versus double-bundle reconstruction.

Sanders et al. reported a 6.7-fold increase in the risk of cyclops syndrome when the ACL reconstruction was performed more than 4 weeks after the injury [22]. Others, however, did not replicate this finding [17,28].

The following factors were not associated with the occurrence of cyclops syndrome: arthroscopy or open surgery for the ACL reconstruction [26], use of patellar tendon or hamstring tendon grafts [5,17,22,28], preservation of the residual ACL fibres [18], concomitant meniscal repair [22,25], or concomitant antero-lateral reconstruction [28].

3.3.3. Post-operative factors

During the immediate post-operative period, loss of active knee extension and stunning of the quadriceps were associated with a higher risk of cyclops syndrome [24]. Pinto et al. reported that hamstring spasm 3 to 6 weeks after surgery was a risk factor for cyclops syndrome: hamstring spasm was present in 58% of the 45 patients who required revision surgery for cyclops syndrome compared to only 24% of the controls ($P<0.001$) [28]. The time from surgery to the initiation of post-operative rehabilitation was not associated with the risk of cyclops syndrome [17,25].

4. Discussion

The incidence of symptomatic cyclops lesion after ACL reconstruction ranged from 1.9% to 10.9% and remained stable over time (Table 1). This systematic literature review helped to identify several specific risk factors including knee inflammation and knee stiffness before surgery, a narrow intercondylar notch and excessively anterior tibial tunnel position, and hamstring spasm after

surgery. Knowledge of these risk factors may help to decrease the risk of cyclops syndrome [31].

Factors that were not associated with the occurrence of cyclops syndrome included age [22,28], level of sports activities [28], bone bruising [28], use of arthroscopy or open surgery [26], use of patellar or hamstring graft material [5,17,22,28], preservation of residual ACL fibres [18], concomitant meniscal repair [22,25], concomitant antero-lateral ligament reconstruction [28], and time from surgery to the initiation of rehabilitation therapy [17,25].

The best treatment of cyclops syndrome is the institution of preventive measures that decrease the incidence and impact of the condition. When revision surgery to perform anterior release is required, the range-of-motion outcomes are disappointing [32]. Revision surgery must be performed within 1 year after ACL reconstruction and must be followed by an appropriate rehabilitation programme [11,33].

Among pre-operative factors, the time from injury to surgery does not seem associated with the risk of cyclops syndrome [8]. In contrast, knee inflammation (oedema, effusion) and knee stiffness are risk factors [12,34].

During surgery, preventive measures include anatomical graft positioning, evaluation for anterior dislocation of the ACL stump that might be overlooked, notch reconstruction if needed, the use of gradually increasing drill bit diameters to create the tunnels, and routine removal of bone and cartilage fragments, notably at the tibial ACL insertion [26].

Proximally, hypertrophic scarring of the synovial membrane or ligament suspending the Hoffa fat pad to the roof of the notch should be sought by arthroscopic inspection at the end of the procedure, with the knee extended. Inverted cyclops lesions have been described and require careful attention to the roof of the notch during surgery [35,36]. A narrow notch and excessively anterior position of the tibial tunnel were also identified as risk factors [25,37]. The risk was not associated with the type of graft material (patellar vs. hamstring tendons) or preservation of the residual ACL fibres [17,38–41].

After surgery, hamstring spasm and stunning of the quadriceps were associated with a higher risk of cyclops syndrome [24,28].

One of the limitations of this study is the low level of evidence provided by the selected articles. Furthermore, cyclops syndrome is defined as loss of knee extension combined with imaging study evidence of a fibrous nodule in the intercondylar notch [8,18]. Sole reliance on arthroscopy tends to overestimate the incidence of cyclops syndrome, and MRI has a high false-positive rate [16,17].

5. Conclusion

The incidence of symptomatic cyclops syndrome ranged from 1.9% to 10.9%. The following risk factors were identified: before surgery, knee inflammation and knee motion range restriction at the time of surgery; during surgery, a narrow intercondylar notch and excessively anterior tibial tunnel; and after surgery, persistent hamstring spasm. Knowledge of these risk factors allows the development of preventive measures.

Disclosure of interest

The authors declare that they have no competing interest.

Funding

None.

Authors' contributions

TN and AH selected the articles.
All authors contributed to create and validate the manuscript.

Acknowledgements

We thank Yannick Minvielle and Isabelle Chapelle-Marcillac for editorial assistance.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.otsr.2019.07.007>.

References

- [1] Mundi R, Bhandari M. Cochrane in CORR (®): double-bundle versus single-bundle reconstruction for anterior cruciate ligament rupture in adults (Review). *Clin Orthop Relat Res* 2016;474:1099–101.
- [2] Colombet P, Dejour D, Panisset JC, Siebold R. The French Arthroscopy Society Current concept of partial anterior cruciate ligament ruptures. *Orthop Traumatol Surg Res* 2010;96(Suppl):S109–18.
- [3] Löcherbach C, Zayni R, Chambat P, Sonnery-Cottet B. Biologically enhanced ACL reconstruction. *Orthop Traumatol Surg Res* 2010;96:810–5.
- [4] Abat F, Gelber PE, Erquicia JJ, Pelfort X, Tey M, Monllau JC. Promising short-term results following selective bundle reconstruction in partial anterior cruciate ligament tears. *Knee* 2013;20:332–8.
- [5] Ahn JH, Yoo JC, Yang HS, Kim JH, Wang JH. Second-look arthroscopic findings of 208 patients after ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2007;15:242–8.
- [6] Salzler MJ, Lin A, Miller CD, Herold S, Irrgang JJ, Harner CD. Complications after arthroscopic knee surgery. *Am J Sports Med* 2014;42:292–6.
- [7] Tan SH, Lau BP, Khin LW, Lingaraj K. The importance of patient sex in the outcomes of anterior cruciate ligament reconstructions: a systematic review and meta-analysis. *Am J Sports Med* 2016;44:242–54.
- [8] Facchetti L, Schwaiger BJ, Gersing AS, Guimaraes JB, Nardo L, Majumdar S, et al. Cyclops lesions detected by MRI are frequent findings after ACL surgical reconstruction but do not impact clinical outcome over 2 years. *Eur Radiol* 2017;27:3499–508.
- [9] Jackson DW, Schaefer RK. Cyclops syndrome: loss of extension following intra-articular anterior cruciate ligament reconstruction. *Arthroscopy* 1990;6:171–8.
- [10] Runyan BR, Bancroft LW, Peterson JJ, Kransdorf MJ, Berquist TH, Ortiguera CJ. Cyclops lesions that occur in the absence of prior anterior ligament reconstruction. *Radiographics* 2007;27:e26.
- [11] Delcogliano A, Franzese S, Branca A, Magi M, Fabbriani C. Light and scan electron microscopic analysis of cyclops syndrome: etiopathogenic hypothesis and technical solutions. *Knee Surg Sports Traumatol Arthrosc* 1996;4:194–9.
- [12] Mayr HO, Brandt CM, Weig T, Koehne M, Bernstein A, Suedkamp NP, et al. Long-term results of arthroscopic arthrolysis for arthrofibrosis after anterior cruciate ligament reconstruction. *Arthroscopy* 2017;33:408–14.
- [13] Samaan MA, Facchetti L, Podoia V, Tanaka MS, Link TM, Souza RB, et al. Cyclops lesions are associated with altered gait patterns and medial knee joint cartilage degeneration at 1 year after ACL-reconstruction. *J Orthop Res* 2017;35:2275–81.
- [14] Irisawa H, Takahashi M, Hosokawa T, Nagano A. Cyclops syndrome occurring after chronic partial rupture of the anterior cruciate ligament without surgical reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2007;15:144–6.
- [15] Tonin M, Saciri V, Veselko M, Rotter A. Progressive loss of knee extension after injury. Cyclops syndrome due to a lesion of the anterior cruciate ligament. *Am J Sports Med* 2001;29:545–9.
- [16] Ahn JH, Wang JH, Lee YS, Kim JG, Kang JH, Koh KH. Anterior cruciate ligament reconstruction using remnant preservation and a femoral tensioning technique: clinical and magnetic resonance imaging results. *Arthroscopy* 2011;27:1079–89.
- [17] Wang J, Ao Y. Analysis of different kinds of cyclops lesions with or without extension loss. *Arthroscopy* 2009;25:626–31.
- [18] Gohil S, Falconer TM, Bredahl W, Annear PO. Serial MRI and clinical assessment of cyclops lesions. *Knee Surg Sports Traumatol Arthrosc* 2014;22:1090–6.
- [19] McMahon PJ, Dettling JR, Yocum LA, Glousman RE. The cyclops lesion: a cause of diminished knee extension after rupture of the anterior cruciate ligament. *Arthroscopy* 1999;15:757–61.
- [20] Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6 [e1000097].
- [21] Wright RW, Brand RA, Dunn W, Spindler KP. How to write a systematic review. *Clin Orthop Relat Res* 2007;455:23–9.
- [22] Sanders TL, Kremers HM, Bryan AJ, Kremers WK, Stuart MJ, Krych AJ. Procedural intervention for arthrofibrosis after ACL reconstruction: trends over two decades. *Knee Surg Sports Traumatol Arthrosc* 2017;25:532–7.
- [23] Sonnery-Cottet B, Panisset JC, Colombet P, Cucurulo T, Graveleau N, Hulet C, et al. Partial ACL reconstruction with preservation of the posterolateral bundle. *Orthop Traumatol Surg Res* 2012;98(Suppl):S165–70.
- [24] Sonnery-Cottet B, Lavoie F, Ogassawara R, Kasmaoui H, Scussiato RG, Kidder JF, et al. Clinical and operative characteristics of cyclops syndrome after double-bundle anterior cruciate ligament reconstruction. *Arthroscopy* 2010;26:1483–8.
- [25] Fujii M, Furumatsu T, Miyazawa S, Okada Y, Tanaka T, Ozaki T, et al. Intercondylar notch size influences cyclops formation after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2015;23:1092–9.
- [26] Delincé P, Kralliss P, Descamps PY, Fabeck L, Hardy D. Different aspects of the cyclops lesion following anterior cruciate ligament reconstruction: a multifactorial etiopathogenesis. *Arthroscopy* 1998;14:869–76.
- [27] Herbst E, Hoser C, Gföller P, Hepperger C, Abermann E, Neumayer K, et al. Impact of surgical timing on the outcome of anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2017;25:569–77.
- [28] Pinto FG, Thauant M, Daggett M, Kajetanek C, Marques T, Guimaraes T, et al. Hamstring contracture after ACL reconstruction is associated with an increased risk of cyclops syndrome. *Orthop J Sports Med* 2017;5, 2325967116684121.
- [29] Watanabe BM, Howell SM. Arthroscopic findings associated with roof impingement of an anterior cruciate ligament graft. *Am J Sports Med* 1995;23:616–25.
- [30] Muellner T, Kdolsky R, Grossschmidt K, Schabus R, Kwasny O, Plenck Jr H. Cyclops and cyclopop formation after anterior cruciate ligament reconstruction: clinical and histomorphological differences. *Knee Surg Sports Traumatol Arthrosc* 1999;7:284–9.
- [31] Noyes FR, Berrios-Torres S, Barber-Westin SD, Heckmann TP. Prevention of permanent arthrofibrosis after anterior cruciate ligament reconstruction alone or combined with associated procedures: a prospective study in 443 knees. *Knee Surg Sports Traumatol Arthrosc* 2000;8:196–206.
- [32] Said S, Christensen SE, Faunoe P, Lund B, Lind M. Outcome of surgical treatment of arthrofibrosis following ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2011;19:1704–8.
- [33] Eckenrode BJ. An algorithmic approach to rehabilitation following arthroscopic surgery for arthrofibrosis of the knee. *Physiother Theory Pract* 2018;34:66–74.
- [34] Mayr HO, Weig TG, Plitz W. Arthrofibrosis following ACL reconstruction—reasons and outcome. *Arch Orthop Trauma Surg* 2004;124:518–22.
- [35] Kambhampati S, Ware H. Inverted cyclops lesion after anterior cruciate ligament reconstruction. *Ann R. Coll Surg Engl* 2012;94:e99–100.
- [36] Rubin LE, Yeh PC, Medvecky MJ. Extension loss secondary to femoral-sided inverted cyclops lesion after anterior cruciate ligament reconstruction. *J Knee Surg* 2009;22:360–3.
- [37] Ahn JH, Lee SH, Choi SH, Lim TK. Magnetic resonance imaging evaluation of anterior cruciate ligament reconstruction using quadrupled hamstring tendon autografts: comparison of remnant bundle preservation and standard technique. *Am J Sports Med* 2010;38(9):1768–77.
- [38] Cha J, Choi SH, Kwon JW, Lee SH, Ahn JH. Analysis of cyclops lesions after different anterior cruciate ligament reconstructions: a comparison of the single-bundle and remnant bundle preservation techniques. *Skeletal Radiol* 2012;41(8):997–1002.
- [39] Hu J, Qu J, Xu D, Zhang T, Zhou J, Lu H. Clinical outcomes of remnant preserving augmentation in anterior cruciate ligament reconstruction: a systematic review. *Knee Surg Sports Traumatol Arthrosc* 2014;22(9):1976–85.
- [40] Ryu K, Iriuchishima T, Oshida M, Saito A, Kato Y, Tokuhashi Y, et al. Evaluation of the morphological variations of the meniscus: a cadaver study. *Knee Surg Sports Traumatol Arthrosc* 2015;23:15–9.
- [41] Tie K, Chen L, Hu D, Wang H. The difference in clinical outcome of single-bundle anterior cruciate ligament reconstructions with and without remnant preservation: a meta-analysis. *Knee* 2016;23:566–74.