



Discharge destination following hip fracture in Canada among previously community-dwelling older adults, 2004–2012: database study

L. Beaupre¹ · B. Sobolev² · P. Guy³ · J. D. Kim² · L. Kuramoto⁴ · K. J. Sheehan⁵ · J. M. Sutherland³ · E. Harvey⁶ · S. N. Morin⁶ · for The Canadian Collaborative Study of Hip Fractures

Received: 1 August 2018 / Accepted: 15 March 2019 / Published online: 1 April 2019
© International Osteoporosis Foundation and National Osteoporosis Foundation 2019

Abstract

Summary Little is known about post-acute care following hip fracture surgery. We investigated discharge destinations from surgical hospitals for nine Canadian provinces. We identified significant heterogeneity in discharge patterns across provinces suggesting different post-acute recovery pathways. Further work is required to determine the impact on patient outcomes and health system costs.

Introduction To examine discharge destinations by provinces in Canada, adjusting for patient, injury, and care characteristics.

Methods We analyzed population-based hospital discharge abstracts from a national administrative database for community-dwelling patients who underwent hip fracture surgery between 2004 and 2012 in Canada. Discharge destination was categorized as rehabilitation, home, acute care, and continuing care. Multinomial logistic regression modeling compared proportions of discharge to rehabilitation, acute care, and continuing care versus home between each province and Ontario. Adjusted risk differences and risk ratios were estimated.

Results Of 111,952 previously community-dwelling patients aged 65 years or older, 22.5% were discharged to rehabilitation, 31.6% to home, 27.0% to acute care, and 18.2% to continuing care, with significant variation across provinces ($p < 0.001$). The proportion of discharge to rehabilitation ranged from 2.4% in British Columbia to 41.0% in Ontario while the proportion discharged home ranged from 20.3% in Prince Edward Island to 52.2% in British Columbia. The proportion of discharge to acute care ranged from 15.2% in Ontario to 58.8% in Saskatchewan while the proportion discharged to continuing care ranged from 9.3% in Manitoba and Prince Edward Island to 22.9% in New Brunswick. Adjusting for hospital type changed the direction of the provincial effect on discharge to continuing care in two provinces, but statistical significance remained consistent with the primary analysis.

Conclusions Discharge destination from the surgical hospital after hip fracture is highly variable across nine Canadian provinces. Further work is required to determine the impact of this heterogeneity on patient outcomes and health system costs.

Keywords Discharge destination · Health services research · Hip fracture · Variation

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00198-019-04943-6>) contains supplementary material, which is available to authorized users.

✉ L. Beaupre
lauren.beaupre@ualberta.ca

¹ University of Alberta, 2-50 Corbett Hall, Edmonton, AB T6G 2G4, Canada

² School of Population and Public Health, University of British Columbia, Vancouver, Canada

³ Centre for Hip Health and Mobility, University of British Columbia, Vancouver, Canada

⁴ Centre for Clinical Epidemiology and Evaluation, Vancouver Coastal Health Research Institute, Vancouver, Canada

⁵ Academic Department of Physiotherapy, Division of Health and Social Care Research, King's College London, London, UK

⁶ McGill University, Montreal, Canada

Introduction

Hip fractures are significant injuries that typically occur in seniors; the incidence of hip fracture is increasing with our aging population [1–3]. In addition to the substantial risk of mortality and morbidity faced by patients after a hip fracture [4–6], healthcare costs are also considerable [7, 8]. In 2005, patients with hip fractures made up only 15% of overall osteoporotic fractures in the USA, but represented 72% of the costs associated with the care of patients with osteoporotic fracture [9]. To date, much of the interventional research to improve patient outcomes after hip fracture has focused on acute care management strategies such as time to surgery [10, 11].

Recovery after hip fracture is prolonged extending over at least the first year, well beyond the care delivered in the surgical hospital setting [12–14]. There is growing recognition that care delivery in the post-acute period, including where patients are discharged to after the surgical hospital stay, is under-studied and whose impact on patient recovery and health services utilization is unclear [15–25]. We hypothesized that, in Canada, hospital type (teaching versus community) might affect discharge decision-making with teaching hospitals more likely to send patients to other sites for further recovery (i.e., rehabilitation or acute care) due to surgical bed demands and availability of community-based care. Unlike the USA, the use of post-acute rehabilitation or skilled nursing facilities is uncommon in Canada [26]. Dis-coordinated discharge decision-making is, at least, in part due to current payment policies where providers across different settings are not incentivized to coordinate care [27, 28]. Instead, care providers work in their respective treatment silos where payment is given for setting-specific services [29]. This may lead to inefficient use of health resources where total episode of care length of stay increases or avoidable readmissions occur [30]. In Ontario, Pitzul et al. [31] identified 49 unique post-discharge pathways utilized following hip fracture with significant regional and hospital variation. Although inpatient rehabilitation was associated with higher initial costs, patients who were discharged to the community were more likely to be readmitted or die within 1 year of hip fracture than patients discharged to inpatient rehabilitation [32]. Others have reported similar findings both in the USA and internationally; rehabilitation may be associated with increased costs, but often leads to improved patient outcomes [19, 33–35].

Thus, data are lacking on how discharge destination varies after discharge from the surgical hospital following hip fracture surgery. Knowledge of variability in discharge practices can inform future policy on post-acute care pathways to enhance patient recovery while utilizing constrained health resources efficiently. The objective of this study was to determine, among previously community-dwelling patients surgically treated for first hip fracture in Canada (excluding Quebec), the following: (1) the proportion of discharge

destinations in each province; (2) the standardized difference in discharge destinations between each province and Ontario, adjusting for patient, injury, and care characteristics; and (3) the role of hospital type in inter-provincial differences in discharge destinations. We hypothesized that there would be significant variation in discharge destinations after hip fracture across provinces, even after risk adjustment and that this variation would be mitigated by hospital type.

Methods

Study and patient setting

We examined discharge abstracts of 121,507 previously community-dwelling patients 65 years or older who underwent surgery for non-pathological first hip fracture between January 1, 2004, and December 31, 2012, in Canadian hospitals, except for the province of Quebec. Quebec compiles hospital discharge data in a separate database and does not contribute to the Canadian Institute for Health Information (CIHI) Discharge Abstracts Database. We focused on those who received surgery as only a small number ($n = 5396$ [4.4%]) did not receive surgical intervention (see [Supplementary Tables](#) for discharge destination of non-operative patients). Abstracts were selected from the Discharge Abstract Database maintained by the CIHI using procedure codes for hip fracture surgery (CCI: 1VA74^^, 1VA53^^, 1VC74^^, 1SQ53^^, or CCP: 9054, 9114, 9134, 9351, 9359, 9361, 9362, 9363, 9364, 9369). Multiple abstracts with the same patient identifier were combined into one care episode using the CIHI rules for hospital transfers [36, 37]. We considered patients to have been previously community dwelling if they were admitted from home or home with support services. Admissions from acute care, ambulatory care, or the emergency department were also considered proxies for previously community-dwelling patients as these settings are not pre-admission residences. In provinces with larger rural populations, patients were frequently initially admitted to non-surgical hospitals and then discharged and admitted to a surgical hospital, which could be several hours later due to transfer time. These were not considered as “preoperative transfers,” but rather “admissions from acute care settings” [36, 37]. Patients were not considered community dwelling if they were admitted from continuing care (i.e., long-term residential care), which included rehabilitation facilities, chronic care facilities, and nursing homes. From our initial cohort of 121,507 potentially eligible patients, we excluded 7716 patients who died in hospital after hip fracture and 1839 patients who underwent surgery in a hospital with an annual surgical volume of less than 24 surgeries [38]. This resulted in an analytical cohort of 111,952 community-dwelling patients who were surgically treated for first hip fracture.

Outcome

The outcome was discharge destination from the surgical hospital, categorized as rehabilitation, home, acute care, continuing care, and other. CIHI “discharge disposition” codes were used to identify discharges to home (discharged home or discharged to home setting with support services), acute care (transferred to another facility providing inpatient hospital care), continuing care (transferred to another facility or level of care other than acute), and other (transferred to palliative care or a hospice, signed out against medical advice, or did not return from pass). CIHI “institution to type” codes indicating transfer to a general or special rehabilitation facility were used to capture discharges to rehabilitation. CIHI re-abstraction shows high reliability for both “discharge disposition” and “institution to type” data elements that were used to characterize the outcome [39]. We estimated the proportion of patients who were discharged to each destination in the overall study population and by province. Due to the small number of patients discharged to other, we did not report the effect of province on this outcome category.

Study variable

The province where surgery was performed was the primary study variable. We coded the available provinces as follows: Alberta, British Columbia, Manitoba, New Brunswick, Newfoundland and Labrador, Nova Scotia, Ontario, Prince Edward Island, and Saskatchewan. Ontario was the reference province as nearly half of all patients underwent surgery in that province (47.4%). The secondary study variable was hospital type at surgery: teaching hospital, large community hospital, or medium community hospital. CIHI classifies non-teaching community hospitals based on the yearly volume of inpatient cases, total weighted cases, and inpatient days [40]. Community hospitals are classified as large if they meet two of the following three criteria: more than 8000 inpatient cases; more than 10,000 weighted cases; more than 50,000 inpatient days. Community hospitals are classified as medium if they have a volume of 2000 weighted cases or more.

Statistical analysis

We reported frequencies and percentages of patient characteristics, the type of fracture, and care delivery across provinces.

The proportion of discharge to rehabilitation, home, acute care, and continuing care was estimated by dividing the number of discharges to each destination by the number of total discharges in the overall population or in a given province. We used Pearson’s chi-square test of independence to compare the distribution of discharge destinations across provinces. We used a multinomial logistic regression model to test the difference in proportions of discharge to rehabilitation, acute care,

continuing care versus home between the provinces and Ontario. We adjusted for age (< 85 years or 85 years and older), sex, comorbidity (none or at least one) [16, 41], fracture type (transcervical or trochanteric), timing of admission (early weekday, late weekday, or weekend), preoperative transfer history (none or at least one hospital transfer occurring between the date of initial hospitalization and the date of hip fracture surgery) [36], preoperative procedures (none or at least one procedure before hip fracture surgery not related to a medical reason for delay) [42], hospital volume (above or below median number of hip fracture surgeries at the treating hospital during the fiscal year when the patient was admitted), excessive surgical demand (excessive demand defined as the number of admissions for hip fracture in the week of initial hospitalization exceeding the hospital weekly capacity for surgery), surgery type (internal fixation or implantation/other), and timing of surgery (day of admission, admission day 2, admission day 3, admission day 4, or after 4 days) and by calendar year of admission. Standard errors of the regression coefficients were estimated by relaxing the independence assumption between patients within hospitals. Based on regression analysis, we estimated absolute risk differences and risk ratios of discharge destination for each province compared with Ontario, standardizing the distribution of patient, injury, and care characteristics to that of Ontario [43].

In secondary analysis, we additionally adjusted for hospital type at surgery in our multinomial logistic regression model to determine if provincial variation would persist. Stata release 15 was used for statistical analyses [44].

Results

Patient characteristics

Most patients were women (73.7%) and were less than 85 years old (59.8%) (Table 1). More than half of patients were admitted for transcervical hip fracture (52.8%) and a quarter of patients had at least one comorbidity (25.0%). Most patients underwent internal fixation (59.8%), underwent surgery at a large community hospital (44.9%), were treated on inpatient day 2 (42.2%), and did not experience a preoperative transfer (91.0%). Alberta and Saskatchewan exhibited the largest proportion of surgeries at a teaching hospital (73.4% and 91.9% respectively).

Discharge destination overall and by province

Overall, 22.5% of patients were discharged to rehabilitation, 31.6% to home, 27.0% to acute care, 18.2% to continuing care, and 0.8% to other (Table 2). These proportions varied significantly across provinces ($X^2 = 33,284.2$, $df = 32$, $p < 0.001$) (Fig. 1). Ontario most frequently discharged

Table 1 Characteristics of 111,952 community-dwelling patients surgically treated for first hip fracture in 2004–2012, overall and for the Canadian Provinces, excluding Quebec

	Province; No. (%) of Patients																			
	Overall (N = 111,952)	AB (N = 12,232)	BC (N = 21,389)	MB (N = 6302)	NB (N = 4137)	NL (N = 2666)	NS (N = 5210)	ON (N = 53,032)	PE (N = 772)	SK (N = 6212)										
	n	%	n	%	n	%	n	%	n	%	n	%								
Age																				
< 85	66,979	(59.8)	12,366	(57.8)	3537	(56.1)	2399	(58.0)	1691	(63.4)	3027	(58.1)	32,561	(61.4)	466	(60.4)	3392	(54.6)		
≥ 85	44,973	(40.2)	4692	(38.4)	9023	(42.2)	2765	(43.9)	1738	(42.0)	975	(36.6)	2183	(41.9)	20,471	(38.6)	306	(39.6)	2820	(45.4)
Sex ^a																				
Women	82,469	(73.7)	8837	(72.2)	15,525	(72.6)	4665	(74.0)	3193	(77.2)	2110	(79.1)	3983	(76.4)	39,023	(73.6)	600	(77.7)	4533	(73.0)
Men	29,470	(26.3)	3393	(27.7)	5862	(27.4)	1637	(26.0)	943	(22.8)	556	(20.9)	1227	(23.6)	14,001	(26.4)	172	(22.3)	1679	(27.0)
Comorbidity																				
None	84,013	(75.0)	9134	(74.7)	16,235	(75.9)	4789	(76.0)	3014	(72.9)	2040	(76.5)	4054	(77.8)	39,263	(74.0)	609	(78.9)	4875	(78.5)
At least one ^b	27,939	(25.0)	3098	(25.3)	5154	(24.1)	1513	(24.0)	1123	(27.1)	626	(23.5)	1156	(22.2)	13,769	(26.0)	163	(21.1)	1337	(21.5)
Fracture type																				
Transcervical	59,126	(52.8)	6519	(53.3)	11,524	(53.9)	3348	(53.1)	2301	(55.6)	1496	(56.1)	2855	(54.8)	27,212	(51.3)	442	(57.3)	3429	(55.2)
Intertrochanteric or subtrochanteric	52,826	(47.2)	5713	(46.7)	9865	(46.1)	2954	(46.9)	1836	(44.4)	1170	(43.9)	2355	(45.2)	25,820	(48.7)	330	(42.7)	2783	(44.8)
Timing of admission ^c																				
Early weekday	37,126	(33.2)	4149	(33.9)	7117	(33.3)	2162	(34.3)	1353	(32.7)	951	(35.7)	1955	(37.5)	17,407	(32.8)	263	(34.1)	1769	(28.5)
Late weekday	43,561	(38.9)	4657	(38.1)	8445	(39.5)	2358	(37.4)	1670	(40.4)	1029	(38.6)	1725	(33.1)	20,779	(39.2)	265	(34.3)	2633	(42.4)
Weekend	31,211	(27.9)	3426	(28.0)	5826	(27.2)	1782	(28.3)	1113	(26.9)	686	(25.7)	1530	(29.4)	14,845	(28.0)	218	(28.2)	1785	(28.7)
Preoperative transfer history ^d																				
No	101,914	(91.0)	10,936	(89.4)	18,957	(88.6)	5535	(87.8)	3960	(95.7)	2554	(95.8)	4997	(95.9)	48,820	(92.1)	741	(96.0)	5414	(87.2)
Yes	10,038	(9.0)	1296	(10.6)	2432	(11.4)	767	(12.2)	177	(4.3)	112	(4.2)	213	(4.1)	4212	(7.9)	31	(4.0)	798	(12.8)
Preoperative procedures ^e																				
No	99,749	(89.1)	10,907	(89.2)	20,050	(93.7)	5895	(93.5)	3846	(93.0)	2414	(90.5)	4836	(92.8)	45,228	(85.3)	725	(93.9)	5848	(94.1)
Yes	12,203	(10.9)	1325	(10.8)	1339	(6.3)	407	(6.5)	291	(7.0)	252	(9.5)	374	(7.2)	7804	(14.7)	47	(6.1)	364	(5.9)
Hospital type at treatment ^f																				
Teaching	44,424	(39.7)	8984	(73.4)	8638	(40.4)	587	(9.3)	901	(21.8)	1663	(62.4)	2202	(42.3)	15,739	(29.7)	0	(0.0)	5710	(91.9)
Community-large	50,279	(44.9)	2984	(24.4)	9960	(46.6)	4511	(71.6)	2249	(54.4)	453	(17.0)	934	(17.9)	28,140	(53.1)	772	(100.0)	276	(4.4)
Community-medium	16,149	(14.4)	264	(2.2)	2791	(13.0)	1204	(19.1)	987	(23.9)	550	(20.6)	2074	(39.8)	8053	(15.2)	0	(0.0)	226	(3.6)
Hospital volume ^g																				
Below median in its type	27,646	(24.7)	2170	(17.7)	6412	(30.0)	1196	(19.0)	2921	(70.6)	453	(17.0)	151	(2.9)	12,735	(24.0)	621	(80.4)	987	(15.9)
Above median in its type	84,306	(75.3)	10,062	(82.3)	14,977	(70.0)	5106	(81.0)	1216	(29.4)	2213	(83.0)	5059	(97.1)	40,297	(76.0)	151	(19.6)	5225	(84.1)
Excess surgical demand ^h																				

Table 1 (continued)

	Province; No. (%) of Patients																			
	AB (N = 12,232)		BC (N = 21,389)		MB (N = 6302)		NB (N = 4137)		NL (N = 2666)		NS (N = 5210)		ON (N = 53,032)		PE (N = 772)		SK (N = 6212)			
	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%		
Overall (N = 111,952)																				
No	34,019	(30.4)	4189	(34.2)	6349	(29.7)	1935	(30.7)	1017	(24.6)	801	(30.0)	1657	(31.8)	15,854	(29.9)	207	(26.8)	2010	(32.4)
Yes	77,933	(69.6)	8043	(65.8)	15,040	(70.3)	4367	(69.3)	3120	(75.4)	1865	(70.0)	3553	(68.2)	37,178	(70.1)	565	(73.2)	4202	(67.6)
Type of surgery																				
Internal fixation	66,994	(59.8)	7122	(58.2)	13,289	(62.1)	3947	(62.6)	2262	(54.7)	1545	(58.0)	3084	(59.2)	31,693	(59.8)	391	(50.6)	3661	(58.9)
Implantation or other	44,958	(40.2)	5110	(41.8)	8100	(37.9)	2355	(37.4)	1875	(45.3)	1121	(42.0)	2126	(40.8)	21,339	(40.2)	381	(49.4)	2551	(41.1)
Timing of surgery																				
Admission day	24,759	(22.1)	2349	(19.2)	4511	(21.1)	1058	(16.8)	1387	(33.5)	642	(24.1)	1608	(30.9)	12,155	(22.9)	266	(34.5)	783	(12.6)
Inpatient day 2	47,209	(42.2)	5377	(44.0)	9677	(45.2)	2928	(46.5)	1571	(38.0)	1093	(41.0)	1823	(35.0)	21,817	(41.1)	262	(33.9)	2661	(42.8)
Inpatient day 3	23,306	(20.8)	2766	(22.6)	4596	(21.5)	1197	(19.0)	605	(14.6)	560	(21.0)	882	(16.9)	11,048	(20.8)	140	(18.1)	1512	(24.3)
Inpatient day 4	8335	(7.4)	946	(7.7)	1452	(6.8)	420	(6.7)	285	(6.9)	188	(7.1)	395	(7.6)	3988	(7.5)	61	(7.9)	600	(9.7)
After 4 days	8343	(7.5)	794	(6.5)	1153	(5.4)	699	(11.1)	289	(7.0)	183	(6.9)	502	(9.6)	4024	(7.6)	43	(5.6)	656	(10.6)
Year of admission																				
2004–2006	38,159	(34.1)	3971	(32.5)	7052	(33.0)	1973	(31.3)	1455	(35.2)	862	(32.3)	1731	(33.2)	18,732	(35.3)	254	(32.9)	2129	(34.3)
2007–2009	36,345	(32.5)	4059	(33.2)	6955	(32.5)	2096	(33.3)	1396	(33.7)	897	(33.6)	1755	(33.7)	16,928	(31.9)	257	(33.3)	2002	(32.2)
2010–2012	37,448	(33.5)	4202	(34.4)	7382	(34.5)	2233	(35.4)	1286	(31.1)	907	(34.0)	1724	(33.1)	17,372	(32.8)	261	(33.8)	2081	(33.5)

AB, Alberta; BC, British Columbia; MB, Manitoba; NB, New Brunswick; NL, Newfoundland and Labrador; NS, Nova Scotia; ON, Ontario; PE, Prince Edward Island; SK, Saskatchewan

^a For 13 patients, sex was unknown

^b Comorbidities included heart failure, chronic obstructive pulmonary disorder, ischemic heart disease (acute and chronic), dysrhythmias, hypertension, diabetes, and cancer (breast—female, prostate, renal, lung, multiple myeloma, and metastatic cancer) identified by diagnostic codes from all hospitalizations in 1 year prior to index admission, and cancer and Paget’s disease, identified by diagnostic codes from all hospitalizations during the hip fracture care episode

^c For 54 patients, timing of admission was unknown

^d Defined as at least one hospital transfer occurring between the date of initial hospitalization and the date of hip fracture surgery

^e Defined as at least one intervention before hip fracture surgery not related to a medical reason for delay

^f For 1100 patients, hospital type was unassigned

^g Defined as the number of hip fracture surgeries at the treating hospital during the fiscal year when the patient was admitted

^h Measured as the number of admissions for hip fracture in the week of initial hospitalization compared with the hospital weekly capacity for surgery

Table 2 Discharge destination for community-dwelling patients surgically treated for first hip fracture, for overall, and for the Canadian Provinces

Discharge destination	Province; No. (%) of patients									
	Overall (N = 111,952)	AB (N = 12,232)	BC (N = 21,389)	MB (N = 6302)	NB (N = 4137)	NL (N = 2666)	NS (N = 5210)	ON (N = 53,032)	PE (N = 772)	SK (N = 6212)
Rehabilitation	25,193 (22.5)	431 (3.5)	510 (2.4)	1312 (20.8)	224 (5.4)	292 (11.0)	247 (4.7)	21,759 (41.0)	115 (14.9)	303 (4.9)
Home	35,353 (31.6)	3424 (28.0)	11,169 (52.2)	2057 (32.6)	1563 (37.8)	1251 (46.9)	2480 (47.6)	11,811 (22.3)	157 (20.3)	1441 (23.2)
Acute care	30,211 (27.0)	7118 (58.2)	4869 (22.8)	2329 (37.0)	1390 (33.6)	758 (28.4)	1649 (31.7)	8041 (15.2)	404 (52.3)	3653 (58.8)
Continuing care	20,327 (18.2)	1147 (9.4)	4545 (21.2)	587 (9.3)	947 (22.9)	315 (11.8)	819 (15.7)	11,120 (21.0)	72 (9.3)	775 (12.5)
Other ^a	868 (0.8)	112 (0.9)	296 (1.4)	17 (0.3)	13 (0.3)	50 (1.9)	15 (0.3)	301 (0.6)	24 (3.1)	40 (0.6)

AB, Alberta; BC, British Columbia; MB, Manitoba; NB, New Brunswick; NL, Newfoundland and Labrador; NS, Nova Scotia; ON, Ontario; PE, Prince Edward Island; SK, Saskatchewan

^a Patients were classified as discharged to other if transferred to palliative care or a hospice, signed out against medical advice, or did not return from pass

patients to rehabilitation (41.0%). Discharge to home was most prevalent in British Columbia (52.2%), Nova Scotia (47.6%), Newfoundland and Labrador (46.9%), and New Brunswick (37.8%). Most patients were commonly discharged to acute care in Saskatchewan (58.8%), Alberta (58.2%), Prince Edward Island (52.3%), and Manitoba (37.0%).

Discharge to rehabilitation

The proportion of discharge to rehabilitation ranged from 2.4% in British Columbia to 41.0% in Ontario (Fig. 1). Compared with Ontario, the standardized proportion of discharge to rehabilitation was significantly lower in all provinces, with the greatest differences in British Columbia (difference = -39.9%; 95% CI -46.0, -33.8), Alberta (difference = -39.2; 95% CI -45.3, -33.2), Saskatchewan (difference = -38.2; 95% CI -44.9, -31.4), Nova Scotia (difference = -38.1; 95% CI -46.0, -30.1), New Brunswick (difference = -36.5; 95% CI -44.6, -28.4), and Newfoundland and Labrador (difference = -32.1; 95% CI -41.2, -23.0) (Table 3). Statistical significance was consistent between risk differences and risk ratios.


Discharge to home

The proportion of discharge to home ranged from 20.3% in Prince Edward Island to 52.2% in British Columbia (Fig. 1). Compared with Ontario, the proportion of discharge to home was higher in British Columbia (difference = 32.2%; 95% CI 23.2, 41.3), Nova Scotia (difference = 23.8%; 95% CI 6.6, 41.0), Newfoundland and Labrador (difference = 22.9%; 95% CI 15.8, 30.0), New Brunswick (difference = 14.7%; 95% CI 6.7, 22.7), and Manitoba (difference = 11.6%; 95% CI 4.8, 18.4), and was significantly lower in Prince Edward Island (difference = -4.0%; 95% CI -7.1, -1.0) (Table 3). Statistical significance was consistent between risk differences and risk ratios.

Discharge to acute care

The proportion of discharge to acute care ranged from 15.2% in Ontario to 58.8% in Saskatchewan (Fig. 1). Compared with Ontario, the proportion of discharge to acute care after standardization was higher in Saskatchewan (difference = 47.4%; 95% CI 34.6, 60.1), Alberta (difference = 47.0%; 95% CI 40.9, 53.0), Prince Edward Island (difference = 42.4%; 95% CI 38.4, 46.3), Manitoba (difference = 22.4%; 95% CI 7.8, 37.0), New Brunswick (difference = 21.9%; 95% CI 13.1, 30.7), Nova Scotia (difference = 21.1%; 95% CI 3.1, 39.1), and Newfoundland and Labrador (difference = 17.7%; 95% CI 12.7, 22.8) (Table 3). Statistical significance was consistent between risk differences and risk ratios.

Fig. 1 Percentage of patients discharged by destination across the Canadian provinces. Discharge destination represented by shade, percentage represented by the length of bars. Provinces are in ascending order of percentage for discharge to rehabilitation. Percentage of patients discharged to destination other than rehabilitation, home, acute care, or continuing care not shown (transferred to palliative care or hospice, signed out against medical advice, or did not return from pass). AB, Alberta; BC, British Columbia; MB, Manitoba; NB, New Brunswick; NL, Newfoundland and Labrador; NS, Nova Scotia; ON, Ontario; PE, Prince Edward Island; SK, Saskatchewan



Discharge to continuing care

The proportion of discharge to continuing care ranged from 9.3% in Manitoba to 22.9% in New Brunswick (Fig. 1). Compared with Ontario, the standardized proportion of discharge to continuing care was lower in Prince Edward Island (difference = -12.8%; 95% CI -15.1, -10.5), Alberta (difference = -12.2%; 95% CI -15.2, -9.2), Manitoba (difference = -12.1%; 95% CI -14.9, -9.3), Newfoundland and Labrador (difference = -9.8%; 95% CI -12.8, -6.8), Saskatchewan (difference = -9.4%; 95% CI -15.5, -3.3), and Nova Scotia (difference = -6.6%; 95% CI -11.6, -1.7) (Table 3). Statistical significance was consistent between risk differences and risk ratios.

Discharge destination by province, accounting for hospital type

After adjusting for hospital type, the direction of the provincial effect on discharge to continuing care changed in British Columbia and New Brunswick (Table 4). The statistical significance of all provincial effects on discharge destination remained consistent with the primary analysis.

Discussion

We found significant variation in discharge destinations from the surgical hospital after hip fracture in Canada even after standardizing patient, injury, and care characteristics.

Discharge to rehabilitation accounted for less than 6% of patient discharges in five provinces. Within these five provinces, discharge destinations were commonly directly to home or transfers to other acute care facilities. Discharge from the surgical hospital to continuing care, which typically indicates a transition from the community to permanent residential care, was also heterogeneous, varying from as low as 9.3% to as high as 22.9%. Based on our previous work, where care and outcomes of hip fracture patients varied by hospital type (e.g., length of stay and mortality) [45, 46], we hypothesized that accounting for teaching versus community hospital would mitigate provincial variation, but hospital type appeared to have little impact on differences in discharge destinations. This may suggest that discharge destination is resource driven (i.e., availability of beds or rehabilitation access) rather than driven by clinical personnel decision-making or hospital status.

Pitzul et al.'s recent work raises concern regarding our findings as patients in Ontario who received post-acute inpatient rehabilitation had lower mortality and readmissions within the first year after hip fracture than those who were discharged directly to the community [32]. A recent systematic review also identified that both patient and facility factors lead to increased readmissions after hip fracture [30]. International reports support improved patient outcomes from discharge to rehabilitation facilities, but at a higher direct cost than discharge to home [15, 19, 33, 35, 47].

In Canada, inpatient rehabilitation facilities for patients with hip fractures are very limited outside of Ontario. In a direct comparison of Canada and the USA using clinical trial

Table 3 Relation between the Canadian provinces and discharge destinations among community-dwelling patients surgically treated for first hip fracture

Province	No. of patients	No. of discharges	% of discharges	Unadjusted RD, % (95% CI)	Adjusted RD, % (95% CI) ^{ab}	Unadjusted RR (95% CI)	Adjusted RR (95% CI) ^{ab}
Discharge to rehabilitation							
ON	53,032	21,759	41.0	–	–	1.00	1.00
AB	12,232	431	3.5	–37.5 (–43.8, –31.2)	–39.2 (–45.3, –33.2)	0.09 (0.08, 0.09)	0.07 (0.00, 0.14)
BC	21,389	510	2.4	–38.6 (–44.8, –32.5)	–39.9 (–46.0, –33.8)	0.06 (0.05, 0.06)	0.06 (0.00, 0.13) ^c
MB	6302	1312	20.8	–20.2 (–30.8, –9.6)	–21.7 (–31.4, –12.0)	0.51 (0.48, 0.53)	0.49 (0.29, 0.69)
NB	4137	224	5.4	–35.6 (–43.3, –28.0)	–36.5 (–44.6, –28.4)	0.13 (0.12, 0.15)	0.14 (0.00, 0.28) ^c
NL	2666	292	11.0	–30.1 (–41.4, –18.8)	–32.1 (–41.2, –23.0)	0.27 (0.24, 0.30)	0.24 (0.06, 0.42)
NS	5210	247	4.7	–36.3 (–44.4, –28.2)	–38.1 (–46.0, –30.1)	0.12 (0.10, 0.13)	0.10 (0.00, 0.24) ^c
PE	772	115	14.9	–26.1 (–31.5, –20.8)	–27.8 (–33.3, –22.3)	0.36 (0.30, 0.42)	0.34 (0.29, 0.39)
SK	6212	303	4.9	–36.2 (–42.9, –29.4)	–38.2 (–44.9, –31.4)	0.12 (0.11, 0.13)	0.10 (0.00, 0.20) ^c
Discharge to home							
ON	53,032	11,811	22.3	–	–	1.00	1.00
AB	12,232	3424	28.0	5.7 (–2.5, 14.0)	4.0 (–3.7, 11.7)	1.26 (1.22, 1.30)	1.18 (0.83, 1.53)
BC	21,389	11,169	52.2	29.9 (21.3, 38.6)	32.2 (23.2, 41.3)	2.34 (2.30, 2.39)	2.42 (1.94, 2.91)
MB	6302	2057	32.6	10.4 (4.3, 16.5)	11.6 (4.8, 18.4)	1.47 (1.41, 1.52)	1.51 (1.18, 1.85)
NB	4137	1563	37.8	15.5 (8.3, 22.7)	14.7 (6.7, 22.7)	1.70 (1.62, 1.77)	1.65 (1.26, 2.04)
NL	2666	1251	46.9	24.7 (17.6, 31.7)	22.9 (15.8, 30.0)	2.11 (2.02, 2.20)	2.01 (1.63, 2.40)
NS	5210	2480	47.6	25.3 (8.9, 41.7)	23.8 (6.6, 41.0)	2.14 (2.07, 2.21)	2.05 (1.26, 2.85)
PE	772	157	20.3	–1.9 (–4.7, 0.9)	–4.0 (–7.1, –1.0)	0.91 (0.78, 1.04)	0.82 (0.71, 0.93)
SK	6212	1441	23.2	0.9 (–4.4, 6.2)	0.0 (–6.4, 6.5)	1.04 (0.99, 1.09)	1.00 (0.72, 1.29)
Discharge to acute care							
ON	53,032	8041	15.2	–	–	1.00	1.00
AB	12,232	7118	58.2	43.0 (36.8, 49.2)	47.0 (40.9, 53.0)	3.84 (3.74, 3.93)	4.50 (3.23, 5.78)
BC	21,389	4869	22.8	7.6 (–1.2, 16.4)	6.5 (–1.7, 14.7)	1.50 (1.45, 1.55)	1.48 (0.81, 2.16)
MB	6302	2329	37.0	21.8 (7.2, 36.4)	22.4 (7.8, 37.0)	2.44 (2.34, 2.53)	2.67 (1.39, 3.95)
NB	4137	1390	33.6	18.4 (9.6, 27.3)	21.9 (13.1, 30.7)	2.22 (2.11, 2.32)	2.63 (1.72, 3.55)
NL	2666	758	28.4	13.3 (6.5, 20.0)	17.7 (12.7, 22.8)	1.88 (1.76, 1.99)	2.32 (1.64, 3.00)
NS	5210	1649	31.7	16.5 (–1.6, 34.6)	21.1 (3.1, 39.1)	2.09 (1.99, 2.18)	2.58 (1.08, 4.07)
PE	772	404	52.3	37.2 (33.0, 41.3)	42.4 (38.4, 46.3)	3.45 (3.21, 3.69)	4.16 (3.04, 5.29)
SK	6212	3653	58.8	43.6 (32.0, 55.3)	47.4 (34.6, 60.1)	3.88 (3.77, 3.99)	4.53 (3.01, 6.06)
Discharge to continuing care							
ON	53,032	11,120	21.0	–	–	1.00	1.00
AB	12,232	1147	9.4	–11.6 (–14.5, –8.7)	–12.2 (–15.2, –9.2)	0.45 (0.42, 0.47)	0.42 (0.32, 0.53)
BC	21,389	4545	21.2	0.3 (–5.2, 5.8)	0.3 (–5.2, 5.7)	1.01 (0.98, 1.04)	1.01 (0.75, 1.27)
MB	6302	587	9.3	–11.7 (–14.3, –9.0)	–12.1 (–14.9, –9.3)	0.44 (0.41, 0.48)	0.43 (0.33, 0.52)
NB	4137	947	22.9	1.9 (–7.2, 11.1)	0.1 (–9.0, 9.3)	1.09 (1.03, 1.16)	1.01 (0.57, 1.44)
NL	2666	315	11.8	–9.2 (–12.2, –6.1)	–9.8 (–12.8, –6.8)	0.56 (0.50, 0.62)	0.53 (0.42, 0.64)
NS	5210	819	15.7	–5.2 (–10.2, –0.3)	–6.6 (–11.6, –1.7)	0.75 (0.70, 0.80)	0.69 (0.47, 0.91)
PE	772	72	9.3	–11.6 (–13.9, –9.4)	–12.8 (–15.1, –10.5)	0.44 (0.35, 0.54)	0.39 (0.35, 0.44)
SK	6212	775	12.5	–8.5 (–14.3, –2.6)	–9.4 (–15.5, –3.3)	0.59 (0.55, 0.64)	0.55 (0.28, 0.83)

RD, risk difference; CI, confidence interval; RR, risk ratio; AB, Alberta; BC, British Columbia; MB, Manitoba; NB, New Brunswick; NL, Newfoundland and Labrador; NS, Nova Scotia; ON, Ontario; PE, Prince Edward Island; SK, Saskatchewan

^a Standardized to the distribution of age, sex, comorbidity, fracture type, timing of admission, preoperative transfer history, preoperative procedures, hospital volume, surgical demand, surgery type, timing of surgery, and calendar year of admission of Ontario

^b Excludes 13 patients with unknown sex and 54 patients with unknown timing of admission

^c Lower confidence limit truncated at 0

Table 4 Relation between the Canadian provinces and discharge destinations among community-dwelling patients surgically treated for first hip fracture, secondary analysis adjusting for hospital type

Province	No. of patients	No. of discharges	% of discharges	Unadjusted RD, % (95% CI)	Adjusted RD, % (95% CI) ^{ab}	Unadjusted RR (95% CI)	Adjusted RR (95% CI) ^{ab}
Discharge to rehabilitation							
ON	53,032	21,759	41.0	–	–	1.00	1.00
AB	12,232	431	3.5	–37.5 (–43.8, –31.2)	–39.3 (–45.1, –33.5)	0.09 (0.08, 0.09)	0.07 (0.00, 0.14) ^c
BC	21,389	510	2.4	–38.6 (–44.8, –32.5)	–40.0 (–45.8, –34.2)	0.06 (0.05, 0.06)	0.05 (0.00, 0.12) ^c
MB	6302	1312	20.8	–20.2 (–30.8, –9.6)	–21.0 (–31.0, –11.1)	0.51 (0.48, 0.53)	0.50 (0.29, 0.72)
NB	4137	224	5.4	–35.6 (–43.3, –28.0)	–37.4 (–44.6, –30.3)	0.13 (0.12, 0.15)	0.11 (0.00, 0.24) ^c
NL	2666	292	11.0	–30.1 (–41.4, –18.8)	–31.9 (–41.7, –22.1)	0.27 (0.24, 0.30)	0.25 (0.04, 0.45)
NS	5210	247	4.7	–36.3 (–44.4, –28.2)	–36.8 (–46.1, –27.4)	0.12 (0.10, 0.13)	0.13 (0.00, 0.32) ^c
PE	772	115	14.9	–26.1 (–31.5, –20.8)	–31.8 (–37.3, –26.4)	0.36 (0.30, 0.42)	0.25 (0.19, 0.30)
SK	6212	303	4.9	–36.2 (–42.9, –29.4)	–37.8 (–44.1, –31.4)	0.12 (0.11, 0.13)	0.11 (0.01, 0.20)
Discharge to home							
ON	53,032	11,811	22.3	–	–	1.00	1.00
AB	12,232	3424	28.0	5.7 (–2.5, 14.0)	4.1 (–3.2, 11.4)	1.26 (1.22, 1.30)	1.18 (0.85, 1.50)
BC	21,389	11,169	52.2	29.9 (21.3, 38.6)	32.1 (23.0, 41.2)	2.34 (2.30, 2.39)	2.39 (1.91, 2.87)
MB	6302	2057	32.6	10.4 (4.3, 16.5)	12.0 (4.5, 19.5)	1.47 (1.41, 1.52)	1.52 (1.17, 1.88)
NB	4137	1563	37.8	15.5 (8.3, 22.7)	13.4 (5.8, 21.0)	1.70 (1.62, 1.77)	1.58 (1.22, 1.95)
NL	2666	1251	46.9	24.7 (17.6, 31.7)	24.7 (15.4, 34.0)	2.11 (2.02, 2.20)	2.07 (1.61, 2.54)
NS	5210	2480	47.6	25.3 (8.9, 41.7)	29.5 (15.4, 43.6)	2.14 (2.07, 2.21)	2.28 (1.62, 2.95)
PE	772	157	20.3	–1.9 (–4.7, 0.9)	–7.3 (–11.1, –3.5)	0.91 (0.78, 1.04)	0.68 (0.55, 0.82)
SK	6212	1441	23.2	0.9 (–4.4, 6.2)	1.4 (–5.6, 8.3)	1.04 (0.99, 1.09)	1.06 (0.75, 1.37)
Discharge to acute care							
ON	53,032	8041	15.2	–	–	1.00	1.00
AB	12,232	7118	58.2	43.0 (36.8, 49.2)	47.2 (41.4, 53.0)	3.84 (3.74, 3.93)	4.72 (3.56, 5.89)
BC	21,389	4869	22.8	7.6 (–1.2, 16.4)	7.1 (–0.9, 15.2)	1.50 (1.45, 1.55)	1.56 (0.88, 2.25)
MB	6302	2329	37.0	21.8 (7.2, 36.4)	21.6 (6.8, 36.5)	2.44 (2.34, 2.53)	2.71 (1.40, 4.01)
NB	4137	1390	33.6	18.4 (9.6, 27.3)	25.3 (17.2, 33.5)	2.22 (2.11, 2.32)	3.00 (2.10, 3.90)
NL	2666	758	28.4	13.3 (6.5, 20.0)	15.5 (6.1, 24.9)	1.88 (1.76, 1.99)	2.22 (1.36, 3.09)
NS	5210	1649	31.7	16.5 (–1.6, 34.6)	12.7 (2.8, 22.6)	2.09 (1.99, 2.18)	2.00 (1.11, 2.89)
PE	772	404	52.3	37.2 (33.0, 41.3)	52.1 (47.0, 57.3)	3.45 (3.21, 3.69)	5.11 (3.88, 6.35)
SK	6212	3653	58.8	43.6 (32.0, 55.3)	45.0 (34.0, 56.0)	3.88 (3.77, 3.99)	4.55 (3.20, 5.90)
Discharge to continuing care							
ON	53,032	11,120	21.0	–	–	1.00	1.00
AB	12,232	1147	9.4	–11.6 (–14.5, –8.7)	–12.4 (–15.6, –9.3)	0.45 (0.42, 0.47)	0.42 (0.31, 0.54)
BC	21,389	4545	21.2	0.3 (–5.2, 5.8)	–0.2 (–5.3, 5.0)	1.01 (0.98, 1.04)	0.99 (0.75, 1.23)
MB	6302	587	9.3	–11.7 (–14.3, –9.0)	–12.4 (–15.2, –9.6)	0.44 (0.41, 0.48)	0.42 (0.34, 0.51)
NB	4137	947	22.9	1.9 (–7.2, 11.1)	–1.1 (–9.9, 7.8)	1.09 (1.03, 1.16)	0.95 (0.54, 1.36)
NL	2666	315	11.8	–9.2 (–12.2, –6.1)	–9.7 (–12.8, –6.6)	0.56 (0.50, 0.62)	0.55 (0.43, 0.67)
NS	5210	819	15.7	–5.2 (–10.2, –0.3)	–5.3 (–10.0, –0.6)	0.75 (0.70, 0.80)	0.75 (0.55, 0.96)
PE	772	72	9.3	–11.6 (–13.9, –9.4)	–14.6 (–17.0, –12.1)	0.44 (0.35, 0.54)	0.32 (0.27, 0.38)
SK	6212	775	12.5	–8.5 (–14.3, –2.6)	–8.8 (–14.7, –2.9)	0.59 (0.55, 0.64)	0.59 (0.33, 0.86)

RD, risk difference; CI, confidence interval; RR, risk ratio; AB, Alberta; BC, British Columbia; MB, Manitoba; NB, New Brunswick; NL, Newfoundland and Labrador; NS, Nova Scotia; ON, Ontario; PE, Prince Edward Island; SK, Saskatchewan

^a Standardized to the distribution of age, sex, comorbidity, fracture type, timing of admission, preoperative transfer history, preoperative procedures, hospital type, hospital volume, surgical demand, surgery type, timing of surgery, and calendar year of admission of Ontario

^b Excludes 13 patients with unknown sex and 54 patients with unknown timing of admission

^c Lower confidence limit truncated at 0

data, the acute care length of stay was higher in Canada with significantly less use of inpatient rehabilitation; however, mortality up to 3 years after hip fracture was significantly lower in Canada [26]. These findings appear contrary to those reported in within-country comparisons, suggesting that further work is required regarding both treatment settings and services provided during the recovery period.

Rehabilitation may also be offered in acute care facilities that are not designated as rehabilitation facilities, or via home care, but it is likely that rehabilitation intensity is lower due to more limited rehabilitation resources relative to rehabilitation facilities. Indeed, a recent national UK audit of rehabilitation after hip fracture surgery reported wait times of up to 3 months for rehabilitation at home following discharge from acute care [48]. If rehabilitation is beneficial to patient outcomes, approaches to deliver timely higher intensity rehabilitation in settings other than formal rehabilitation facilities should be developed as infrastructure for multiple new rehabilitation facilities is likely not feasible for most healthcare systems.

Other Canadian work evaluating total joint arthroplasty and stroke patient populations reported that post-acute service variation led to substantial differences in healthcare costs due to longer length of stays and higher readmissions [27, 29]. Similarly, studies in the USA have also identified substantial variation in post-acute care with resultant differences in both costs and outcomes [18, 19, 23, 33]. Clearly, further evaluation of post-acute care is needed to determine how best to develop and organize post-acute care to maximize patient outcomes while using resources efficiently.

Although this is the first study to evaluate discharge destinations using a national evaluation approach in Canada, there are some limitations. Based on data available, we can describe discharge patterns, but not the patient outcomes associated with these care patterns. We also did not attempt to separate secular trends in care from provincial variation in discharge practices. Canada has provincial physician and hospital-based insurance programs that independently organize patient care, payments, and data reporting, limiting our ability to determine the impact of inter-provincial service variation and organizational changes that might have occurred over time.

We also only followed patients for their initial hospital stay, so we are unable to determine how discharge practices affected patients' ability to ultimately return to live in the community. Data from rehabilitation settings following surgical hospital discharge are not in the national hospital discharge database, so we could not compare overall lengths of stays in facilities after a hip fracture. Non-mandatory data reporting in rehabilitation facilities or reporting of rehabilitation intensity in hospital settings in Canada substantially limits our ability to understand the association between rehabilitation intensity and duration relative to patient outcomes and health service utilization. We excluded patients who underwent surgery in a hospital with an annual volume of < 24 surgeries. This led

to the exclusion of small community hospitals from our analysis. Therefore, results may not be generalizable to patients undergoing surgery at these sites.

Finally, we classified patients admitted from acute, emergency, or ambulatory care as previously community dwelling, which may have led to some misclassification. We made this decision to capture an additional 25,814 patients in the analysis (Table 2 vs. Supplementary Table 2). In particular, we substantially increased the proportion of patients included in the analysis from provinces with large rural populations where patients may take longer to move from a non-surgical hospital to a surgical hospital and, thus, are considered as admissions from acute care rather than transfers. We believe the potential for misclassification is low as patients admitted from continuing care are likely to return to continuing care after their hospital stay [49]. We conducted a series of sensitivity analyses and noted a similar proportion of patients discharged to continuing care among those who were admitted from home or home with support, acute, emergency, or ambulatory care, and those who were admitted from home or home with support. (Supplementary Tables 1–3). We also examined the discharge status of the small proportion (< 5%) of patients who did not receive surgical management in our sensitivity analysis, which also demonstrated similar heterogeneity in discharge patterns (Supplementary Table 4).

This work is foundational as the first to examine discharge patterns following hip fracture across Canada to increase our understanding of the national variation in post-acute recovery pathways. This baseline knowledge will facilitate further evaluations and comparisons across and within provinces and/or countries to determine which pathways are most effective and cost-effective. We demonstrated that there are currently no broadly standardized pathways in provinces' post-acute care delivery with marked heterogeneity in discharge decision-making, similar to that noted in other countries [19, 23, 33, 34]. Further evaluations in various post-acute treatment settings (i.e., rehabilitation, acute care, or home) as well as by rehabilitation intensity (e.g., frequency, duration) and services provided would facilitate evaluation of the impact of various post-discharge pathways on patient outcomes and health system costs.

Acknowledgements LB, BS, PG, JDK, LK, and KJS contributed to the conception and design of the study. SB, PG, LK, and KJS arranged the acquisition of data. JDK and LK conducted the statistical analysis. LB, BS, PG, JDK, LK, KJS, JMS, EH, and SNM contributed to the interpretation of data. LB and JDK drafted the manuscript. LB, BS, PG, JDK, LK, KJS, JMS, EH, and SNM critically revised the manuscript for important intellectual content and approved the final version for submission.

We gratefully acknowledge the guidance from the Canadian Institute for Health Information experts in understanding the discharge abstracts.

Funding source This research was funded by the Canadian Institutes of Health Research (MOP 133629). This funder had no role in the design of this study, execution, analyses, data interpretation, or decision to submit results for publication.

Author contributions The following are members of the Canadian Collaborative Study of Hip Fractures: Lauren Beaupre, Eric Bohm, Michael Dunbar, Donald Griesdale, Pierre Guy, Edward Harvey, Erik Hellsten, Susan Jaglal, Hans Kreder, Lisa Kuramoto, Adrian Levy, Suzanne N. Morin, Katie J. Sheehan, Boris Sobolev, Jason M. Sutherland, and James Waddell.

Compliance with ethical standards

Conflicts of interest The following competing interests are declared: (1) LB is the David Magee Endowed Chair in Musculoskeletal Research supported by the Faculty of Rehabilitation Medicine at the University of Alberta. (2) PG has received grants from the Canadian Institutes of Health Research related to this work. PG also receives funding from the Natural Sciences and Engineering Research Council of Canada, the Canadian Foundation for Innovation, and the British Columbia Specialists Services Committee for work around hip fracture care not related to this manuscript. He has also received fees from the BC Specialists Services Committee (for a provincial quality improvement project on redesign of hip fracture care) and from Stryker Orthopedics (as a product development consultant). He is a board member and shareholder in Traumis Surgical Systems Inc. and a board member for the Canadian Orthopedic Foundation. He also serves on the speakers' bureau of AO Trauma North America and Stryker Canada. (3) SNM reports research grants from Amgen Canada and from Merck. (4) BS, JDK, LK, KJS, JMS, and EH declare they have no conflicts of interest.

References

- Brauer CA, Coca-Perraillon M, Cutler DM, Rosen AB (2009) Incidence and mortality of hip fractures in the United States. *JAMA* 302:1573–1579
- Johnell O, Kanis JA (2006) An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporos Int* 17:1726–1733
- Papadimitropoulos EA, Coyte PC, Josse RG, Greenwood CE (1997) Current and projected rates of hip fracture in Canada. *CMAJ* 157:1357–1363
- Nazrun AS, Tzar MN, Mokhtar SA, Mohamed IN (2014) A systematic review of the outcomes of osteoporotic fracture patients after hospital discharge: morbidity, subsequent fractures, and mortality. *Ther Clin Risk Manag* 10:937–948
- Paksima N, Koval KJ, Aharonoff G, Walsh M, Kubiak EN, Zuckerman JD, Egol KA (2008) Predictors of mortality after hip fracture: a 10-year prospective study. *Bull NYU Hosp Jt Dis* 66:111–117
- Smith T, Pelpola K, Ball M, Ong A, Myint PK (2014) Pre-operative indicators for mortality following hip fracture surgery: a systematic review and meta-analysis. *Age Ageing* 43:464–471
- Haentjens P, Magaziner J, Colon-Emeric CS et al (2010) Meta-analysis: excess mortality after hip fracture among older women and men. *Ann Intern Med* 152:380–390
- Stukel TA, Fisher ES, Alter DA, Guttman A, Ko DT, Fung K, Wodchis WP, Baxter NN, Earle CC, Lee DS (2012) Association of hospital spending intensity with mortality and readmission rates in Ontario hospitals. *JAMA* 307:1037–1045
- Burge R, Dawson-Hughes B, Solomon DH, Wong JB, King A, Tosteson A (2007) Incidence and economic burden of osteoporosis-related fractures in the United States, 2005–2025. *J Bone Miner Res* 22:465–475
- Moja L, Piatti A, Pecoraro V, Ricci C, Virgili G, Salanti G, Gennari L, Liberati A, Banfi G (2012) Timing matters in hip fracture surgery: patients operated within 48 hours have better outcomes. A meta-analysis and meta-regression of over 190,000 patients. *PLoS One* 7:e46175
- Simunovic N, Devereaux PJ, Sprague S, Guyatt GH, Schemitsch E, DeBeer J, Bhandari M (2010) Effect of early surgery after hip fracture on mortality and complications: systematic review and meta-analysis. *CMAJ* 182:1609–1616
- Bertram M, Norman R, Kemp L, Vos T (2011) Review of the long-term disability associated with hip fractures. *Inj Prev* 17:365–370
- Dyer SM, Crotty M, Fairhall N et al (2016) A critical review of the long-term disability outcomes following hip fracture. *BMC Geriatr* 16:158
- Gonzalez-Zabaleta J, Pita-Fernandez S, Seoane-Pillado T, Lopez-Calvino B et al (2015) Dependence for basic and instrumental activities of daily living after hip fractures. *Arch Gerontol Geriatr* 60:66–70
- Armstrong JJ, Zhu M, Hirdes JP, Stolee P (2015) Rehabilitation therapies for older clients of the Ontario home care system: regional variation and client-level predictors of service provision. *Disabil Rehabil* 37:625–631
- Auais M, Morin S, Nadeau L, Finch L, Mayo N (2013) Changes in frailty-related characteristics of the hip fracture population and their implications for healthcare services: evidence from Quebec, Canada. *Osteoporos Int* 24:2713–2724
- Bentler SE, Liu L, Obrizan M, Cook EA, Wright KB, Geweke JF, Chrischilles EA, Pavlik CE, Wallace RB, Ohsfeldt RL, Jones MP, Rosenthal GE, Wolinsky FD (2009) The aftermath of hip fracture: discharge placement, functional status change, and mortality. *Am J Epidemiol* 170:1290–1299
- Blackburn J, Locher JL, Morrisey MA, Becker DJ, Kilgore ML (2016) The effects of state-level expenditures for home- and community-based services on the risk of becoming a long-stay nursing home resident after hip fracture. *Osteoporos Int* 27:953–961
- Buntin MB, Colla CH, Deb P, Sood N, Escarce JJ (2010) Medicare spending and outcomes after postacute care for stroke and hip fracture. *Med Care* 48:776–784
- Cary MP Jr, Pan W, Sloane R, Bettger JP, Hoenig H, Merwin EI, Anderson RA (2016) Self-care and mobility following postacute rehabilitation for older adults with hip fracture: a multilevel analysis. *Arch Phys Med Rehabil* 97:760–771
- DeJong G (2014) Are we asking the right question about postacute settings of care? *Arch Phys Med Rehabil* 95:218–221
- Fisher ES, Wennberg JE (2003) Health care quality, geographic variations, and the challenge of supply-sensitive care. *Perspect Biol Med* 46:69–79
- Fisher ES, Wennberg DE, Stukel TA, Gottlieb DJ, Lucas FL, Pinder EL (2003) The implications of regional variations in Medicare spending. Part 1: the content, quality, and accessibility of care. *Ann Intern Med* 138:273–287
- Huang TT, Liang SH (2005) A randomized clinical trial of the effectiveness of a discharge planning intervention in hospitalized elders with hip fracture due to falling. *J Clin Nurs* 14:1193–1201
- Nordstrom P, Michaelsson K, Hommel A, Norrman PO et al (2016) Geriatric rehabilitation and discharge location after hip fracture in relation to the risks of death and readmission. *J Am Med Dir Assoc* 17:91–97
- Beaupre LA, Wai EK, Hoover DR, Noveck H, Roffey DM, Cook DR, Magaziner JS, Carson JL (2018) A comparison of outcomes between Canada and the United States in patients recovering from hip fracture repair: secondary analysis of the FOCUS trial. *Int J Qual Health Care* 30:97–103

27. Sutherland JM, Hellsten E, Yu K (2012) Bundles: an opportunity to align incentives for continuing care in Canada? *Health Policy* 107:209–217
28. Sutherland JM, Busse R (2016) Canada: focus on a country's health system with provincial diversity. *Health Policy* 120:729–731
29. Hellsten E, Chu S, Crump RT, Yu K, Sutherland JM (2016) New pricing approaches for bundled payments: leveraging clinical standards and regional variations to target avoidable utilization. *Health Policy* 120:316–326
30. Ali AM, Gibbons CE (2017) Predictors of 30-day hospital readmission after hip fracture: a systematic review. *Injury* 48:243–252
31. Pitzul KB, Wodchis WP, Carter MW, Kreder HJ, Voth J, Jaglal SB (2016) Post-acute pathways among hip fracture patients: a system-level analysis. *BMC Health Serv Res* 16:275
32. Pitzul KB, Wodchis WP, Kreder HJ, Carter MW, Jaglal SB (2017) Discharge destination following hip fracture: comparative effectiveness and cost analyses. *Arch Osteoporos* 12:87
33. Fisher ES, Wennberg DE, Stukel TA, Gottlieb DJ, Lucas FL, Pinder ÉL (2003) The implications of regional variations in Medicare spending. Part 2: health outcomes and satisfaction with care. *Ann Intern Med* 138:288–298
34. Ireland AW, Kelly PJ, Cumming RG (2015) Total hospital stay for hip fracture: measuring the variations due to pre-fracture residence, rehabilitation, complications and comorbidities. *BMC Health Serv Res* 15:17
35. Jung HY, Trivedi AN, Grabowski DC, Mor V (2016) Does more therapy in skilled nursing facilities lead to better outcomes in patients with hip fracture? *Phys Ther* 96:81–89
36. Sheehan KJ, Sobolev B, Guy P, Bohm E, Hellsten E, Sutherland JM, Kuramoto L, Jaglal S, for the Canadian Collaborative Study on Hip Fractures (2016) Constructing an episode of care from acute hospitalization records for studying effects of timing of hip fracture surgery. *J Orthop Res* 34:197–204
37. Sobolev B, Guy P, Sheehan KJ et al (2016) Time trends in hospital stay after hip fracture in Canada, 2004–2012: database study. *Arch Osteoporos* 11:13
38. Sheehan KJ, Filliter C, Sobolev B et al (2018) Time to surgery after hip fracture across Canada by timing of admission. *Osteoporos Int* 29:653–663
39. Canadian institutes for Health Information (2009) CIHI data quality study of the 2007–2008 discharge abstract database. Canadian Institutes for Health Information. https://secure.cihi.ca/free_products/DAD_2007-08_Main_Report_en_web.pdf. Accessed 9 May 2018
40. Canadian institutes for Health Information. (2013) Canadian Institutes for Health Information (2013) Canadian Hospital Reporting Project Technical Notes - Clinical Indicators. http://publications.gc.ca/collections/collection_2013/icis-cihi/H118-86-1-2013-eng.pdf Accessed 10 Oct 2018
41. Nikkel LE, Fox EJ, Black KP, Davis C, Andersen L, Hollenbeak CS (2012) Impact of comorbidities on hospitalization costs following hip fracture. *J Bone Joint Surg Am* 94:9–17
42. Guy P, Sheehan KJ, Morin SN, Waddell J, Dunbar M, Harvey E, Sirett S, Sobolev B, Kuramoto L, Tang M, Canadian Collaborative Study on Hip Fractures (2017) Feasibility of using administrative data for identifying medical reasons to delay hip fracture surgery: a Canadian database study. *BMJ Open* 7:e017869
43. Wu H, Gao L, Zhang Z (2014) Analysis of crash data using quantile regression for counts. *J Transport Eng* 140:04013025
44. StataCorp (2017) Statistical Software: Release 15. StataCorp LP, College Station
45. Sobolev B, Guy P, Sheehan KJ, Bohm E, Beaupre L, Morin SN, Sutherland JM, Dunbar M, Griesdale D, Jaglal S, Kuramoto L, Canadian Collaborative Study on Hip Fractures (2017) Hospital mortality after hip fracture surgery in relation to length of stay by care delivery factors: a database study. *Medicine* 96:e6683
46. Sheehan KJ, Sobolev B, Guy P, Kuramoto L, Morin SN, Sutherland JM, Beaupre L, Griesdale D, Dunbar M, Bohm E, Harvey E (2016) In-hospital mortality after hip fracture by treatment setting. *CMAJ* 188:1219–1225
47. Ireland AW, Kelly PJ (2013) Total length of stay, costs and outcomes at final discharge for admitted patients with hip fracture: linked episode data for Australian veterans and war widows. *Int Med J* 43:1280–1286
48. National Hip Fracture Database (NHFD) (2018) Recovering after a hip fracture: helping people understand physiotherapy in the NHS. Physiotherapy 'Hip Sprint' audit report 2017. Royal College of Physicians, editor. London. <https://www.rcplondon.ac.uk/projects/outputs/recovering-after-hip-fracture-helping-people-understand-physiotherapy-nhs>. Accessed 23 Apr 2018
49. Beaupre LA, Cinats JG, Jones CA, Scharfenberger AV, William C, Johnston D, Senthilselvan A, Saunders LD (2007) Does functional recovery in elderly hip fracture patients differ between patients admitted from long-term care and the community? *J Gerontol A-Bio Sci Med Sci* 62:1127–1133

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.