

a statistically significant difference in any of the outcome measures. However, at latest follow-up rTSA patients with scapula notching had significantly worse outcome scores according to all 5 metrics as well as forward flexion and abduction (Table 1). Evaluation of the pre- to post-operative improvement also demonstrated statistically significant differences in clinical outcomes between the notching and no notching cohorts as depicted in Table 2.

**Discussion and Conclusion:** This study identifies a 16% notching rate after a minimum of five years follow up. Based on our analysis, inferior scapular notching is associated with a statistically significant degradation of clinical outcomes and range of motion compared to patients with an absence of scapular notching. Based on this, it is advisable to limit the risk of inferior scapular notching through patient selection, technique variation and implant choice.

**Reference**

1. Mollon B, Mahure SA, Roche CP, Zuckerman JD. Impact of scapular notching on clinical outcomes after reverse total shoulder arthroplasty: an analysis of 476 shoulders. *J Shoulder Elbow Surg* 2017;26:1253-61. <http://dx.doi.org/10.1016/j.jse.2016.11.043>

**Paper #14 CHARACTERISTICS OF PATIENTS WHO RATE THEIR SUBJECTIVE SATISFACTION AS UNCHANGED OR WORSE AFTER REVERSE TOTAL SHOULDER ARTHROPLASTY**

Ira M. Parsons, MD<sup>a</sup>, Howard D. Routman, DO<sup>b</sup>, Christopher Roche, MBA, MS<sup>c</sup>, Richard J. Friedman, MD, FRCSC<sup>d</sup>, <sup>a</sup>The Knee, Hip and Shoulder Center, Portsmouth, New Hampshire, USA; <sup>b</sup>Atlantis Orthopaedics, Palm Beach Gardens, Florida, USA; <sup>c</sup>Exactech, Gainesville, Florida, USA; <sup>d</sup>Medical University of South Carolina, Charleston, South Carolina, USA

**Introduction:** Patient reported outcome measures are an important determinant of surgical results as objective measures of function may not correlate with patient satisfaction. The prevalence of reverse total shoulder arthroplasty (rTSA) has increased over the past several years and for many surgeons now represents a majority of arthroplasty cases. It is important to understand who is at risk for unsatisfactory outcomes after rTSA so that surgeons can appropriately counsel patients as part of a shared decision-making process. The purpose of this study was to compare characteristics of patients who reported to be unchanged or worse to those who reported to be better or much better after rTSA.

**Methods:** Data was derived from a prospective registry of patients who underwent primary rTSA with a minimum 2-year follow-up. The same implant was used in all cases (Equinoxe, Exactech, Inc.). This registry collects information on demographics, diagnosis, comorbidities, implant information, preoperative pain and function and postoperative pain and function. Patients are also asked to rate their subjective satisfaction from which patients were divided into 2 groups: those who rated themselves as worse or unchanged (Unimproved Group, UG) and those who rated themselves as better or much

better (Improved Group, IG). Groups were compared for differences using a 2-tailed t-test with significant set at  $P < .05$ .

**Results:** There were 851 patients overall including 768 (90%) in IG and 83 (10%) in UG. There were no differences in age, gender, ethnicity or BMI between groups. Patients in IG were more likely to report no comorbidities (43% vs 25%,  $P = .001$ ), while those in UG were more likely to report a history of coronary artery disease (22% vs 10%,  $P = .03$ ) and diabetes (17% vs 10%,  $P = .04$ ). Patients in IG were more likely to have a diagnosis of osteoarthritis (56% vs 41%,  $P = .03$ ). Patients in UG were more likely to have rheumatoid arthritis (10% vs 3%,  $P = .03$ ). There were no differences between groups for diagnoses of rotator cuff tear or cuff tear arthropathy. Patients in UG were significantly more likely to have undergone prior surgery (42% vs 24%,  $P = .0001$ ). There were no differences in any implant configuration between groups. Patients in UG were much more likely to report residual moderate to severe pain (48% vs 6%,  $P < .001$ ) and 25% of patients in UG reported pain that was unchanged or worse after rTSA. Outcome scores were significantly worse for all measures for patients in the UG.

**Discussion:** This study demonstrates that up to 10% of patients rate themselves as unchanged or worse after surgery. These patients are more likely to have had prior surgery and more likely to have certain comorbidities. Patients with a diagnosis of osteoarthritis are more likely to rate themselves as better. While subjective function scores and abduction did not differ between groups preoperatively, those in UG did have significantly lower scores on all other preoperative measures. Interestingly, while outcomes were significantly worse for all measures in UG, there was improvement in all measures despite patients subjectively being worse or unchanged (Table 1).

**Paper #15 PRIMARY REVERSE SHOULDER ARTHROPLASTY USING CONTEMPORARY IMPLANTS IS ASSOCIATED WITH VERY LOW REOPERATION RATES**

Joaquin Sanchez-Sotelo, MD, PhD, Jason R. Kang, MD, Matthew J. Dubiel, MD, Robert H. Cofield, MD, Scott P. Steinmann, MD, Bassem T. Elhassan, MD, Mark E. Morrey, MD, John W. Sperling, MD, MBA, Mayo Clinic, Department of Orthopaedic Surgery, Rochester, Minnesota, USA

**Introduction:** The early results of reverse shoulder arthroplasty (RSA) were influenced to some extent by the use of first generation implants as well as surgeons' learning curves, resulting in relatively high reoperation rates. As experience has been gained with the procedure and new implants have been introduced, reasons for reoperation and complication rates may have changed. The purpose of this study was to quantify the burden and identify the indications for reoperation after primary RSA using contemporary implants and techniques.

**Methods:** A retrospective review of 1,649 primary RSAs implanted consecutively during a 7-year period (2009-2015) at a single institution was conducted. Study dates were chosen to minimize learning curve associated bias and to have a minimum follow-up of 2 years. All arthroplasties were performed by five fellowship trained shoulder

**Table 1**

	UG Preop	IG Preop	P	UG Postop	IG Postop	P
<b>Abduction</b>	66 + 31	70 + 33	NS	91 + 31	114 + 27	.0001
<b>Forward Elevation</b>	79 + 35	88 + 38	NS	109 + 36	141 + 24	.0001
<b>External Rotation</b>	11 + 20	16 + 22	.02	26 + 20	35 + 17	.0001
<b>Pain Score</b>	6.7 + 2.2	5.8 + 2.2	.003	4.1 + 3.0	0.9 + 1.7	.0001
<b>Function Score</b>	3.4 + 2.1	3.7 + 2.0	NS	5.5 + 2.8	8.2 + 1.7	.0001
<b>SST</b>	2.7 + 2.4	3.6 + 2.7	.003	6.1 + 3.8	10.3 + 2.2	.0001
<b>ASES</b>	31 + 15	37 + 15	.001	54 + 27	85 + 15	.0001

**Table 1** Demographics

Demographics	
Total number of primary RSA (2009-2015)	1,649
Number of RSAs requiring reoperation	39 (2.37%)
Average age at primary RSA	63.6 years old (range 20-87, SD 12.5)
Gender	18 male, 21 female
Laterality	16 right, 23 left
Average time to reoperation	20.6 months (range 0.4-71.2, SD 19.0)
Indications for primary RSA* (more than one indication for some patients)	
Cuff tear arthropathy	15
Proximal humerus fracture sequela	10
• Acute fracture	3
• Failed fixation	5
• Failed non-operative	2
Septic arthritis	4
Massive rotator cuff tear	3
Rheumatoid arthritis	3
Charcot arthropathy	2
AVN with rotator cuff tear	1
OA severe glenoid bone loss	1
Tumor	1

**Table 2** Indications for RSA reoperation

Indications for RSA reoperation	Number of Reoperations (n = 39)	Percent of primary RSAs (n = 1,649)
Instability	9 (23.1%)	0.55%
Infection	9 (23.1%)	0.55%
Glenoid component	6 (15.4%)	0.36%
• Baseplate loosening	5 (12.8%)	0.30%
• Glenosphere dissociation	1 (2.56%)	0.06%
Humeral component	17 (43.6%)	1.03%
• Humeral tray fracture	9 (23.1%)	0.55%
• Humeral stem aseptic loosening	4 (10.3%)	0.24%
• Periprosthetic humerus fracture	4 (10.3%)	0.24%
Deltoid failure	1 (2.56%)	0.06%

surgeons at a tertiary referral center. Demographics and indications for primary RSA were collected (Table 1), and all reoperations were identified, analyzed and categorized for trends associated with each type reoperation performed.

**Results:** 39 (2.37%) reoperations were performed for a variety of indications (Table 2). Overall, only a few patients with infection or instability required reoperation, (0.55%). The most common indications for reoperation were related to the humeral component (1.02%); the majority of humeral component complications were related to a specific design flaw of one implant system. RSAs performed for proximal humeral fracture sequela more commonly underwent reoperation due to instability or humeral component related issues; all four cases of aseptic humeral stem loosening occurred in the setting of proximal humeral fracture sequela treatment. Only 0.36% of all primary RSAs required reoperation due to glenoid complications.

**Conclusion:** Primary RSA performed with contemporary implants and surgical techniques seems to be associated with a very low rate of reoperation. The most common reasons for reoperation were humeral component fracture for one particular implant, humeral loosening, dislocation, infection, and glenoid failure, all under 1% each.

#### Paper #16 INCIDENCE OF POST-OPERATIVE ACROMIAL FRACTURES WITH ONLAY VS INLAY REVERSE SHOULDER ARTHROPLASTY

Ryan LeDuc, BS, Dane H. Salazar, MD, Nickolas G. Garbis, MD, Loyola University Medical Center, Maywood, Illinois, USA

**Background:** Acromial fractures in reverse shoulder arthroplasty (RSA) have been reported at rates up to 7%. Although these patients are reported to have improved function compared to their preoperative state, acromial insufficiency may negatively impact the final postoperative outcome in RSA compared to a cohort of patients without postoperative acromial complication. There are multiple implant systems used for reverse total shoulder arthroplasty. The purpose of this study was to retrospectively compare the rates of acromial fractures in two of such systems, using the outcomes of two surgeons at a single institution.

**Methods:** Patients that underwent surgery from August 2013-July 2016 were retrospectively reviewed. Postoperative clinic notes were reviewed to determine the incidence of acromial fracture after surgery, and all imaging studies obtained were reviewed to confirm presence or absence of acromial fracture. Patients with at least six months of follow up were included in our study population, as this has been identified as an early postoperative complication. Preoperative and one-year ASES scores were calculated and compared for subjects who had sufficient follow up.

**Results:** Our study population included 109 total patients, 58 who have an onlay reverse shoulder and 51 with an inlay system in place. There were 9 total acromial fractures recorded (8.26%). The onlay cohort had a 12.07% acromial fracture rate (7 out of 58), compared to a 3.92% rate for the inlay group (2 out of 51). Compared to patients receiving an inlay system, those receiving an onlay prosthesis were 3.33 times more likely to experience a complication, though this did not reach statistical significance (95% CI: 0.60-34.38;  $P = .23$ ).

**Conclusion:** There may be a trend towards higher rate of post-operative acromial fractures when an onlay reverse shoulder is used as opposed to an inlay system. This may be due to the potentially increased tension placed on the deltoid resulting from the increased proximal size of typical onlay prostheses. We encourage further investigation into the acromial complication rates between different systems (Table 1).