



## Ultrastructure of the anterior adhesive apparatus of the gill parasite *Macrogyrodactylus clarii* and skin parasite *M. congolensis* (Monogenea; Gyrodactylidae) from the catfish *Clarias gariepinus*

Mohammed Mohammed El-Naggar<sup>a,b</sup>, Safaa Zaky Arafa<sup>c,\*</sup>, Samir Ahmed El-Abbassy<sup>a</sup>,  
Graham C. Kearn<sup>d</sup>, Jo Cable<sup>b</sup>

<sup>a</sup> Zoology Department, Faculty of Science, Mansoura University, Mansoura, Egypt

<sup>b</sup> School of Biosciences, Cardiff University, CF10 3AX, UK

<sup>c</sup> Department of Basic Sciences, Biology Section, Deanship of Preparatory Year and Supporting Studies, Imam Abdulrahman Bin Faisal University, Saudi Arabia

<sup>d</sup> School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich NR4 7TJ, UK

### ARTICLE INFO

#### Keywords:

Platyhelminthes  
Monogenea  
Fish ectoparasite  
Temporary adhesion  
Adhesive apparatus  
Ultrastructure

### ABSTRACT

Transmission electron microscopy (TEM) was used for the first time to study the anterior adhesive apparatus of the monogeneans *Macrogyrodactylus clarii* Gussev, 1961 and *M. congolensis* (Prudhoe, 1957) Yamaguti, 1963 inhabiting gills and skin respectively of the same catfish *Clarias gariepinus*. Despite the different microhabitats occupied by these parasites, the present study revealed that they have a similar anterior adhesive system. In both parasites, the anterior adhesive apparatus consists of three types of gland cells: G1 cells that produce rod-shaped bodies (S1), G2 cells manufacture irregularly shaped bodies (S2) and G3 cells form mucoid-like secretions (S3). In the cytoplasm of G1 cells, a single layer of microtubules encloses each developing rod-shaped body. A unique feature of S1 secretory bodies is that some fully developed S1 bodies are attached to each other, forming large condensed globules in the cytoplasm of G1 gland cells and terminal portion of the G1 ducts, but none were detected in the adhesive sacs outside the ducts. In the adhesive sacs, G1 ducts open with multiple apertures whereas each of the G2 and G3 ducts have a single opening. The adhesive sacs are lined with two types of tegument (st1 and st2). A third tegument type (st3) connects the st2 tegument with the general body tegument. Only st1 has microvilli. Each adhesive sac is provided with a spike-like sensillum and single unciliated sense organ. The possible functions of microvilli in increasing the surface area and assistance in spreading and mixing of the adhesive secretion, and the role of sense organs associated with the adhesive sacs are discussed.

### 1. Introduction

Monogenean ectoparasites attach to their hosts primarily with their posterior attachment organ (haptor), which is equipped with hamuli and marginal hooklets [1], but in order to move from one position to another they rely on their anterior adhesive apparatus [2]. Typically, monogeneans move on the host or artificial substrates by stretching out their bodies and attaching with head lobes to the host tissue, releasing and moving the haptor to attach close to the adhesive areas of the head lobes, and then they detach the head lobes to move anteriorly where they attach again to a new site. Some can move in a similar leech-like manner upside down, using the water surface tension [3].

The anterior adhesive apparatus has been studied with transmission electron microscopy (TEM) and/or scanning electron microscopy (SEM)

in many monogenean parasites, including the gyrodactylids [4–6], dactylogyrids [7,8], entobdellids [2,9], acanthocotyliids [10], monocoelyids [11,12] and ancyrocephalids [13]. They have various kinds of gland cells that open either into the outer syncytial tegumental layer [7,14], or onto the specialized haptor [13,15–17] or ventrally-located head regions (see for example, El-Naggar and Khidr [8], Wong et al. [13]). The monogenean anterior adhesive apparatus produces one to three types of secretion bodies. Species with rod-shaped bodies (S1), spherical bodies (S2) and irregularly-shaped, electron-lucent vesicles (S3) include *Gyrodactylus eucaliae* (see Kritsky, 1978) [4], *G. sprostonae* (see Yuan and Long [18]), *Dactylogyryus amphibothrium* and *D. hemi-amphibothrium* (see El-Naggar and Kearn [7]), *D. aristichthys* (see Yuan and Long [19]), *Cichlidogyryus halli* (see El-Naggar and Khidr [8]) and *Merizocotyle icopae* (see Cribb et al. [20]). Two types of secretion, rod-

\* Corresponding author.

E-mail address: [sarafah@iau.edu.sa](mailto:sarafah@iau.edu.sa) (S.Z. Arafa).

<https://doi.org/10.1016/j.parint.2019.03.005>

Received 29 September 2018; Received in revised form 26 February 2019; Accepted 7 March 2019

Available online 08 March 2019

1383-5769/ © 2019 Elsevier B.V. All rights reserved.






Fig. 2. G1 gland cell of *Macrogryodactylus clarii* containing S1 rod-shaped bodies and large spherical globules (lg). dS1, Developing rod-shaped bodies; ch, chromatin; GER, granular endoplasmic reticulum; N, nucleus; Nu, nucleolus; r, ribosomes; S1, rod-shaped secretory bodies.

anterior unbranched region of the intestine. The G2 cells are larger than the G1 cells and produce irregularly-shaped secretory bodies (S2). In both *M. clarii* and *M. congolensis*, the G3 cells are three in number located lateral to the cerebral region and produce translucent mucoid secretory bodies (S3) (Fig. 1).

### 3.1. Gland cells

TEM of both *M. clarii* and *M. congolensis* revealed that each G1 gland cell has a nearly spherical nucleus with granular nucleoplasm, conspicuous nucleolus and condensed chromatin (Fig. 2). The cytoplasm is moderately electron-dense and contains abundant granular endoplasmic reticulum (GER), numerous ribosomes (Figs. 2–5), a few Golgi bodies, small electron-lucent vesicles and mitochondria. The dilated cisternae of the GER enclose an amorphous, finely granular material with an electron density slightly higher than that of the basal cytoplasm (Fig. 2). Generally, each fully-developed, rod-shaped S1 body is of high electron density and measures 0.3–0.5 (average 0.4)  $\mu\text{m}$  in diameter. The maximum length measured in sections is 4–6.5




Fig. 3. Cytoplasm of the G1 gland cell of *Macrogryodactylus clarii* containing fully formed S1 rod-shaped bodies and granular endoplasmic reticulum (GER).




Fig. 4. G1 gland cell of *Macrogryodactylus congolensis* containing fully formed rod-shaped bodies (S1), large globule (lg) and developing S1 (dS1) secretory bodies. Note that the developing rod-shaped bodies (dS1) have different sizes and are surrounded by microtubules (mt). Note also that some of the developing S1 bodies contain a peripheral layer of small electron-dense granules (arrows) and have no bounding membrane. GER, granular endoplasmic reticulum.




Fig. 5. G1 gland cell of *Macrogryodactylus congolensis* containing fully formed rod-shaped bodies (S1) and large globules (lg) each containing many S1 bodies (arrow). Note the presence of small translucent vesicles (v) and developing rod-shaped bodies (dS1).

(average 6)  $\mu\text{m}$ . These bodies are membrane bounded and contain a finely granular dense matrix in which small particles are embedded within higher electron-dense material (Figs. 2–5). In sections, immature S1 bodies have a greater diameter than that of the fully developed ones (Figs. 4, 6). They measure 0.4–0.7 (average 0.6)  $\mu\text{m}$  in *M. congolensis* and *M. clarii* and contain granular material, with electron-density lower than that of the smaller S1 bodies (Figs. 4, 6). Each of the large, immature, less electron-dense bodies and some of the small highly electron-dense bodies are enclosed by a single layer of microtubules, which appear to be parallel with each other and with the long axis of the rod (Figs. 4–6). Some of the less electron-dense S1 bodies contain a peripheral layer of small electron-dense granules and have no bounding membrane (Fig. 4). In cross sections, some fully developed S1 bodies attach to each other, forming large condensed globules with various sizes and shapes (Figs. 2, 4, 5). Their number varies from 3 to 7 S1 bodies in each globule. In a few sections of *M. congolensis*, some fully formed S1 secretory bodies with peripheral translucent vesicles were detected (Fig. 7).

Each G2 cell is enclosed by a layer of fibrous interstitial material. They have an irregularly shaped nucleus with a relatively large, conspicuous nucleolus, granular nucleoplasm, small chromatin patches and nuclear membrane with characteristic nuclear pores (Fig. 8). The cytoplasm is moderately electron-dense, but it is slightly darker than that of the G1 cells. It contains abundant GER, free ribosomes, and




Fig. 6. G1 gland cell of *Macrogyrodactylus congolensis* showing longitudinal sections of the large developing rod-shaped bodies (dS1) and fully formed rod-shaped bodies (S1). Note the microtubules (mt) associated with dS1.




Fig. 7. Magnified S1 secretory bodies of *Macrogyrodactylus congolensis* with translucent vesicles (arrow heads).

numerous Golgi complexes, which in many sections are aggregated (in groups of 2–4) in close proximity to the nuclear membrane (Fig. 9). Each Golgi complex consists of 3–5 narrow parallel cisternae terminating with small and large vesicles (Fig. 9). Both cisternae and vesicles are filled with homogeneous, highly electron-dense material. In sections, the irregularly shaped bodies (S2) have different sizes ranging from 0.7 to 1.5 (average 1.2)  $\mu\text{m}$  in diameter. They are abundant and contain granular, highly electron-dense material (Figs. 8, 9). However, in *M. clarii* with higher magnification, each S2 body contains tubular structures with lower electron-density, which are embedded in highly electron-dense ground substance (Fig. 10). In most regions of the G2 cells, fully developed S2 bodies are surrounded by cytoplasm characterized by translucent ground substance (Figs. 8, 10).

Each G3 gland cell has a nearly oval nucleus with granular nucleoplasm, conspicuous nucleolus and condensed chromatin patches (Fig. 11). Some GER have dilated cisternae. The mucoid secretory bodies (S3) are abundant, irregularly shaped (1–1.9, average 1.4,  $\mu\text{m}$ ) and contain granular moderately electron-dense material (Fig. 11).




Fig. 8. G2 gland cell of *Macrogyrodactylus clarii* surrounded by fibrous interstitial material (fm) and containing large nucleus (N) with conspicuous nucleolus (Nu) and irregularly shaped secretory bodies (S2) surrounded by translucent area (\*).




Fig. 9. Magnified part of G2 gland cell of *Macrogyrodactylus clarii* with Golgi bodies (Go), granular endoplasmic reticulum (GER), ribosomes (r) and irregularly shaped secretory bodies (S2).

### 3.2. Gland ducts and adhesive areas

Ducts of the G1, G2 and G3 gland cells carrying the secretory bodies S1, S2 and S3, respectively, extend anteriorly as cytoplasmic processes where they converge on adhesive papillae through which they open into the adhesive sacs (Figs. 1, 12–21). As the gland ducts approach the adhesive sac, they dilate and become closely packed (Figs. 12,19). At this point, some of the gland ducts are associated with muscle fibres that are present beneath the tegument lining the adhesive sac (Fig. 13). Most ducts of the G1 cells are filled with completely formed rod-shaped bodies (Figs. 12,13,14), but in some sections, a few condensed globules




Fig. 10. Magnified S2 of *Macrogyrodactylus clarii* containing tubular structures with lower electron density and surrounded by a translucent area (\*).




Fig. 11. G3 gland cell of *Macrogyrodactylus clarii* containing nucleus (N) with chromatin (ch), dilated cisternae of granular endoplasmic reticulum (dGER) and translucent mucoid secretory bodies (S3).




Fig. 12. Section through adhesive sac (as) of *Macrogyrodactylus clarii* showing the ventral surface of adhesive papillae (ap) covered with st1 tegument. S1, rod-shaped bodies.

of attached rods are found beside S1 bodies (Fig. 15). There are no microtubules in any of the gland ducts. Each G1 gland duct opens to the exterior via multiple apertures (Figs. 14–16). At the openings of the G1 ducts, five layers, three electron-dense and two electron-lucent (Fig. 16), bound each aperture. The outer layer membrane connects with the surrounding tegument by means of septate desmosomes (Fig. 14). Each one of the multiple apertures allows passage of a single rod (Fig. 14). Although large globules were detected in the terminal portion of the G1 ducts just beneath the multiple apertures, none of them were seen passing through the openings or outside the body




Fig. 13. Section through adhesive sac (as) of *Macrogyrodactylus clarii* showing muscle fibres (mf) in between G1 ducts. S1, rod-shaped bodies.




Fig. 14. Adhesive papilla of *Macrogyrodactylus clarii* showing S1 body protruding from its aperture. Note the st1 tegument covering the ventral surface of the adhesive papillae and st2 tegument covering the lateral surface of adhesive papillae. as, Adhesive sac; mi, microvilli; \*, homogeneous material around S1 bodies.




Fig. 15. Duct of G1 gland cell of *Macrogyrodactylus congolensis* containing rod-shaped bodies (S1) and large globules (lg) close to the multiple apertures. Note that the membrane bounding the outer layer of the multiple apertures is connected to the adjacent tegument (st1) by means of septate desmosomes (d) and presence of homogeneous material (\*) around S1 bodies. mi, Microvilli; st1, tegument covering the ventral surface of adhesive papillae.

(Fig. 15). Each of the G2 and G3 gland ducts opens to the exterior by a single aperture (Figs. 17–20).

The adhesive sac is lined with three types of tegumental layer (st1,




Fig. 16. Cross section of the multiple apertures of a G1 gland duct of *Macrogyrodactylus congolensis* showing that each aperture is bounded by five layers (l), three electron-dense and two electron-lucent. S1, rod-shaped bodies.




Fig. 17. Terminal portion of G2 gland duct of *Macrogyrodactylus congolensis* carrying S2 secretory bodies. f, fibrous layer; mi, microvilli; st1, microvillous tegument.




Fig. 18. Section through adhesive sac of *Macrogyrodactylus clarii* showing terminal portions of gland ducts carrying S1, S2 and S3 bodies. Note that secretory bodies of S3 bodies form particulate material.

st2 and st3) (Figs. 14, 15, 17, 22, 24). The first (st1) represents the outer tegumental layer covering the ventral surface of the adhesive papillae surrounding the gland duct openings (Figs. 12, 14, 15), while st2 represents the outer tegumental layer covering the lateral surfaces of the adhesive papillae (Figs. 12, 14, 22). The third type (st3) is the outer tegumental layer of the inner rim of the adhesive sac and connects st2




Fig. 19. Terminal portions of G1, G2 and G3 gland cells of *Macrogyrodactylus clarii* containing S1, S2 and S3 secretory bodies, respectively. Note that the component of S2 bodies (\*) diffuse into the lumen of the duct. mf, muscle fibres.




Fig. 20. Duct of G2 gland cell carrying S2 secretory bodies of *Macrogyrodactylus clarii*. c, Cilium; nb, nerve bulb; nt, neurotubule; St1, tegument covering the ventral surface of adhesive papillae.

and the tegumental layer of the general body surface (i.e. the outer surface of the head lobe) (Figs. 12, 22). Comparing the three tegumental layers, the st1 layer is relatively thin, electron-dense and has numerous microvilli but lacks secretory bodies (Figs. 14, 15). The st2 layer is highly electron-dense and contains abundant electron-dense bodies (Figs. 14, 22). No cytoplasmic organelles like mitochondria, Golgi bodies, GER or free ribosomes were found in st1 or st2 tegument. The st3 tegumental layer connects with the st2 tegument by means of junctional complexes (Fig. 22) and contains a few translucent vesicles containing moderately electron-dense particles (Fig. 24). These vesicles are restricted to the outer region of the tegument. Some electron-dense granular bodies, abundant rod-shaped, electron-dense bodies and a few mitochondria were also seen (Fig. 22 inset). The general body tegument contains abundant translucent vesicles and some electron-dense granular secretory bodies, but no rod-shaped bodies (Fig. 22).

No experimental work was performed to study the mechanism of attachment and detachment of the head lobes of *Macrogyrodactylus* species. However, in most sections the terminal portions of G1 ducts,




Fig. 21. The terminal portions of G2 gland ducts of *Macrogyrodactylus clarii* showing that the components of S2 bodies (\*) diffuse into the duct lumen.




Fig. 22. Intervening tegument (st3) connecting the sac tegument of type st2 with the general body tegument (gt) of *Macrogyrodactylus congolensis*. Note that st3 tegument and st2 are connected by a junctional complex (j). gb, Electron-dense granular secretory bodies; rb, rod-shaped electron-dense bodies; v, translucent vesicle with electron-dense granule. Inset: magnified st3 with abundant electron-dense, rod-shaped bodies (rb), few electron-lucent vesicles with dark granules (v) and electron-dense granular bodies (gb), similar to those in the general body tegument.

homogeneous particulate material was detected around the S1 bodies (Figs. 14, 15) while sections of the terminal portions of G2 and G3 ducts revealed considerable change in appearance of the secretory bodies particularly S2 and S3. The S3 bodies lose their membranes and their secretory components form homogeneous particulate material (Figs. 18,19), while S2 bodies become slightly smaller in size and their particulate components diffuse into the lumen of the duct in-between bodies that are still membrane-bounded (Figs. 19,21). Moreover, in the same region, these sections show a network of homogeneous material covering the surface of the adhesive papilla (Figs. 19, 21).

TEM revealed the presence of a single sensillum on each adhesive papilla (Fig. 20), in the intervening region between the adhesive sac and general body tegument (Fig. 23) and on the anterior region, which is covered by general body tegument (Fig. 24). Each sensillum has an elongated nerve bulb, which terminates in a single opening through which a single cilium protrudes (Figs. 20, 23, 24). Close to the opening, there is an electron-dense thickening and the lining of the opening is connected with the intervening tegument st3 via desmosomes (Figs. 23, 24). The nerve bulb contains neurotubules, electron-dense bodies and




Fig. 23. Section through the terminal part of ciliary sensillum of *Macrogyrodactylus clarii*. c, Cilium; edb, electron-dense bodies; et, electron-dense thickening; m, mitochondria; nb, nerve bulb; nt, neurotubules.



Fig. 24. Section through ciliary sensillum of *Macrogyrodactylus clarii*. c, cilium; d, desmosomes; et, electron-dense thickening; gt, general body tegument; m, mitochondria; r, root of the cilium; st3, intervening tegument.

mitochondria (Figs. 23, 24).

#### 4. Discussion

This is the first ultrastructural study of the anterior adhesive apparatus of the monogeneans *Macrogyrodactylus clarii* [22] from the gills of *Clarias gariepinus* and *M. congolensis* [23,24] from the skin and fins of the same host. TEM revealed that the anterior adhesive apparatus of both parasites consists of three types of gland cells (G1, G2 and G3). The G1 cells produce rod-shaped bodies (S1) and roughly spherical large globules, G2 cells secrete irregularly shaped, highly electron-dense bodies with tubular contents (S2) and G3 cells manufacture irregularly

shaped, mucoid-like secretion (S3). These glands resemble those of the anterior adhesive apparatus of other monogenean parasites [28–30]. Previous studies illustrated that congeners in the same microhabitat tend to have similar types of anterior adhesive secretions [7,29,31]. In the present study, the anterior adhesive apparatus of *M. clarii* and *M. congolensis* have the similar morphological features, despite the differences in their microhabitat, with the exception of the number of G2 cells: 10 pairs in *M. congolensis* and 7 pairs in *M. clarii*. Morphological similarities, however, do not exclude the possibility of chemical and/or functional differences [30].

In *D. amphibothrium*, El-Naggar and Kearn [7] found that S1 bodies in the G1 ducts connect with each other by membrane-like structures and a similar feature of interlinking band-like structures was observed between S1 bodies and S2 bodies in *Bravohollisia gussevi* and *Caballeria liewi* (see Wong et al. [13,16], respectively). Also, the bounding membranes of S1 bodies in *Entobdella australis* and *Entobdella* spp. (see Whittington and Cribb [29]) showed periodic dense bandings. None of these structures, however, were observed in either *M. congolensis* or *M. clarii*. A unique feature of these parasites though is the presence of large globular bodies in the cytoplasm and ducts of the G1 cells, in addition to fully formed S1 bodies. With TEM, it became evident that these globules are aggregations of S1 bodies. There was no evidence that S2 or S3 bodies in *M. clarii* and *M. congolensis* aggregate and coalesce in the cytoplasm of their cells but they become closely packed as they reach the terminal portions of the ducts. Another important feature of *M. clarii* and *M. congolensis* is that S2 bodies contain tubular structures, a feature not reported in any other monogeneans studied by TEM. In addition, the present study indicates that the fully formed rods in *M. clarii* and *M. congolensis* are considerably larger than the S1 bodies in *Entobdella* spp. (see Whittington and Cribb [29]).

During the early stage of assembly, the large, less electron-dense rods, and some of the smaller highly electron-dense rods of *M. clarii* and *M. congolensis* are enclosed by microtubules. The microtubules disappear when the rods are fully formed and become bounded by membrane. Microtubules have been reported in most other monogeneans studied (see for example Wong et al. [13]) except for monocotylids [11,12] and *Benedenia* spp. [32]. Moreover, the rods of *Monocotyle spiremae* have no bounding membrane and possess an outer electron-dense cortex and a more electron-lucent core [11]. El-Naggar and Kearn [7] suggested that encircling microtubules may play a role in transporting products from different parts of the cell prior to assembly of the secretory bodies. In addition, the microtubules may orientate the rods during their passage from within the gland cells to the lumen of their gland ducts, and help to maintain the parallel arrangement of rods into bundles [7].

The monogeneans *M. clarii* and *M. congolensis* resemble other gyrodactylids in that the secretions of the anterior adhesive apparatus open into a single pair of adhesive sacs, one situated antero-ventrally on each of the two head lobes [4]. Other monogeneans, with the exception of gyrodactylids and some monocotylids, have three distinct zones on each side of the head for the release of secretions (see, for example, El-Naggar and Kearn [7]). Such organization into six separate points of contact presumably allows the parasite to be more resistant to detachment caused by water currents [10].

In the present study of *Macrogryodactylus* spp., it has been established that the rod-shaped bodies and roughly spherical large globules produced by G1 gland cells are transported through ducts terminating with multiple apertures. Each aperture apparently permits the passage of only one rod but there is no evidence that the large globules pass through multiple apertures. Multiple apertures were reported in *G. eucaliae*, *Entobdella soleae* and *M. spiremae* (see Kritsky [4]). [2,11], respectively. However, in *D. amphibothrium* and *D. hemiamphibothrium* the rod-shaped bodies are released from ducts with single apertures (see El-Naggar and Kearn [7]). The unique feature of *M. clarii* and *M. congolensis* is that the ducts that carry the rod-shaped bodies also carry larger globules of the same secretion, but the globules were not seen

passing through duct apertures. It is possible that the multilayered boundary of one of the small multiple openings dilate to permit passage of the larger globules. Alternatively, the large spherical globules may liquefy or fragment before passing through the multiple apertures. Presence of material similar to the contents of S1 bodies and large globules in the terminal portion of G1 ducts (Figs. 10, 11) supports the latter suggestion. There is also some evidence that the large globules in the G1 cells are composite structures, perhaps made by accumulation of rod-shaped bodies or components of them. If correct, then the globules might escape from the duct openings after disintegration into their small rod-like components. Moreover, in *M. clarii* and *M. congolensis*, the S2 and S3 bodies showed considerable change in their appearance inside the terminal portions of the ducts indicating that they are released from duct openings in a liquid form. A similar feature was reported by Kearn and Evans-Gowing [9] who found that the spheroidal secretory bodies associated with the anterior adhesive apparatus of *E. soleae* transform within the duct terminations immediately prior to attachment of the head region.

A characteristic feature of *M. clarii* and *M. congolensis* is that each adhesive sac is lined with three types of tegument (st1, st2, and st3) that are different from the general body surface. The first kind (st1) is thickly covered with microvilli, a feature that has been reported in the adhesive areas of many monogeneans [2,4,5,7,9,11,33,34]. These specialized microvilli may be important during attachment of the head lobes by increasing the surface area available for binding the adhesive secretions to the head region. Lyons [34] suggested that these microvilli in *Gyrodactylus* spp. may assist in spreading the adhesive secretion of the head glands over the skin of the host into a thin “tacky” film. The microvilli may help to mix the products of different gland cells, which might have to interact with each other or with water before the sticky properties are developed.

Rod-shaped bodies are the most abundant component of the anterior adhesive apparatus of *M. clarii* and *M. congolensis*. These bodies also represent the main component of the anterior adhesive secretions of many monogeneans, which produce two or three types of secretory bodies such as *D. amphibothrium* and *D. hemiamphibothrium* (see El-Naggar and Kearn [7]) and *E. soleae* (see Kearn and Evans-Gowing [9]). Furthermore, rod-shaped bodies are the only secretory body recorded in the anterior adhesive apparatus of the monocotylid, *Monocotyle spiremae* (see Cribb et al. 11)).

The mechanism of attachment of *M. clarii* and *M. congolensis* may involve adhesion of the adhesive sac rim to the host tissues, protrusion of the adhesive papillae by means of associated muscles and release of secretory bodies through gland duct openings. The spike sensillum and other ciliary structures may serve as chemoreceptors that control attachment of the adhesive sacs. In monogeneans, it was suggested that stickiness could be a property of one type of secretory body or could develop by mixing between two types of secretion [2,7,20,35]. Interaction between secretory bodies and water [2] or between secretory bodies and host mucus [11] are possible alternative mechanisms.

Detachment of the head lobes of *M. clarii* and *M. congolensis* may occur mechanically by contraction of muscle fibres attached to the anterior region [11]. In *E. soleae*, tegument of the adhesive pads may play a part in detachment of the head region, by release of secretory bodies, which are abundant in this layer, or by some other physical or chemical change mediated via the tegumentary membrane [9]. In *M. spiremae*, where only one type of secretion (rods) was found, detachment may involve additional glue, physical detachment by muscle contraction or extrusion of material surrounding the rods [11]. Experimental studies are still needed in this field to determine which secretion is responsible for attachment and how detachment takes place: a potentially lucrative area for industry in relation to binding agents in water.

Regarding parasite-host specificity, it has been reported that the epidermal mucous cells of specific fish hosts may influence parasite attachment (see review in Whittington et al. [31]). The anterior

attachment region of *Gyrodactylus derjavini* contains mannose-rich glycoproteins, which are implicated in stimulating the alternative complement pathway in the host [36]. Specific differences in host fish epithelium and differences in monogenean anterior adhesive chemistry or in the chemistry of the specialized tegument of the anterior adhesive area may all contribute to host specificity amongst monogeneans [30].

## Acknowledgements

The study was funded by a Royal Society Developing World Visit Research grant to MME and a Natural Environment Research Council (NERC), Advanced Research Fellowship (NER/J/S/2002/00706) to JC

## References

- G.C. Kearn, Parasitism and the Platyhelminths, Chapman and Hall, London, 1998, p. 544.
- M.M. El-Naggar, G.C. Kearn, Glands associated with the anterior adhesive areas and body margins in the skin-parasitic monogenean *Entobdella soleae*, Int. J. Parasitol. 13 (1983) 67–81, [https://doi.org/10.1016/S0020-7519\(83\)80067-8](https://doi.org/10.1016/S0020-7519(83)80067-8).
- M.M. El-Naggar, A.M. El-Naggar, S.A. El-Abbassy, Microhabitat and movement of the viviparous monogeneans *Gyrodactylus alberti*, *Macrogyrodactylus clarii* and *M. congolensis* from the Nile catfish *Clarias gariepinus*, J. Egypt Ger. Soc. Zool. 35 (2001) 169–187.
- D.C. Kritsky, The cephalic glands and associated structures in *Gyrodactylus eucaliae* Ikezaki and Hoffman, 1957 (Monogenea: Gyrodactylidae), Proc. Helminthol. Soc. Wash. 45 (1978) 37–49.
- M.M. El-Naggar, Scanning electron microscope studies on the head lobes and haptor of the monogenean *Gyrodactylus groschafti* Ergens, 1973, J. Egypt Ger. Soc. Zool. 8 (1992) 435–445.
- M.M. El-Naggar, Scanning electron microscope observations on the head lobes and haptor of the monogenean *Macrogyrodactylus clarii* Gussev, 1961, J. Egypt Ger. Soc. Zool. 10 (1993) 143–155.
- M.M. El-Naggar, G.C. Kearn, Ultrastructural observations on the anterior adhesive apparatus in the monogeneans *Dactylogyrus amphibothrium* Wagner, 1957 and *D. hemiamphibothrium* Ergens, 1956, Z. Parasitenkd. 61 (1980) 223–241.
- M.M. El-Naggar, A.A. Khidr, Ultrastructural observations on the anterior adhesive apparatus of the monogenean gill parasite *Cichlidogyrus halli typicus*, J. Egypt Ger. Soc. Zool. 26 (1998) 309–324.
- G.C. Kearn, R. Evans-Gowing, Attachment and detachment of the anterior adhesive pads of the monogenean (Platyhelminth) parasite *Entobdella soleae* from the skin of the common sole (*Solea solea*), Int. J. Parasitol. 28 (1998) 1595–1607, [https://doi.org/10.1016/S0020-7519\(98\)00059-9](https://doi.org/10.1016/S0020-7519(98)00059-9).
- J.A. Rees, G.C. Kearn, The anterior adhesive apparatus and an associated compound sense organ in the skin-parasitic monogenean *Acanthocotyle lobianchi*, Z. Parasitenkd. 70 (1984) 609–625, <https://doi.org/10.1007/BF00926591>.
- B.W. Cribb, I.D. Whittington, L.A. Chisholm, Observations on the ultrastructure of the anterior glands in the monogenean, *Monocotyle spiremae* (Monocotylidae), from the gills of *Himantura fai* (Dasyatididae), Int. J. Parasitol. 27 (1997) 907–917, [https://doi.org/10.1016/S0020-7519\(97\)00061-1](https://doi.org/10.1016/S0020-7519(97)00061-1).
- B.W. Cribb, I.D. Whittington, L.A. Chisholm, Observations on the ultrastructure of the anterior adhesive areas and other anterior gland cells in the monogenean *Merizocotyle australensis* (Monocotylidae) from the nasal fossae of *Himantura fai* (Dasyatididae), Microsc. Res. Tech. 42 (1998) 200–211, [https://doi.org/10.1002/\(SICI\)1097-0029\(19980801\)42:3%3C200::AID-JEMT4%3E3.0.CO;2-W](https://doi.org/10.1002/(SICI)1097-0029(19980801)42:3%3C200::AID-JEMT4%3E3.0.CO;2-W).
- W.L. Wong, G.P. Brennan, D.W. Halton, A.G. Maule, L.H. Lim, Ultrastructure of head organs (anterior adhesive apparatus) and posterior secretory systems of *Caballeria lievi* Lim, 1995 (Monogenea, Ancyrocephalidae), Parasitol. Res. 113 (2014) 3935–3946, <https://doi.org/10.1007/s00436-014-4057-8>.
- M.M. El-Naggar, A.A. Khidr, G.C. Kearn, Ultrastructural observations on the tegument and associated structures of the monogenean *Cichlidogyrus halli typicus* (Price & Kirk, 1967) Paperna, 1979, Int. J. Parasitol. 21 (1991) 707–713.
- M.M. El-Naggar, G.C. Kearn, Haptor glands in the gill-parasitic, ancyrocephaline monogenean *Cichlidogyrus halli typicus* and the report of a possible prokaryotic symbiont, Int. J. Parasitol. 19 (1989) 401–408, [https://doi.org/10.1016/0020-7519\(89\)90096-9](https://doi.org/10.1016/0020-7519(89)90096-9).
- W.L. Wong, G.P. Brennan, D.W. Halton, L.H.S. Lim, Fine structure of the anterior adhesive apparatus (head organs) of *Bravohollisia gussevi* Lim, 1995 (Monogenea: Ancyrocephalidae), Parasitology 132 (2006) 427–438, <https://doi.org/10.1017/S0031182005009054>.
- L.H.S. Lim, D. Gibson, Species of *Triacanthinella* Bychowsky and Nagibina, 1968 (Monogenea: Ancyrocephalidae) from Triacanthid Teleosts off Peninsular Malaysia, with a generic revision, amended diagnosis and key, Syst. Parasitol. 70 (2008) 191–213, <https://doi.org/10.1007/s11230-008-9137-7>.
- W.J. Yuan, S. Long, Ultrastructure of the adhesive apparatus in the monogenean *Gyrodactylus sprostanae* (In Chinese), Acta Zool. Sin. 43 (1997) 209–210.
- W.J. Yuan, S. Long, Ultrastructure of the adhesive apparatus in some monogeneans (In Chinese), Acta Zool. Sin. 42 (1996) 3–9.
- B.W. Cribb, W.D. Armstrong, I.D. Whittington, Mechanism of adhesion and detachment at the anterior end of *Merizocotyle icopae* (Monogenea: Monocotylidae) including ultrastructure of the anterior adhesive matrix, Parasitology 129 (2004) 181–190, <https://doi.org/10.1017/S0031182004005530>.
- A.A. Khidr, S.H. Hassan, G.C. Kearn, Redescription of *Enterogyrus cichlidarum* Paperna 1963 (Monogenea: Ancyrocephalinae) from the stomach of *Tilapia* spp. in Egypt, Proc. Zool. Soc. Arab Repub. of Egypt 18 (1990) 269–280.
- A.V. Gussev, A Viviparous Monogenetic Trematode from Freshwater Basins of Africa, vol. 136, Doklady Akademii Nauk SSSR, 1961, pp. 490–493.
- S. Prudhoe, Trematoda. Exploration du Parc National de l'Upemba. Mission F Witte, 48 (1957), pp. 1–28.
- S. Yamaguti, Systema Helminthum, Monogenea and Aspidocotylea, Vol. IV Interscience, New York, 1963, p. 699.
- M.M. El-Naggar, H.M. Serag, Redescription of *Macrogyrodactylus clarii* Gussev 1961, a monogenean gill parasite of *Clarias lazera* in Egypt, Arab Gulf J. Sci. Res. Agric. Biol. Sci. 5 (1987) 257–271.
- M.M. El-Naggar, G.C. Kearn, A.E. Hagra, S.Z. Arafa, On some anatomical features of *Macrogyrodactylus congolensis*, a viviparous monogenean ectoparasite of the catfish *Clarias gariepinus* from Nile water, J. Egypt Ger. Soc. Zool. 29 (1999) 1–24.
- S.Z. Arafa, M.M. El-Naggar, G.C. Kearn, Scanning electron microscope observations on the monogenean skin parasite *Macrogyrodactylus congolensis* (Prudhoe, 1957) Yamaguti, Acta Parasitol. 48 (2003) (1963) 163–171.
- P. Ghallii, G. Strona, R. Giovannoni, M. Lavitrano, Head glands of Monogenoidea: morphology, functionality, and potentialities in industrial production of surgery bioadhesives, J. Parasitol. 95 (2009) 1330–1341, <https://doi.org/10.1645/GE-2077.1>.
- I.D. Whittington, B.W. Cribb, Glands associated with the anterior adhesive areas of the monogeneans, *Entobdella* sp. and *Entobdella australis* (Capsalidae) from the skin of *Himantura fai* and *Taeniura lymma* (Dasyatididae), Int. J. Parasitol. 28 (1998) 653–665, [https://doi.org/10.1016/S0020-7519\(98\)00016-2](https://doi.org/10.1016/S0020-7519(98)00016-2).
- I.D. Whittington, B.W. Cribb, Adhesive secretions in the Platyhelminthes, Adv. Parasitol. 48 (2001) 101–224, [https://doi.org/10.1016/S0065-308X\(01\)48006-7](https://doi.org/10.1016/S0065-308X(01)48006-7).
- I.D. Whittington, B.W. Cribb, T.E. Hamwood, J.A. Halliday, Host-specificity of monogenean (Platyhelminth) parasites: a role for anterior adhesive areas? Int. J. Parasitol. 30 (2000) 305–320, [https://doi.org/10.1016/S0020-7519\(00\)00006-0](https://doi.org/10.1016/S0020-7519(00)00006-0).
- I.D. Whittington, B.W. Cribb, Morphology and ultrastructure of the anterior adhesive areas of the capsalid monogenean parasites *Benedenia rohdei* from the gills and *B. lutjani* from the pelvic fins of *Lutjanus carponotatus* (Pisces: Lutjanidae), Parasitol. Res. 85 (1999) 399–408, <https://doi.org/10.1007/s004360050566>.
- K.M. Lyons, The fine structure and function of the adult epidermis of two skin-parasitic monogeneans, *Entobdella soleae* and *Acanthocotyle elegans*, Parasitology 60 (1970) 39–52, <https://doi.org/10.1017/S0031182000077234>.
- K.M. Lyons, The fine structure of the outer epidermis of the viviparous monogenean, *Gyrodactylus* sp. from the skin of *Gasterosteus aculeatus*, J. Parasitol. 56 (1970) 1110–1117, <https://doi.org/10.2307/3277554>.
- I.D. Whittington, W.D. Armstrong, B.W. Cribb, Mechanism of adhesion and detachment at the anterior end of *Neoheterocotyle rhinobatidis* and *Troglocephalus rhinobatidis* (Monogenea: Monopisthocotylea: Monocotylidae), Parasitol. Res. 94 (2004) 91–95, <https://doi.org/10.1007/s00436-004-1171-z>.
- K. Buchmann, Histochemical characteristics of *Gyrodactylus derjavini* parasitizing the fins of rainbow trout (*Oncorhynchus mykiss*), Folia Parasitol. 45 (1998) 312–318.