

HOSTED BY



Contents lists available at ScienceDirect

International Journal of Pediatrics and Adolescent Medicine

journal homepage: <http://www.elsevier.com/locate/ijpam>

Original research article

Severity of Salmonella infection among sickle cell diseases pediatric patients: Description of the infection pattern

Tariq AlFawaz^{a,*}, Omar Alzumar^b, Dayel AlShahrani^a, Mohammed Alshehri^a^a Pediatric Infectious Diseases Section-children Specialized Hospital, King Fahad Medical City, P.O. Box 59046, Riyadh, 11525, Saudi Arabia^b Pediatric infectious Diseases Section-children Hospital, King Saud Medical City, Riyadh, Saudi Arabia

ARTICLE INFO

Article history:

Received 26 December 2018

Received in revised form

18 April 2019

Accepted 26 May 2019

Available online 15 June 2019

Keywords:

Severity

Salmonella

Sickle cell diseases

Children

ABSTRACT

Introduction: Sickle cell disease (SCD) affects millions worldwide. It has a spectrum of clinical manifestations. However, SCD is more prone to have invasive infection compared with normal individual, and one of the main pathogen of concern is salmonella, where the individual with SCD is more susceptible to salmonella infection. Furthermore, several distinct clinical syndromes can develop in children infected with salmonella, depending on both host factors and the specific serotype involved.

Objectives: We aim to describe the infection patterns and whole range of potential complications in children with SCD exposed to invasive salmonella infection.

Method: This is a retrospective observational cohort study which was conducted at King Fahad Medical City (KFMC), Riyadh, Saudi Arabia between 2012 and 2018. All sickle cell patients who are exposed to invasive salmonella infections and treated in our hospital over the last 6 years were included in our study.

Results: Six patients were enrolled in the study, five males and one female with ratio of (M: F) 5:1, age range from 20 months–14 years, and the diagnosis at admission were as follows: (three as vasoocclusive crisis, three as infection) with different kind of infections (three sepsis, three septic arthritis, four osteomyelitis, one meningitis, one myositis, one periorbital cellulitis, one diskitis), where three (50%) suffered multiple sites of infections and the other three (50%) with one site of infection, two (50%) of osteomyelitis patients suffered multifocal infection. Species identification is as follows: (three group D, one group C, and two were not specified), only two occasions where resistant to ciprofloxacin while all others were pan sensitive. Fever was prolonged (take more than seven days to subside even with appropriate therapy and intervention) in five out of six.

Conclusions: Multiple site of infection, severe osteomyelitis, and delay in fever response consolidated the fact of high virulence of salmonella in SCD patients. We did not encounter significant resistant rate to both quinolone and cephalosporin.

© 2019 Publishing services provided by Elsevier B.V. on behalf of King Faisal Specialist Hospital & Research Centre (General Organization), Saudi Arabia. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Sickle cell disease (SCD) affects millions worldwide [1], particularly those of black African and Afro-Caribbean descent and also those from the Mediterranean, Middle East, and parts of

India [2]. It has a wide range of clinical manifestations. However, SCD is more prone to have invasive infection compared with normal individual, and many reasons were encountered [3]. One of the main pathogens of concern is salmonella, where the individual with SCD is more susceptible to salmonella infection due to recurrent vaso-occlusion with intestinal infarction leads to necrosis and increased gut permeability in addition to decreased neutrophil killing [3]. Furthermore, children infected with salmonella are prone to developing multisystem infection such as osteomyelitis, liver and splenic abscesses, and overwhelming sepsis, sometimes with fatal consequences [4].

* Corresponding author.

E-mail addresses: t_alfawaz@yahoo.com (T. AlFawaz), alzomor55@gmail.com (O. Alzumar), daalshahrani@kfmc.med.sa (D. AlShahrani), malshhri@kfmc.med.sa (M. Alshehri).

Peer review under responsibility of King Faisal Specialist Hospital & Research Centre (General Organization), Saudi Arabia.

2. Objectives

Using a case series, we aim to highlight the invasive and virulent nature of salmonella and demonstrate the infection patterns and whole range of potential complications.

3. Method

This is a retrospective observational cohort study which was conducted at King Fahad Medical City (KFMC), Riyadh, Saudi Arabia between 2012 and 2018. All sickle cell patients who are exposed to invasive salmonella infections and treated in our hospital over the last 6 years were included in our study.

Structured data sheet were prepared, and electronic file were reviewed for all included patients.

Ethical approval was obtained from the Institutional Review Board at KFMC.

4. Results

Six patients were enrolled in the study, five males and one female with ratio of (M: F) 5:1, age range from 20 months–12 years, and the diagnosis at admission were as follows: (three as vaso-occlusive crisis, three as infection), patients suffered different kind of infections (three sepsis, three septic arthritis, five osteomyelitis, one meningitis, one myositis, one spondylodiskitis) diagnosed by direct culture or by supported imaging with positive culture at other sites. Four patients (66.7%) suffered multiple site of infections and two (33.3%) with one site of infection, two (40%) of osteomyelitis patients suffered of disseminated osteomyelitis, with most common bone affected is femur (44.4%) (Table 1). In general, preceded illnesses were fever and body pain.

None of our patients was found with apparent immunodeficiency, while two of them were on hydroxyurea. Maximum ESR reported on admission was 120 mm/h while the lowest was 29 mm/h, and maximum WBC was reported is 17.6 (10e/L), five out of six patients (83.3%) ended with surgical intervention with incision and drainage. Salmonella species were isolated from different sites as follow (blood, CSF, joints, bone, Para spinal abscess), and species identification as follows: (three group D, one group C, and two were not specified only non typhi), only two occasions where resistant to ciprofloxacin while all others were pan sensitive. Fever

was prolonged (take more than seven days to subside even with appropriate therapy and intervention) in five out of six (83%). However, four (80%) of osteomyelitis patients ended with chronic osteomyelitis, where three (75%) cured with disabilities which include either limb deformity or vertebral destruction while two (40%) cured with neither complications nor disabilities (Table 1).

5. Discussion

Patients with SCD are known to have an increased susceptibility to severe bacterial infections compared with the population at large [5,6]. Also SCD patients are at higher risk for invasive salmonella disease at both pediatric and adult population [7,8]. In a previous Saudi study, [4] the incidence of *Salmonella* infections was 11.5% compared to only 0.64% during the same period in the general hospital population. In London, 52.3% of all bacteremia over a 15-year period in SCD patients were found to be due to *Salmonella*, compared to 0.4% in non-SCD individuals. Most of these infections were *Salmonella typhimurium*, a common food-borne pathogen [9].

In our review, *Salmonella* can affect all childhood age with diversity of clinical presentation in addition to a variety of clinical diseases predominantly (15 out of 19 episodes = 79%) of musculo-skeletal diseases where osteomyelitis consists of 60%.

In SCD, *Salmonella* is the most common agent causing osteomyelitis, followed by *S. aureus*, then Gram-negative enteric bacteria [10]. In long-term retrospective reviews from the USA [11] and Saudi Arabia [12], *Salmonella* accounted for 57% and 41.7% of cases of acute osteomyelitis, respectively. In our study, osteomyelitis encountered significant number (47.3%) of invasive salmonella episodes in SCD, corresponding with what have been reported in the previous studies [4,9] which were 42% and one third respectively, nevertheless; the difference in population age and number of cases should be considered. However, in our study, osteomyelitis was much severe than other series where 40% of osteomyelitis patients suffered of disseminated osteomyelitis, and the majority underwent multiple interventions and required prolonged therapy, and the majority ended with chronic osteomyelitis. However, B. L Atkins et al., described such severity [13].

The extraordinary susceptibility of patients with SCD to *Salmonella* osteomyelitis was first described fully in 1951 by Hodges and Holt [14] and has since been confirmed in several studies [15,16]. However, the susceptibility of SCD patients to *Salmonella*

Table 1
Patients description.

	Age	Sex	Diagnosis	Fever	Complication	Species	Treatment	Duration	Outcome
Case 1	3 Y	M	Femur Osteomyelitis	Short	Chronic Osteomyelitis	Group D	Ceftriaxone I.V. Cefdinir P.O.	Four months	Cure
Case 2	10 Y	M	Sepsis	Prolonged	None	Group D	Ceftriaxone 4 weeks/Cipro 2 weeks	Four weeks	Cured (Fever continued 3 weeks)
Case 3	3 Y	M	Knee septic arthritis, Femur Osteomyelitis	Prolonged	Chronic osteomyelitis	Group D	Ceftriaxone I.V. + Cefdinir P.O.	Four months	Cure with disabilities*
Case 4	12 Y	M	Sepsis, Meningitis, Knee septic arthritis, Shoulder, Femur, Humerus: Disseminated osteomyelitis,	Prolonged	Disseminated infection, Disseminated osteomyelitis, chronic osteomyelitis	Non-Typhi	Ceftriaxone I.V.+ Ciprofloxacin I.V., then Ciprofloxacin P.O.	5 months IV, 10 months PO	Cure with disabilities*
Case 5	20 months	F	Psoas myositis, Hip septic arthritis, Femur osteomyelitis	Prolonged	None	Group C	Ceftriaxone I.V. (Cipro R)*	six weeks	Cure
Case 6	11 Y	M	Sepsis, shoulder, Mandible, Lumbar osteomyelitis, Spondylodiskitis, Para spinal abscess	Prolonged	Disseminated infection, Disseminated osteomyelitis, intervertebral destruction,	Non-typhi	Ceftriaxone I.V. Bactrim I.V. 6 weeks, then Cefdinir P.O., Bactrim P.O. (4 months), (Cipro R)*	six months	Cure with disabilities*

*R = Resistant.

*Disabilities = Limb deformity – Vertebral destruction.

osteomyelitis has not been fully explained, but certain factors deserve emphasis. It has been proposed that repeated intravascular sickling and vaso-occlusive episodes devitalize the gut wall, leading to frequent seeding of *Salmonella* organisms from the bowel into the blood stream [16]. The presence of repeated marrow thrombosis, infarction and necrosis would make the bone a favored site of localization, leading to osteomyelitis and septic arthritis.

Interestingly, non-SCD patients usually became afebrile within 2–3 days after initiation of adequate antimicrobial treatment for salmonella bacteremia [17], unlike in SCD patients where fever persists more which could be explained by prolongation of apoptotic cell death that allows the bacteria to persist in the host cells for a longer period [18], or due to high initial immune response triggered by bacteremia [19].

In our series, we do not encounter multi resistant salmonella species. However, the emergence of *Salmonella* serotypes resistant to quinolones which increasing through the time in multiple area over world [20–22], and to less extent to cephalosporin where both poses a new challenge in treating infected patients. In our series, we encountered only two isolates (33%) of salmonella were resistant to quinolone which consider relatively low in comparison with prevalence in other area of the world where it could reach to 70% [22], but we never encountered resistant to cephalosporin either in SCD patient nor in non SCD patient.

6. Conclusions

Invasive *Salmonella* infection can have catastrophic consequences in pediatric patients with sickle cell disease. There is an emphasis for early recognition and therapeutic intervention in all cases. Multiple site of infection and severe bone destructions in addition to delay in fever response consolidated the fact of high virulence of salmonella in SCD patients. Presently, aggressive treatment with antibiotics and early recognition of complications are the mainstay of management. We encountered relatively low resistant rate to quinolone and no resistant to cephalosporin. Additionally, health promotion strategies to prevent infection play an important role.

We are Tariq AlFawaz, Omar Alzumar, Dayel AlShahrani, Mohammed Alshehri; authors of paper: Severity of *Salmonella* infection among sickle cell diseases pediatric patients: description of the infection pattern:

Agreed to submit our paper to international journal of pediatric and adolescent

And our work was approved by the Institutional Review Board at KFMC.

References

- [1] NHS Sickle cell and thalassaemia screening programme. Available at: <http://www-phm.ums.ac.uk/haemscreening>. [Accessed April 2018].
- [2] Dick M. Sickle cell disease in childhood. Standards and guidelines for clinical care. UK Forum on Haemoglobin Disorders; 2007 October.
- [3] CatherineBooth Babalusa, Obaro Stephen K. Infection in sickle cell disease: a review. Int J Infect Dis January 2010;14(Issue 1). e2-e12.
- [4] Elbasher Ali Magzoub, Al- Salem Ahmed Hassan, Aliama Abdulaziz. *Salmonella* as a causative organism of various infections in patients with sickle cell disease. Ann Saudi Med 2003;23(6). NOVEMBER-DECEMBER.
- [5] Barrett-Conner E. Bacterial infection and sickle cell anemia. Medicine 1971;50:96–112.
- [6] Pearson HA. Sickle cell anemia and severe infections due to encapsulated bacteria. J Infect Dis 1977;136:S25–30.
- [7] Gordon MA. *Salmonella* infections in immunocompromised adults. J Infect 2008;56(6):413–22.
- [8] Sommet J, Missud F, Holvoe t L, Ithier G, Lorrot M, Benkerrou M, et al. Morbidity among child travelers with sickle-cell disease visiting tropical areas: an observational study in a French tertiary care center. Arch Dis Child 2013;98(7):533–6.
- [9] Workman MR, Philpott-Howard J, A.J. Bellingham *Salmonella* bacteremia in sickle cell disease at King's College Hospital: 1976–1991. J Hosp Infect 1997;34:195–9.
- [10] Almeida A, Roberts I. Bone involvement in sickle cell disease. Br J Haematol 2005;129:482–90.
- [11] Chambers JB, Forsythe DA, Bertrand SL, Iwinski HJ, Steflik DE. Retrospective review of osteoarticular infections in a pediatric sickle cell age group. J Pediatr Orthop 2000;20:682–5.
- [12] Sadat-Ali M. The status of acute osteomyelitis in sickle cell disease. A 15-year review. Int Surg 1998;83:84–7.
- [13] Atkins BL, Price EH, Tillyer L, et al. *Salmonella* osteomyelitis in sickle cell disease children in the East End of London. J Infect March 1997;34(Issue 2): 133–8.
- [14] Wright J, Thomas P, Serjant GR. Septicemia caused by salmonella infection: an overlooked complication of sickle cell disease. J Pediatr 1997;130(3):394–9.
- [15] Hook EW, Campbell CG, Weens HS. *Salmonella* osteomyelitis in patients with sickle cell disease anemia. N Engl J Med 1957;257:403–7.
- [16] Adeyokunnu AA, Hendrickse RG. *Salmonella* osteomyelitis in childhood: a report of 63 cases seen in Nigerian children of whom 57 had sickle cell anemia. Arch Dis Child 1980;55:175–84.
- [17] Tsai MH, Huang YC, Lin TY, Huang YL, Kuo CC, Chiu CH. Reappraisal of parenteral antimicrobial therapy for nontyphoidal *Salmonella* enteric infection in children. Clin Microbiol Infect 2011;17:300–5.
- [18] Gulig PA, Danbara H, Guiney DG, Lax AJ, Norel F, Rhen M. Molecular analysis of spv virulence genes of the *Salmonella* virulence plasmids. Mol Microbiol 1993;7:825–30. <https://doi.org/10.1111/j.1365-2958.1993>.
- [19] Eng Shu-Kee, Pusparajah Priyia, Nurul-Syakima Ab Mutalib, et al. Alaysia View further author information *Salmonella*: a review on pathogenesis, epidemiology and antibiotic resistance. J Front Life Sci 2015-issue;8:284–93.
- [20] Frost JA, Kelleher A, Rowe B. Increasing ciprofloxacin resistance in salmonellas in England and Wales 1991–1994. J Antimicrob Chemother 1996;37:85–91.
- [21] Herikstad Hallgeir, Hayes Peggy, Mokhtar Mohamed, et al. Emerging quinolone-resistant *Salmonella* in the United States, emerging infectious diseases, vol. 3; July–September 1997. No. 3.
- [22] Hung-MingChen, YueWang, Lin-HuiSu, Cheng-HsunChiu. Nontyphoid *Salmonella* infection: microbiology, clinical features, and antimicrobial therapy. Pediatr Neonatol June 2013;54(Issue 3):147–52.