



ELSEVIER



PET/MRI vs PET/CT in Head and Neck Imaging: When, Why, and How?

Joshua L. Ryan, MD, Vasantha D. Aaron, MD, and Justin B. Sims, MD

The increasing availability of hybrid PET/MRI systems has led to a breadth of new publications and opportunities for use of PET/MRI. While PET/CT has been a valuable tool for oncologic staging, including head and neck malignancy, there are several theoretical and practical advantages a PET/MRI system would have over PET/CT in head and neck imaging. This review article discusses the established role of PET/CT, early evidence for the role of PET/MRI, and protocol considerations for both PET/CT and PET/MRI as they apply to head and neck imaging. *Semin Ultrasound CT MRI 40:376-390 © 2019 Elsevier Inc. All rights reserved.*

Introduction

Malignancy in the head and neck encompasses a broad array of diagnoses, including mucosal surface (epithelial) malignancies of the sinonasal cavity and aerodigestive tract, primary and secondary salivary gland malignancies, thyroid cancer, skin cancer, orbito-ocular malignancies, skeletal involvement by tumors, and malignancies involving the lymphoid tissues. The most common subtype of head and neck malignancy is mucosal surface squamous cell carcinoma (SCCa), which is the histologic type in greater than 90% of mucosal surface malignancies, and may arise in the sinonasal mucosa, nasopharynx, oral cavity, oropharynx, hypopharynx, and larynx.¹ In 2012, a worldwide estimated 300,400 new cases and 145,400 deaths arose from oral cavity cancer alone.² It is estimated that in 2019, there will be 53,000 new cases of cancer within the oral cavity or pharynx diagnosed in the United States.³ Well known risk factors for head and neck cancer included tobacco use and alcohol, which are also known risk factors for other malignancies in the aerodigestive tract (lung and esophagus).¹ Human Papilloma Virus (HPV) is an increasingly recognized risk factor for head and neck cancer, which has dramatically changed the demographic of head and neck cancer patients and shifted the typical age of onset of head and neck cancer downward.^{1,4,5} Of note, due to increasing incidence of HPV associated head and neck malignancy, the American Joint Committee on

Cancer (AJCC) now includes a separate category for staging of HPV-related oropharyngeal cancer.⁴

In the primary initial workup of a head and neck malignancy, imaging plays an essential role, and is generally believed to be more useful in advanced tumor stages.⁴ TNM staging developed by the AJCC is subtyped into multiple sites of origin.⁴ Accurate staging is important in treatment planning and prognosis. Owing to the complex anatomy of the head and neck region, imaging plays a vital role in supplementing diagnosis, treatment planning, and disease monitoring in head and neck malignancy.⁶ In clinical practice, specific imaging is typically selected on a case by case basis, and usually involves contrast enhanced Neck CT or MRI. For patients who are suspected to have more advanced disease (stage III or IV), FDG PET/CT is a useful tool that provides high sensitivity for distant disease.^{1,6,7}

Whole body PET/MR imaging systems became available in 2010.⁸ Since its introduction, hybrid PET/MR has grown in availability and has begun to be adopted for clinical use, although is still not widely available in the medical landscape. One of the primary theoretical advantages of Positron emission tomography/magnetic resonance imaging (PET/MRI) over PET/CT is the improved soft tissue contrast of MRI compared to CT. This is especially advantageous when imaging the complex anatomical structures that reside within the head and neck.⁸ PET/MRI also allows for a significant reduction in radiation dose to the patient compared to PET/CT.⁹

This chapter will discuss the current role of Fluorine-18 fluorodeoxyglucose positron emission tomography/computed tomography (18F-FDG PET/CT) in head and neck imaging and contrast the more established role of PET/CT against the emerging technology of hybrid PET/MRI.

Indiana University School of Medicine, Indianapolis, IN.

Address reprint requests to Joshua Ryan, MD, Indiana University School of Medicine, Department of Radiology and Imaging Sciences, 550 N. University Blvd. Room 0663, Indianapolis, IN 46202. E-mail: ryanjl@iu.edu

Additionally, this article will briefly review protocolling for PET/CT and PET/MRI.

Established Role of 18F-FDG PET/CT in Head and Neck Cancer

FDG PET/CT is commonly performed in clinical practice for the workup and staging of head and neck malignancies. In many cases, patients have already undergone CT or MRI and endoscopic biopsy prior to undergoing a PET/CT, and an established tissue diagnosis is known. The National Comprehensive Cancer Network (NCCN) guidelines generally recommend FDG PET/CT in workup of stage III or IV disease, or as clinically indicated for most sites of origin.¹

Primary Tumor

A recent study of 35 patients by Chaput et al showed greater diagnostic accuracy with higher sensitivity of FDG PET/CT (83%) compared to MRI (63%) for detection of T1-T2 category head and neck SCCa, suggesting efficacy of FDG PET/CT in lower stage disease.¹⁰ When PET is performed with a contrast enhanced CT, diagnostic CT images can be obtained from a single examination and may obviate the need for additional imaging for staging (CT or MR).¹¹ Of note, at our institution, all PET/CT exams are performed with iodinated IV contrast unless there are medical contraindications for the patient (contrast allergy or renal insufficiency). PET/CT protocols are discussed in detail later in this chapter.

Nodes and Metastases

FDG PET/CT has also been established as the optimal imaging study for staging of nodal metastases and distant metastatic

disease, backed by recommendations from the AJCC and NCCN for evaluation in stage III and IV disease (Fig. 1).^{1,4} A meta-analysis by Sun et al showed greater pooled per neck level sensitivity of FDG PET CT (84%) for detection of regional nodal metastases over CT or MR (63%).¹² In addition, a recent prospective cohort study by Rohde et al demonstrated superiority of PET/CT for detection rates of distant metastases and second primary malignancy over the combination of radiographs, CT, and MRI based on geographic preferences of Europe and United States.¹³

Cancer of Unknown Primary

The use of 18F-FDG PET/CT is also recommended for diagnoses beyond the workup of a known primary site of malignancy within the head and neck. Notably, PET/CT has a role in helping identify the primary site in cancer of unknown primary, which is typically discovered as enlarged cervical lymph nodes with pathology of SCCa, adenocarcinoma or undifferentiated epithelial tumors (Fig. 2).¹ A meta-analysis by Zhu et al noted that 18F-FDG PET/CT had an overall detection rate of 0.44, pooled sensitivity of 0.97, and specificity of 0.68 for primary site detection in an analysis of 7 studies.¹⁴

Mucosal Melanoma

18F-FDG PET/CT should also be considered for staging and screening for metastatic disease in the setting of mucosal melanoma, which is generally seen in an older population with median age of 70 at diagnosis.^{4,15} Mucosal melanoma is a rare entity, encompassing only 1.3% of known melanoma cases, with about 55% occurring in the head and neck region.¹⁶ Most of the head and neck mucosal melanoma cases arise in the paranasal sinuses and nasal cavity.⁴ Unfortunately, mucosal




Figure 1 PET/CT and MRI Nodal Evaluation in a patient with recurrent nasopharyngeal carcinoma. Fused axial FDG-PET/CT (A) shows a hypermetabolic right retropharyngeal lymph node (black arrow). Unfused CECT (B) shows corresponding enhancing retropharyngeal lymph node (black arrow). Precontrast T1-weighted precontrast image (C) shows the metastatic lymph node (blue arrow) to have similar signal intensity to adjacent longus colli muscle. Axial T1-weighted postcontrast fat-saturated image (D) shows enhancement of the nodal metastasis (blue arrow), along with enhancement into the adjacent longus colli muscle (red arrow), suggestive of extra nodal extension. There is also enhancement of the soft tissues adjacent to the right internal carotid artery (yellow arrow), compatible with arterial encasement. (Color version of figure is available online.)




Figure 2 Cancer of unknown primary in a patient with palpable right supraclavicular mass. Biopsy showed metastatic squamous cell carcinoma without known primary. Fused axial FDG-PET/CT (A) for staging shows a right supraclavicular mass (green arrow) with peripheral hypermetabolism and central necrosis. The primary site of malignancy was not identified on this PET/CT. Fused axial FDG-PET/CT (B) also shows an additional, small, hypermetabolic metastasis noted in the right gluteus maximus (blue arrow). Unfused axial CECT (C) at the level of the lesion shows the gluteal metastasis is not visible on CT alone (yellow circle). (Color version of figure is available online.)

melanoma is an aggressive entity, and most patients progress to metastatic disease.¹⁷ Metastases can occur most commonly in lung, bone, peritoneum, and central nervous system; thus PET/CT is an ideal imaging modality to evaluate for distant metastases.¹⁷ A brain MRI is also usually indicated for evaluation of brain metastases in melanoma patients.

Salivary Gland Malignancy

According to the AJCC, there is no established role for PET/CT for the evaluation of primary salivary gland malignancy, with CT and MR potentially providing a complementary role in staging and diagnosing major salivary gland tumors.⁴ A recent prospective study by Park et al showed 18F-FDG PET/CT to be useful in the detection of metastases and additional primary cancers in these patients, along with a predictive role in survival outcomes.¹⁸ This study suggests that expansion of the current role of 18F-FDG PET/CT in primary salivary gland cancer may be an area of further research.

Treatment Evaluation and Post-Treatment FDG PET

In the post-treatment setting following radiation alone or chemotherapy and radiation, NCCN guidelines state that 18F-FDG PET should be considered after therapy in patients who exhibit clinical response to treatment with a wait time of at least 12 weeks to minimize post-treatment uptake and false positive results (Fig. 3).¹ A meta-analysis performed by Gupta et al showed excellent sensitivity (79.9%), specificity (87.5%), and negative predictive value (NPV; 95.1%), but a relatively

low positive predictive value (PPV; 58.6%).¹⁹ This meta-analysis suggests that PET/CT is a more accurate modality for informing clinical decision making when the results are negative.¹⁹ While it has generally been established that post-therapy 18F-FDG PET/CT should occur around 12 weeks or greater post-therapy, the NCCN suggests that 18F-FDG PET/CT can also be considered earlier in patients who have evidence of persistent or progressive disease.¹ However, there is currently no well-established role of 18F-FDG PET/CT in assessing early response while undergoing treatment.²⁰

The recently published PET-NECK randomized controlled trial performed by Mehanna et al showed noninferiority of 18F-FDG PET/CT surveillance after chemotherapy and radiation in patients with advanced nodal (N2 or N3) disease compared to patients who underwent radical neck dissection.²¹ The results of the PET-NECK study show a viable noninvasive alternative for disease surveillance in patients with treated metastases.

Another recently published multicenter trial by Lowe et al, the ACRIN 6685 trial, showed a 18F-FDG PET/CT to have a high NPV (87%) for a clinically N0 neck in patients with SCCa.²² This study also noted that surgical planning was changed in approximately 21% of patients after surgeon review of the PET/CT. The findings from this study demonstrate the value of 18F-FDG PET/CT in determining treatment course for clinically N0 necks.

Prognostic Value of 18F-FDG PET/CT

There has also been a focus in the recent literature on the prognostic value of 18F-FDG PET/CT in patients with head and neck malignancy. The most well-known and widely




Figure 3 Treatment response with staging and postchemotherapy and radiation treatment FDG-PET/CTs in a patient with T3, N0, M0 squamous cell carcinoma of the right palatine tonsil and base of tongue. Fused axial FDG-PET/CT (A) for staging shows a hypermetabolic right palatine tonsil and base of tongue mass (red arrow). Fused axial FDG-PET/CT 3 months postcompletion of chemotherapy and radiation (B) shows no residual hypermetabolic activity in the right tonsillar fossa (yellow circle), compatible complete local response to therapy. Unfortunately, 3 month post-therapy fused axial FDG-PET/CT (C) also shows a new, hypermetabolic and partially necrotic left upper lobe mass compatible with distant metastasis and progression of disease (blue arrow). (Color version of figure is available online.)

reported semiquantitative measurement parameter in PET is the maximum standardized uptake value (SUV) within a volume of interest. SUV has been established as a prognostic indicator in head and neck SCCa, with increased SUV of primary tumors being a poor prognostic factor.²³ However a recently published meta-analysis by Wang et al demonstrated superior prognostic predictive capabilities of metabolic tumor volume and total lesion glycolysis compared to SUV max.²⁴ It should be noted that total lesion glycolysis and metabolic tumor volume are not yet widely used, possibly because they are difficult to perform and require specialized software. Regardless of the measurement parameter employed, 18F-FDG PET/CT has established itself as an important prognostic tool for clinicians. 18F-FDG PET/CT has also shown potential value in the radiation therapy treatment planning, as treatments such as intensity-modulated radiation therapy depend on accuracy of tumor size.²⁰

18F-FDG PET/CT is a versatile imaging modality that can help guide clinical decision making in a variety of settings. However, there are known advantages MRI has over CT in head and neck imaging, including better soft tissue contrast and spatial resolution, and lack of ionizing radiation. Combined with the fact that many patients with head and neck malignancy undergo both, these limitations provide a framework for the clinical use of PET/MRI in many head and neck settings.

PET/MRI in Imaging of Head and Neck

Retrospective Fusion

The earliest studies that examined the feasibility of PET combined with MRI in head and neck imaging were fusion studies between separately performed MRI and PET. Nakamoto et al published a retrospective fusion study in 2009 that showed the benefit of PET fusion with MRI in post-treatment patients suspected of having disease recurrence, though there was minimal benefit in initial staging.²⁵ Retrospective fusion

study by Huang et al showed that fused PET/MR provided greater performance than PET/CT, standalone CT, or standalone MRI in detecting tumor size and local invasion in advanced buccal SCCa.²⁶ Kanda et al performed a retrospective fusion study in 30 patients with oral cavity or hypopharyngeal cancer, which showed greater accuracy of T category assessment with fused PET/MRI and MRI compared to PET/CT, and greater accuracy of N category assessment with fused PET/MRI and PET/CT over MRI alone.²⁷

Initial Experiences With Hybrid PET/MRI

Boss et al were the first to publish a study demonstrating the feasibility of a simultaneous PET/MRI system in a study of 8 patients.²⁸ Their results showed no image degradation of the accompanying MR images, with the added benefit better spatial resolution of PET images performed with MRI as compared to the PET/CT, due to the smaller diameter of PET detector ring and smaller scintillation crystal sizes on their PET/MRI system.²⁸ These initial results were supported by the findings of Platzek et al, who showed no MRI artifacts in PET images in a 20-person prospective study in patients with head and neck SCCa.²⁹ Another early study showed a good qualitative comparison between PET/MRI and PET/CT images and a correlation between SUVs between the PET/MRI and PET/CT.³⁰ Along with correlating SUVs between PET/MRI and PET/CT, Covelto et al demonstrated the multiparametric abilities of PET/MRI in the evaluation of head and neck imaging (SUVs, diffusion, and perfusion parameters).³¹

PET/MRI vs PET/CT and Conventional Imaging: The Evidence

TNM Staging

Following the feasibility studies, the next step in research of PET/MRI for image quality in the head and neck was to

compare PET/MRI to PET/CT and to conventional imaging (MRI and CT). Several studies have shown the noninferiority of PET/MRI in staging these malignancies. For example, a prospective study by Kuhn et al evaluated 150 patients with head and neck cancer presenting for a PET/CT evaluation. Using a trimodality PET/CT-MR setup, their results showed that contrast enhanced PET/MR performed significantly better than contrast enhanced PET/CT in evaluation of primary tumors; however, there was no significant difference in evaluation of lymph nodes.³² Another study by Platzeck et al noted that there was no significant difference between PET/MR and MRI and PET alone in the accuracy of lymph node staging in 38 patients.³³ A small study of 17 patients by Kubiessa et al showed no statistically significant difference in diagnostic performance (sensitivity, specificity, PPV, and NPV) between PET/CT and PET/MRI in evaluation of malignant findings in patients with head and neck cancer.³⁴ Partovi et al also found comparable results between PET/CT and PET/MRI for detection of lymph nodes and metastases in their prospective study of 14 patients with head and neck cancer (Fig. 4).³⁵

A small study of 16 patients with laryngeal cancer by Cavaliere et al showed that PET/MRI is useful for the primary staging of laryngeal cancer and is comparable to PET/CT.³⁶ Their study showed a high correlation between imaging findings and endoscopic/histologic evaluation for PET/MRI. They also reference the ability of MRI in determining cartilage invasion in suggesting potential for PET/MRI in the evaluation of laryngeal cancer.^{36,37} Cavaliere et al also discuss the advantages of being able to collect metabolic, anatomical, and functional parameters in a single imaging study.³⁶

Stronger evidence for the use of PET/MRI over PET/CT is seen in a recently published prospective study of 113 patients by Chan et al, which recommended the use of whole-body PET/MRI as a single imaging modality in the




Figure 4 PET/MRI with lymph node metastases in a patient with oropharyngeal squamous cell carcinoma. Fused coronal FDG-PET/MRI with T1-weighted postcontrast fat-saturated image shows mildly FDG avid cervical lymph nodes (green arrows) due to nodal metastases. (Color version of figure is available online.)

staging of nasopharyngeal carcinoma. In this study, PET/MRI outperformed PET/CT and head and neck MRI for T category assessment and showed higher PPV for assessment of distant metastatic disease.³⁸

Follow-Up Imaging and Disease Recurrence

The theoretical advantage of PET/MRI over PET/CT may be greatest in the post-treatment setting, as the superior soft tissue contrast resolution of MRI may provide the clinician and radiologist more evidence to determine whether FDG uptake is true recurrence, physiological activity, or expected post-treatment related uptake. To this effect, Queiroz et al suggest that PET/MRI may be superior to PET/CT in evaluating unclear FDG findings in follow-up patients with head and neck cancer, but their study showed no significant difference in diagnostic accuracy between PET/CT and PET/MRI.³⁹ A retrospective analysis by Schaarschmidt et al showed no significant difference in correct classification of incidental radiotracer uptake in the head and neck on PET/MRI compared to PET/CT in 81 oncologic patients.⁴⁰ In a separate publication, Schaarschmidt et al also noted no significant difference between PET/MRI, PET/CT, and MRI in local tumor staging or cancer recurrence in a small study of 25 patients.⁴¹

In a study demonstrating the multiparametric capabilities of PET/MRI, Becker et al showed excellent results for detection of recurrent disease and tumor classification of PET/MRI performed with diffusion weighted imaging (DWI) in the post-treatment setting.⁴² This study notes the value of concordant imaging results between PET, MRI, and DWI in predicting tumor recurrence, and discordant results corresponding to benign post-treatment changes.⁴² These results differ from an earlier study by Queiroz et al, which concluded that DWI was not helpful in the evaluation of head and neck malignancy with PET/MRI.⁴³ They also highlight the findings of Rasmussen et al that 18F-FDG uptake and DWI have independent but complementary roles in head and neck imaging.⁴⁴ Based on the preliminary research, further studies will be necessary to define the role of multiparametric imaging as part of a PET/MRI examination in the head and neck.

Cancer of Unknown Primary and Nonsquamous Cell Carcinoma of the Head and Neck

18F-FDG PET/CT has been firmly established as a valuable tool in the work-up of cancer of unknown primary manifesting as cervical lymph node metastases, and PET/MRI hold promise in this setting, as well.^{1,14} In a study of 20 patients, PET/CT and PET/MRI equally identified 11/20 primary cancer sites.⁴⁵ They also noted superior assessment of cervical lesions on PET/MRI and superior detection of pulmonary metastases on PET/CT, although no statistical significance was determined.⁴⁵ To date, no studies have been performed that specifically assess the

utility of PET/MRI in salivary gland malignancies or mucosal melanoma. This may be in part due to the nonestablished role in of FDG-PET/CT in assessing salivary gland malignancies, and the rarity of mucosal melanoma.

Specific Situations Where MRI is Preferred Over CT

There are situations in which the superiority of MRI over CT can be used to infer that PET/MRI may be a preferred modality to PET/CT. One is the evaluation of perineural spread, where MRI is preferred over CT because of its superior soft tissue contrast resolution⁴⁶ (Fig. 5). In reference to PET/MRI, a prospective study by Sekine et al, which used a trimodality PET/CT + MRI system, showed that PET/MRI detected 6 cases of perineural spread, while PET/CT detected only 5.⁴⁷

MRI is also the imaging modality of choice in the evaluation of salivary gland malignancy and helps with accurate localization and locoregional extension of salivary gland tumors, as well as detection of perineural spread.⁴⁸ MRI is also generally regarded as superior to CT in evaluation of orbital disease, due to superior contrast resolution and ability to assess extent of disease involving the orbital compartment, perineural spread, and intracranial extension.⁴⁹ Evaluation of subtle skull base invasion and marrow replacement, along with intracranial extension is also better appreciated by MRI, which is currently the gold standard for evaluation of nasopharyngeal carcinoma.³⁸ MRI is also the preferred modality when imaging the sella and parasellar region.⁵⁰ Specifically, the superior soft tissue contrast resolution would be better for staging in evaluating for carotid encasement and cavernous sinus extension.

Another clinical situation is evaluating the oral cavity and oropharynx in patients with dental hardware. Streak artifact from CT can limit the diagnostic confidence and lesion conspicuity, while MRI can also have signal void due to metallic

artifact, it is generally less of a hindrance in diagnostic confidence than CT artifact (Fig. 6).

Summary of Findings and Suggestions for Clinical Use of PET/MRI

The first portion of this chapter has briefly reviewed the evidence for the use of both 18F-FDG PET/CT and PET/MRI in the setting of head and neck cancer. As described, 18F-FDG PET/CT plays a significant and versatile role in head and neck imaging, and its recommendation by the AJCC and NCCN in multiple settings further emphasizes its established role in medical imaging. However, while the data comparing PET/CT to PET/MRI are limited in number of studies and patients per study, the preliminary results show that PET/MRI is comparable to PET/CT in multiple settings. It is therefore reasonable that PET/MRI should be considered for use in individually selected cases where the benefit of MRI is perceived to be of greater than CT.

Specific examples where PET/MRI could be considered include patients with significant dental artifact that hinders the diagnostic confidence of CT. PET/MRI should also be considered in patients who have suspected perineural disease or have primary tumors that are at high risk for perineural tumor spread, such as salivary gland neoplasms, orbital tumors, and skin cancer. In patients with head and neck malignancy who would be deemed to require a separate MRI and PET/CT for staging or restaging, PET/MRI can provide the convenience of a “one stop shop” and should be considered. PET/MRI should also be considered in evaluation of locally advanced tumors where differentiation between tumor and adjacent structures would have an impact on clinical or surgical decision making, including staging of disease. PET/MRI should also be considered for primary tumors near the skull base, orbits, oral cavity, or in primary nasopharyngeal carcinoma. At our institution, the selection of head and



Figure 5 PET/MRI with perineural tumor spread in a patient with squamous cell carcinoma arising in the maxillary sinus. Fused coronal FDG-PET/MRI with T1-weighted postcontrast fat-saturated image (A) shows a hypermetabolic, enhancing mass in the left maxillary sinus with local extension. Coronal T1-weighted postcontrast fat-saturated image (B) shows soft tissue thickening and enhancement along the superior and inferior orbital fissures (yellow arrow), which is markedly FDG avid on the fused FDG-PET/MRI (C) compatible with perineural tumor spread. (Color version of figure is available online.)




Figure 6 Dental artifact in a patient with suspected recurrent squamous cell carcinoma in the right palatine fossa on clinical exam. Axial CECT (A) shows streak artifact from dental amalgam limiting evaluation of the oropharynx, although there is subtle fullness of the right palatine fossa in the location of clinically suspected recurrence. Axial T1-weighted postcontrast fat-saturated image (B) obtained during FDG-PET/MRI acquisition shows reduction in metal artifact from dental amalgam compared to CT, allowing the asymmetric soft tissue thickening of the right palatine tonsil to be better appreciated. Fused axial FDG-PET/MRI with T1-weighted postcontrast fat-saturated image (C) shows avid FDG uptake in the area of masslike thickening of the right palatine tonsil, compatible with recurrent squamous cell carcinoma in this patient.

neck cancer patients for PET/MRI is typically done in a multidisciplinary approach with multidisciplinary tumor boards providing an ideal platform for physician-to-physician discussions about the capabilities of PET/MRI and the appropriate selection of patients for PET/MRI.

PET/CT Protocol Considerations

Background

PET/CT imaging protocols vary greatly, and imaging of head and neck malignancies highlights some of the differences. In this section, we will review the PET and CT parameters that can be manipulated in order to optimize head and neck scans. More general PET imaging questions, such as appropriate blood glucose levels or 18F-FDG dosing, will not be discussed.

CT Parameters

The most fundamental question in protocoling the CT portion of a PET/CT study is determining whether the CT should be “diagnostic” or performed only for anatomical localization and attenuation correction (AC). The latter “low-dose CT” protocol is the most common approach in the United States with some predominantly academic institutions favoring the former. Some institutions perform a combination of both low-dose and diagnostic CTs. For example, performance of a low-dose CT from the skull to the thigh may be followed by a dedicated, diagnostic, breath-hold chest CT in patients with lung cancer.

Localization and AC protocols employ low mAs (30-80) and varying kVp settings and may reduce radiation dose by as much as 50%-80%⁵¹; however, they come at the cost of increased noise and artifact and do not typically use

iodinated contrast. This means that there is little soft tissue contrast and minimal information regarding tumor characterization or extent of local spread can be gleaned from such studies.

The exact definition of diagnostic CT may differ among radiologists; however, adequate spatial resolution with acceptable signal to noise ratio, increased soft tissue contrast, and enough radiation dose to allow for diagnostic characterization of pathologic processes are necessary to consider a study diagnostic.⁵² This requires higher mAs, which is ideally modulated based on the patient topogram, and in patients without contraindications, iodinated contrast administration. The advantages of contrast-enhanced CT (CECT) over non-contrast CT (NECT) in evaluating many pathologic processes are well established and include increased lesion conspicuity, lesion characterization, and diagnostic confidence. However, there are very few clinical studies evaluating its benefits in the setting of 18F-FDG PET/CT and no study specifically addresses this question in the setting of head and neck malignancies. There are a few studies of other tumor sites, which have shown the benefits of using CECT with PET imaging. A small retrospective study comparing CECT vs low dose, NECT PET/CT whole body imaging in patients with different tumor types showed that CECT added value to a majority of scans.⁵³ The greatest benefit in this study was in exact localization of abnormal 18F-FDG uptake on the PET and precise delineation of tumor, which led to a change in management in 21% of patients. The same group confirmed that CECT PET/CT changed the delineation of primary tumor in 63% of patients with nonsmall cell lung cancer compared to NECT PET/CT imaging, and in 20% of patients the CECT led to the detection of other important clinical findings.⁵⁴ Another small retrospective study showed that CECT PET/CT more accurately identified nodal metastatic spread of rectal cancer over NECT PET/CT.⁵⁵

To date, only two studies have been published comparing CECT and NECT PET/CT in head and neck malignancies, however, neither study was designed to evaluate the benefits specific to CECT. One study was performed to determine whether the use of CECT would change interpretation of the PET portion of the scan compared to low-dose NECT PET/CT due to potential differences in AC.⁵⁶ While the study authors concluded that there was no clinically significant difference in interpretation of the PET when using CECT or NECT for AC, the study did not address changes in interpretation or management based on the CTs themselves. The second study compared whole body NECT PET/CT to dedicated head/neck CECT PET/CT.⁵⁷ The dedicated head/neck CECT PET/CT was performed after longer equilibration compared to the whole body NECT PET/CT. In addition, the PET was obtained over a longer time period in the dedicated protocol and other PET acquisition and reconstruction parameters were altered as well. This study design confounded the ability to determine whether the CECT itself led to improved detection of nodal metastases or NECT when performed with PET/CT. Regardless, the study did determine that dedicated head/neck CECT PET/CT identified more cervical nodal metastases compared to whole body NECT PET/CT and suggests that optimizing both PET and CT protocols may significantly alter interpretation and patient management.

We do know that identifying and defining perineural spread, cartilage involvement, necrosis, and nodal extracapsular extension of tumor is crucial in the staging and treatment planning of patients with these malignancies and may only be delineated with diagnostic CECT or MRI. CECT performed with PET may yield all the necessary information and negate the need for additional, separately performed CECT or MRI, which lead to additional cost and medical visits for the patient. Another reason to consider CECT over NECT PET/CT in head and neck malignancies is the potential for avoiding false negative interpretations. Cystic and mucinous tumors or cystic

nodal metastases may show little uptake of 18F-FDG and may not be identified or characterized as abnormal based upon the PET data alone (Fig. 7). Finally, the ability to reconstruct the diagnostic CT into sagittal and coronal views, which are standard series for all head/neck imaging at our institution, and review the images as fused and unfused data sets gives additional, useful information to interpreting physicians.

There are pros and cons to both low-dose NECT and diagnostic CECT PET/CT protocols. The major benefits to performing low-dose NECT with PET include lower radiation dose, limited prescan patient evaluation, minimal CT training for technologists, and no need for physician presence during scans. As mentioned previously, CT images are noisy and show increased artifact without additional information beyond basic localization of tumor sites. Diagnostic CT protocols employing intravenous contrast require fully trained technologists in both PET and CT, prescan evaluation of renal function and contrast allergies, physician presence in case of allergic reaction, and increased radiation dose to the patient. In addition, if iodinated contrast is administered, interpreting physicians must be aware of possible pitfalls to interpretation. While there is no clinically significant change in SUVs when CECT is used for AC of the PET data, artifacts can arise due to large collections of contrast.^{58,59} For example, contrast pooling within a vein may lead to overcorrection of the PET data and what appears to be focal accumulation of the radiopharmaceutical. Careful evaluation of the underlying CT images and review of the nonattenuation corrected PET data are required to determine whether the “hot” spot is a true pathologic lesion or artifact (Fig. 8).

PET Parameters

There is also flexibility in protocoling the PET portion of PET/CTs for head and neck malignancies. The simplest protocol includes a CECT or NECT skull to thigh CT followed by a




Figure 7 Primary tumor and cystic nodal metastasis evaluation. Axial CECT (A) demonstrates fullness of the left nasopharyngeal mucosa with heterogeneous attenuation. A small, hypodense lesion in the right retropharyngeal space (yellow arrow) is able to be appreciated due to the surrounding, enhancing normal tissues and corresponds to a necrotic retropharyngeal lymph node. Fused axial FDG-PET/CT demonstrates a hypermetabolic left nasopharyngeal mucosa consistent with primary malignancy. The necrotic right retropharyngeal lymph node does not show FDG avidity, but remains suspicious for malignancy given its characteristics on CT. (Color version of figure is available online.)




Figure 8 Attenuation correction. Axial 18F-FDG PET attenuation corrected image (A) demonstrates 2 foci of abnormal uptake in the right supraclavicular fossa (white arrows). Axial 18F-FDG nonattenuation corrected PET (B) shows minimal uptake in the right supraclavicular fossa on the same slice (black arrows). Review of the fused axial 18F-FDG PET/CT dataset at the same level (C) demonstrates a large bolus of iodinated contrast material in the right supraclavicular fossa (yellow arrow) causing artifactual accumulation of radiotracer activity.

skull to thigh PET. In this protocol, there is no separate, dedicated neck imaging. This protocol is fast, minimizes the possibility of patient motion, and potentially exposes patients to the least amount of radiation. In head and neck malignancies, dedicated head and neck PET images commonly are obtained at a delayed time point after performance of the skull to thigh PET/CT. Several studies have demonstrated increased sensitivity for detection of small nodal metastases with these types of protocols, however the equilibration time after 18F-FDG injection varies widely from 60 to 150 minutes (Fig. 9).^{57,60,61} At least 1 study has shown that dedicated, delayed neck PET/CT can lead to decreased specificity in detection of lymph node metastases, presumably because reactive nodes demonstrate abnormal uptake.⁶²

Most dedicated head and neck PET acquisitions include longer bed times with reconstructed smaller field of view and

higher matrix/decreased pixel size compared to the whole-body PET scan. For example, a dedicated neck PET protocol may include 6-8 min/bed with a 30 cm field of view and 256×256 matrix, while the skull-thigh PET protocol may employ 2-4 min/bed with a 50 cm field of view and 128×128 matrix. The higher number of counts obtained with the longer bed time in the dedicated PET should compensate for the smaller pixels and potential for increased noise. In addition, decreasing the size of the field of view without changing the reconstruction matrix or even increasing it improves spatial resolution. Slice thickness and amount of zoom should be matched with the dedicated CT performed with the PET as well.

If a dedicated neck PET/CT is performed separate from the skull to thigh images, established protocols include performance of a skull to thigh low dose, NECT PET/CT followed by the dedicated neck PET/CT, which may also be performed as a low dose, NECT PET/CT or as a diagnostic, CECT PET/CT. Performing the dedicated PET/CT after the skull to thigh allows for longer equilibration time. Alternatively, the dedicated neck PET/CT may be performed after a clavicle to thigh PET/CT or vice versa. In this protocol, the patient's arms are down for the neck portion of the study, which limits beam hardening and streak artifact on the CT and, therefore, subsequent artifact on the attenuation corrected PET images. While the technologist is setting-up the clavicle to thigh portion of the study, the patient is instructed to raise the arms above the head, which then limits artifact upon scanning of the torso. In addition to producing images with significantly less artifact, this protocol decreases radiation dose to the neck, which is included in only 1 CT scan. Finally, parameters such as field of view and reconstruction matrix can be optimized for the dedicated neck PET/CT and adjusted for the clavicle to thigh PET/CT, which may require different parameters such as a larger field of view.

PET/CT Protocol With Diagnostic CECT

There are many ways to optimize both the PET and the CT protocols in the imaging of head and neck malignancies. At our institution, we perform clavicle to thigh PET/CT without contrast followed by dedicated orbit to aortic arch, CECT PET/CT of the neck, which closely mirrors the parameters of standalone diagnostic neck CTs. For example, the soft tissue neck CT protocol at our institution on a 256 slice Philips iCT scanner includes scanning at 120 kVp, 250 mAs, 0.993 pitch, and 0.75 seconds rotation time with 2 mm slice thickness at 2 mm intervals 70 seconds after the intravenous administration of contrast with axial, sagittal, and coronal reconstructions. This protocol is replicated as closely as possible in our dedicated CECT neck PET/CT.

There are advantages and disadvantages to varying CT and PET parameters in patients with head and neck malignancies, and each patient's unique circumstances should be considered in determining the optimal protocol.




Figure 9 Added value of PET/CT fusion. Axial 18F-FDG PET image (A) through the neck does not demonstrate a definitive focus of abnormal uptake. Axial noncontrast CT through the same level (B) shows no pathologically enlarged lymphadenopathy. Fused axial 18F-FDG PET/CT without contrast (C) shows a small left level 2A lymph node with low uptake (yellow arrow), which makes it difficult to determine whether it is an abnormal metastatic node or a reactive lymph node. Fused axial 18F-FDG PET/CT with iodinated contrast in the same patient more conspicuously demonstrates an abnormal, round, mildly hypermetabolic left level 2A lymph node metastasis (white arrow). (Color version of figure is available online.)

PET/MRI Protocol Considerations

Background

Hybrid imaging with PET/MR presents unique protocol challenges not present in either standalone imaging technique. One of the initial technical challenges to be overcome for clinical PET/MRI was the need for AC of the PET images. PET imaging is uniquely affected by the effects of attenuation, resulting in images with dramatically fewer counts in the middle of the imaged object relative to the periphery. With hybrid PET/CT, the CT can serve as a quickly acquired, accurate attenuation (μ) map used for AC of the images. When CT based AC is utilized, attenuation values of tissues are derived from their measured CT numbers, corrected for the expected attenuation values of 511 keV photons.⁶³ The CT-based method of AC has superseded older emission-based methods that used radioactive sources to generate attenuation maps due to the ubiquity of PET/CT and the significant lengthening of imaging times created by emission methods.

MRI does not provide a similar analogous source of AC due to the fact that MR signal is derived from tissue

hydrogen density and relaxation properties rather than the attenuation of x-rays.⁶⁴ For a set of MR images to serve for AC of the PET data, segmentation maps must be generated, most commonly from a DIXON sequence. This sequence is used to segment tissues into air, lung, fat, and soft tissue. Linear attenuation maps based on the expected attenuation of 511 keV photons in the segmented tissues can then be used to perform AC of the PET images (Fig. 10). A fifth tissue—bone—can be added to the AC algorithm based on atlas data. This helps to avoid the quantitative errors introduced when the attenuation effects of bone are ignored; however, errors can still occur when an individual patient's bone anatomy deviates significantly from atlas data.⁶⁵ In addition, errors can be introduced due to incorrect tissue segmentation at air tissue interfaces—for example in the paranasal sinuses (Fig. 11).⁶⁶ There is continued disagreement among experts regarding whether the quantitative errors introduced by MR based attenuation are clinically significant.⁶⁷

The additional sequences required for accurate AC of the images and the time-consuming nature of MRI relative to CT place a critical emphasis on the need to develop an efficient




Figure 10 Tissue segmentation map. Coronal MR tissue segmentation map derived from DIXON MR sequence and Atlas data. This map is used for attenuation correction of the PET data. Note that the spine, pelvis, and femora are included in the segmentation map, however, the upper extremity bones are not segmented in the software algorithm.

set of PET/MRI protocols. It is therefore necessary to be judicious with the sequences selected for imaging.

PET/MRI Protocol Structure and Oncologic Protocols in the Head and Neck

A general PET/MRI protocol structure is as follows. After an equilibration period (typically at least 45 minutes), the patient lies in the PET/MRI gantry (Fig. 12). Once localizer




Figure 11 Tissue segmentation map of the face with incorrect segmentation. Coronal MR tissue segmentation map of the skull shows portions of the frontal bone, orbits, and paranasal sinuses to be incorrectly segmented as soft tissue. The air-filled paranasal sinuses are also mostly incorrectly segmented.



Figure 12 PET/MR Unit. Siemens (Siemens Healthineers AG, Munich, Germany) Biograph MMR PET/MR scanner is shown. The gantry houses an integrated PET detector ring which allows concurrent PET and MR sequence acquisition.

MRI sequences are obtained, PET and MRI data are obtained concurrently in sequential PET “beds.” While PET data are acquired—typically 2-5 minutes of PET data at each bed—basic MRI sequences are obtained in the anatomical regions of each PET bed. These consist of localizer, AC MR sequences (typically DIXON), and a wide field of view sequence designed to depict the relevant anatomy (we use coronal T2 weighted single shot fast spin echo images). These MRI sequences are designed to fit within the time frame of PET imaging, and typically occur over a time frame of 15-20 minutes. Once the whole-body PET/MR imaging has been completed, dedicated MRI of the primary region of interest is acquired with the imaging protocol tailored to help answer the current clinical question, usually with a slightly abbreviated list of sequences to prevent an excessively long duration of imaging. During this period, an additional longer PET acquisition is acquired, resulting in a set of PET images with higher image quality made possible by the longer duration of imaging.

A typical head and neck oncologic PET/MR protocol at our institution begins with skull to thigh PET/MR imaging beginning at the top of the skull and proceeding to the mid thighs. This is then followed by a dedicated neck or brain MRI tailored to answer the relevant clinical questions (eg, presence of perineural tumor spread, intracranial extension of tumor, etc). For example, a neck PET/MRI (Table 1) to evaluate a malignancy of the suprahyoid neck would consist of axial and coronal T1 nonfat saturation fast spin echo and T2 fat saturation fast spin echo images. This would be followed by DWI in the area of clinical concern. After gadolinium administration, a spoiled 3-dimensional gradient echo sequence (VIBE on Siemens scanners, LAVA on GE) is obtained, and axial and coronal images are generated. PET acquisition continues throughout most of the duration of the neck sequences. This is then followed by axial breath-hold spoiled gradient echo sequences at midinspiration at each of the PET bed regions. We have found that these

Table 1 Sample Oncologic Neck Soft Tissue PET/MRI Protocol

Region Scanned	MRI Sequence	Plane	Imaging Time	Concurrent PET Acquisition
Skull vertex to mid thighs	DIXON	Axial	13 s (each bed)	15 min (3 min per bed)
Skull vertex to mid thighs	HASTE	Coronal	2 min	
Neck	T1 TSE 3 mm	Axial	3 min	Continuous PET acquisition, first 10 min of imaging sent to PACS
Neck	T1 TSE 3 mm	Coronal	2 min	
Neck	T2 FS TSE 3 mm	Axial	3 min	
Neck	T2 FS TSE 3 mm	Coronal	5 min	
Neck	T1 TSE FS + C 3 mm	Axial	5 min	
Neck	T1 TSE FS + C 3 mm	Coronal	3 min	
Skull vertex to mid thighs	VIBE	Axial	17 s (per body region)	None

Table 2 Sample Oncologic Facial PET/MRI Protocol

Region Scanned	MRI Sequence	Plane	Imaging Time	Concurrent PET Acquisition
Skull vertex to mid thighs	DIXON	Axial	13 s (each bed)	15 min (3 min per bed)
Skull vertex to mid thighs	HASTE	Coronal	2 min	
Face	T1 TSE 3 mm	Axial	3 min	Continuous PET acquisition, first 10 min of imaging sent to PACS
Face	T1 TSE 3 mm	Coronal	2 min	
Face	T2 FS TSE 3 mm	Axial	3 min	
Face	T2 FS TSE 3 mm	Coronal	5 min	
Face	T1 TSE FS + C 3 mm	Axial	5 min	
Face	T1 TSE FS + C 3 mm	Coronal	3 min	
Face	DWI	Axial	2 min	
Skull vertex to mid thighs	VIBE	Axial	17 s (per body region)	None

sequences only minimally lengthen the examination and facilitate comparison with previous and future PET/CT imaging in the axial plane.

A facial oncologic PET/MR protocol (Table 2) consists of the above sequences centered on the facial region, though a 3-dimensional FLAIR sequence of the brain may be added if there is concern for intracranial involvement. A general orbital PET/MRI protocol consists of axial and coronal T1 FSE, coronal T2 FSE fat-saturated, 3-dimensional T2 SPACE of the orbits, and axial and coronal gadolinium enhanced T1 FS FSE sequences. An oncologic brain protocol (Table 3) would consist of 3D T1 high resolution spoiled gradient echo sequence (MPRAGE on Siemens scanners, SPGR on

GE scanners), Axial FLAIR, Axial T2 FSE FS, Axial DWI, and contrast-enhanced 3D MPRAGE or SPGR sequences.

It should be noted that at our institution, all head and neck PET/MRIs are protocolled with input of both nuclear medicine and head and neck radiologists, who contribute expertise to a patient-specific protocol that is geared toward answering specific staging-related questions. After completion of imaging, this collaborative approach continues with joint read outs of cases by both neuroradiologists and nuclear radiologists. An example case is depicted in Figs. 13–15. This case study of PET/MR imaging in the initial staging of marginal zone lymphoma highlights the complementary nature of information derived from PET and MRI imaging.

Table 3 Sample Oncologic PET/MRI Brain Protocol Including Optional Orbital Sequences

Region Scanned	MRI Sequence	Plane	Imaging Time	Concurrent PET Acquisition
Skull vertex to mid thighs	DIXON	Axial	13 s (each bed)	15 min (3 min per bed)
Skull vertex to mid thighs	HASTE	Coronal	2 min	
Brain	DWI	Axial	2 min	Continuous PET acquisition, first 10 min of imaging sent to PACS
Brain	T2 FLAIR	Axial	2 min	
Brain	T2 FS TSE	Axial	2 min	
Brain	T1 MPRAGE	3D	5 min	
Orbit (optional)	T1 TSE 3 mm	Axial	3 min	
Orbit (optional)	T2 FS TSE 3 mm	Coronal	3 min	
Orbit (optional)	T1 TSE 3 mm	Coronal	4 min	
Orbit (optional)	T2 SPACE	3D	5 min	
Orbit (optional)	T1 TSE FS + C 3 mm	Axial	5 min	
Orbit (optional)	T1 TSE FS + C 3 mm	Coronal	3 min	
Brain	T1 MPRAGE + C	3D	4 min	
Skull vertex to mid thighs	VIBE	Axial	17 s (per body region)	None




Figure 13 Masticator space mass. Fused axial PET/MRI with T1-weighted postcontrast fat-saturated image demonstrates an avidly-enhancing, FDG avid mass (yellow arrow) of the left infrazygomatic masticator space involving the left medial and lateral pterygoid muscles. (Color version of figure is available online.)



Figure 14 Masticator space mass with perineural spread. Fused coronal PET/MRI with T1-weighted postcontrast fat-saturated image shows an infiltrating, FDG avid masticator space mass (yellow arrow) with enhancing, mildly FDG avid perineural tumor spread along the course of cranial nerve V3 (white arrow). (Color version of figure is available online.)

The joint interpretation of this case resulted in the correct identification of perineural tumor spread by the interpreting neuroradiologist as well as the identification of widespread areas of intramuscular and subcutaneous lymphomatous infiltration by the nuclear radiologist.

In summary, a clinical PET/MRI protocols begins with whole body PET/MRI imaging consisting of abbreviated MRI sequences designed to allow AC and provide basic




Figure 15 Distant disease. Fused axial PET/MR with T1 VIBE post-contrast fat-saturated image shows enhancing, FDG avid intramuscular (white arrows) and subcutaneous (yellow arrow) nodules suggestive of lymphomatous infiltration. (Color version of figure is available online.)

anatomical information in a similar amount of time as clinical PET/CT. This is followed by additional MRI sequences that are designed to answer the clinical question in the area of interest. This strategy typically allows a clinical PET/MRI to be completed in under an hour, allowing the radiologist to answer the question at hand in a time frame acceptable for clinical imaging (Fig. 14).

Conclusion

PET/CT is an established modality in oncologic imaging and is widely used in head and neck imaging. The increasing availability of PET/MRI in the medical landscape raises the question about which modality would be most useful in head and neck oncologic imaging. The early literature for PET/MRI largely shows noninferiority when compared to PET/CT in head and neck imaging, and some studies show advantages of PET/MRI over PET/CT in certain situations, such as the evaluation of nasopharyngeal carcinoma. PET/CT and PET/MRI have separate and unique protocol considerations in regard to head and neck imaging.

References

1. National Comprehensive Cancer Network. Clinical practice guidelines in oncology: Head and neck cancers. https://www.nccn.org/professionals/physician_gls/pdf/head-and-neck.pdf. Published 2018. Accessed February 19, 2019.
2. Torre LA, Bray F, Siegel RL: Global cancer statistics, 2012. *CA Cancer J Clin* 65:87-108, 2015
3. American Cancer Society. ACS cancer-facts-and-figures. <https://www.cancer.org/research/cancer-facts-statistics/all-cancer-facts-figures/cancer-facts-figures-2019.html>. Published 2019. Accessed 2/19/2019.
4. Amin MB, Edge S, Greene F, et al (eds): *AJCC Cancer Staging Manual*, ed 8, Chicago IL: American Joint Committee on Cancer, Springer, 2017
5. Rettig EM, D'Souza G: Epidemiology of head and neck cancer. *Surg Oncol Clin N Am* 24:379-396, 2015
6. Seeburg DP, Baer AH, Aygun N: Imaging of patients with head and neck cancer: from staging to surveillance. *Oral Maxillofac Surg Clin North Am* 30:421-433, 2018
7. Goel R, Moore W, Sumer B, et al: Clinical practice in PET/CT for the management of head and neck squamous cell cancer. *AJR Am J Roentgenol* 209:289-303, 2017

8. Platzek I: (18)F-fluorodeoxyglucose PET/MR imaging in head and neck cancer. *PET Clin* 11:375-386, 2016
9. Brix G, Lechel U, Glattig G, et al: Radiation exposure of patients undergoing whole-body dual-modality 18F-FDG PET/CT examinations. *J Nucl Med* 46:608-613, 2005
10. Chaput A, Robin P, Podeur F, et al: Diagnostic performance of (18) fluorodeoxyglucose positron emission/computed tomography and magnetic resonance imaging in detecting T1-T2 head and neck squamous cell carcinoma. *Laryngoscope* 128:378-385, 2018
11. Subramaniam RM, Truong M, Peller P, et al: Fluorodeoxyglucose-positron-emission tomography imaging of head and neck squamous cell cancer. *AJNR Am J Neuroradiol* 31:598-604, 2010
12. Sun R, Tang X, Yang Y, et al: (18)FDG-PET/CT for the detection of regional nodal metastasis in patients with head and neck cancer: A meta-analysis. *Oral Oncol* 51:314-320, 2015
13. Rohde M, Nielsen AL, Johansen J, et al: Head-to-head comparison of chest X-Ray/head and neck MRI, chest CT/head and neck MRI, and (18)F-FDG PET/CT for detection of distant metastases and synchronous cancer in oral, pharyngeal, and laryngeal cancer. *J Nucl Med* 58:1919-1924, 2017
14. Zhu L, Wang N: 18F-fluorodeoxyglucose positron emission tomography-computed tomography as a diagnostic tool in patients with cervical nodal metastases of unknown primary site: A meta-analysis. *Surg Oncol* 22:190-194, 2013
15. Keraliya AR, Krajewski KM, Braschi-Amirfarzan M, et al: Extracutaneous melanomas: A primer for the radiologist. *Insights Imaging* 6:707-717, 2015
16. Chang AE, Karnell LH, Menck HR: The National Cancer Data Base report on cutaneous and noncutaneous melanoma: A summary of 84,836 cases from the past decade. The American College of Surgeons Commission on Cancer and the American Cancer Society. *Cancer* 83:1664-1678, 1998
17. O'Regan K, Breen M, Ramaiya N, et al: Metastatic mucosal melanoma: Imaging patterns of metastasis and recurrence. *Cancer Imaging* 13:626-632, 2013
18. Park MJ, Oh JS, Roh JL, et al: 18F-FDG PET/CT versus contrast-enhanced CT for staging and prognostic prediction in patients with salivary gland carcinomas. *Clin Nucl Med* 42:e149-e156, 2017
19. Gupta T, Master Z, Kannan S, et al: Diagnostic performance of post-treatment FDG PET or FDG PET/CT imaging in head and neck cancer: A systematic review and meta-analysis. *Eur J Nucl Med Mol Imaging* 38:2083-2095, 2011
20. Szyszko TA, Cook GJR: PET/CT and PET/MRI in head and neck malignancy. *Clin Radiol* 73:60-69, 2018
21. Mehanna H, Wong WL, McConkey CC, et al: PET-CT surveillance versus neck dissection in advanced head and neck cancer. *N Engl J Med* 374:1444-1454, 2016
22. Lowe VJ, Duan F, Subramaniam RM, et al: Multicenter trial of [(18)F] fluorodeoxyglucose positron emission tomography/computed tomography staging of head and neck cancer and negative predictive value and surgical impact in the N0 Neck: results from ACRIN 6685. *J Clin Oncol* 2019;Jco1801182
23. Zhang B, Li X, Lu X: Standardized uptake value is of prognostic value for outcome in head and neck squamous cell carcinoma. *Acta Otolaryngol* 130:756-762, 2010
24. Wang L, Bai J, Duan P: Prognostic value of 18F-FDG PET/CT functional parameters in patients with head and neck cancer: A meta-analysis. *Nucl Med Commun* 40(4):361-369, 2019
25. Nakamoto Y, Tamai K, Saga T, et al: Clinical value of image fusion from MR and PET in patients with head and neck cancer. *Mol Imaging Biol* 11:46-53, 2009
26. Huang SH, Chien CY, Lin WC, et al: A comparative study of fused FDG PET/MRI, PET/CT, MRI, and CT imaging for assessing surrounding tissue invasion of advanced buccal squamous cell carcinoma. *Clin Nucl Med* 36:518-525, 2011
27. Kanda T, Kitajima K, Suenaga Y, et al: Value of retrospective image fusion of (1)(8)F-FDG PET and MRI for preoperative staging of head and neck cancer: comparison with PET/CT and contrast-enhanced neck MRI. *Eur J Radiol* 82:2005-2010, 2013
28. Boss A, Stegger L, Bisdas S, et al: Feasibility of simultaneous PET/MR imaging in the head and upper neck area. *Eur Radiol* 21:1439-1446, 2011
29. Platzek I, Beuthien-Baumann B, Schneider M, et al: PET/MRI in head and neck cancer: Initial experience. *Eur J Nucl Med Mol Imaging* 40:6-11, 2013
30. Varoquaux A, Rager O, Poncet A, et al: Detection and quantification of focal uptake in head and neck tumours: (18)F-FDG PET/MR versus PET/CT. *Eur J Nucl Med Mol Imaging* 41:462-475, 2014
31. Covello M, Cavaliere C, Aiello M, et al: Simultaneous PET/MR head-neck cancer imaging: Preliminary clinical experience and multiparametric evaluation. *Eur J Radiol* 84:1269-1276, 2015
32. Kuhn FP, Hullner M, Mader CE, et al: Contrast-enhanced PET/MR imaging versus contrast-enhanced PET/CT in head and neck cancer: how much MR information is needed? *J Nucl Med* 55:551-558, 2014
33. Platzek I, Beuthien-Baumann B, Schneider M, et al: FDG PET/MR for lymph node staging in head and neck cancer. *Eur J Radiol* 83:1163-1168, 2014
34. Kubiessa K, Purz S, Gawlitza M, et al: Initial clinical results of simultaneous 18F-FDG PET/MRI in comparison to 18F-FDG PET/CT in patients with head and neck cancer. *Eur J Nucl Med Mol Imaging* 41:639-648, 2014
35. Partovi S, Kohan A, Vercher-Conejero JL, et al: Qualitative and quantitative performance of (1)(8)F-FDG-PET/MRI versus (1)(8)F-FDG-PET/CT in patients with head and neck cancer. *AJNR Am J Neuroradiol* 35:1970-1975, 2014
36. Cavaliere C, Romeo V, Aiello M, et al: Multiparametric evaluation by simultaneous PET-MRI examination in patients with histologically proven laryngeal cancer. *Eur J Radiol* 88:47-55, 2017
37. Becker M, Zbaren P, Casselman JW, et al: Neoplastic invasion of laryngeal cartilage: Reassessment of criteria for diagnosis at MR imaging. *Radiology* 249:551-559, 2008
38. Chan SC, Yeh CH, Yen TC, et al: Clinical utility of simultaneous whole-body (18)F-FDG PET/MRI as a single-step imaging modality in the staging of primary nasopharyngeal carcinoma. *Eur J Nucl Med Mol Imaging* 45:1297-1308, 2018
39. Queiroz MA, Hullner M, Kuhn F, et al: PET/MRI and PET/CT in follow-up of head and neck cancer patients. *Eur J Nucl Med Mol Imaging* 41:1066-1075, 2014
40. Schaarschmidt BM, Gomez B, Buchbender C, et al: Is integrated 18F-FDG PET/MRI superior to 18F-FDG PET/CT in the differentiation of incidental tracer uptake in the head and neck area? *Diagn Interv Radiol* 23:127-132, 2017
41. Schaarschmidt BM, Heusch P, Buchbender C, et al: Locoregional tumour evaluation of squamous cell carcinoma in the head and neck area: A comparison between MRI, PET/CT and integrated PET/MRI. *Eur J Nucl Med Mol Imaging* 43:92-102, 2016
42. Becker M, Varoquaux AD, Combescure C, et al: Local recurrence of squamous cell carcinoma of the head and neck after radio(chemo)therapy: Diagnostic performance of FDG-PET/MRI with diffusion-weighted sequences. *Eur Radiol* 28:651-663, 2018
43. Queiroz MA, Hullner M, Kuhn F, et al: Use of diffusion-weighted imaging (DWI) in PET/MRI for head and neck cancer evaluation. *Eur J Nucl Med Mol Imaging* 41:2212-2221, 2014
44. Rasmussen JH, Norgaard M, Hansen AE, et al: Feasibility of multiparametric imaging with PET/MR in head and neck squamous cell carcinoma. *J Nucl Med* 58:69-74, 2017
45. Ruhlmann V, Ruhlmann M, Bellendorf A, et al: Hybrid imaging for detection of carcinoma of unknown primary: A preliminary comparison trial of whole-body PET/MRI versus PET/CT. *Eur J Radiol* 85:1941-1947, 2016
46. Badger D, Aygun N: Imaging of perineural spread in head and neck cancer. *Radiol Clin North Am* 55:139-149, 2017
47. Sekine T, de Galiza Barbosa F, Kuhn FP, et al: PET+MR versus PET/CT in the initial staging of head and neck cancer, using a trimodality PET/CT+MR system. *Clin Imaging* 42:232-239, 2017
48. Abdel Razek AAK, Mukherji SK: State-of-the-art imaging of salivary gland tumors. *NeuroImaging Clin N Am* 28:303-317, 2018
49. Tailor TD, Gupta D, Dalley RW, et al: Orbital neoplasms in adults: Clinical, radiologic, and pathologic review. *Radiographics* 33:1739-1758, 2013
50. Go JL, Rajamohan AG: Imaging of the sella and parasellar region. *Radiol Clin North Am* 55:83-101, 2017

51. Akin EA, Torigian DA, Colletti PM, et al: Optimizing oncologic FDG-PET/CT scans to decrease radiation exposure. 2017. <https://www.imagewisely.org/-/media/Image-Wisely/Files/Nuclear-Medicine/Optimizing-Oncologic-FDG-PET-CT-Scans-to-Decrease-Radiation-Exposure.pdf?la=en>. Accessed 3/25/2019.
52. Kuehl H, Veit P, Rosenbaum SJ, et al: Can PET/CT replace separate diagnostic CT for cancer imaging? Optimizing CT protocols for imaging cancers of the chest and abdomen. *J Nucl Med* 48(Suppl 1):45s-57s, 2007
53. Pfannenbergl AC, Aschoff P, Brechtel K, et al: Value of contrast-enhanced multiphase CT in combined PET/CT protocols for oncological imaging. *Br J Radiol* 80:437-445, 2007
54. Pfannenbergl AC, Aschoff P, Brechtel K, et al: Low dose non-enhanced CT versus standard dose contrast-enhanced CT in combined PET/CT protocols for staging and therapy planning in non-small cell lung cancer. *Eur J Nucl Med Mol Imaging* 34:36-44, 2007
55. Tateishi U, Maeda T, Morimoto T, et al: Non-enhanced CT versus contrast-enhanced CT in integrated PET/CT studies for nodal staging of rectal cancer. *Eur J Nucl Med Mol Imaging* 34:1627-1634, 2007
56. Hall LT, Struck AF, Guglielmo CG, et al: Does the use of IV contrast enhanced CT for attenuation correction affect clinical interpretation of head and neck PET/CT. *Open Nucl Med J* 3:12-18, 2011
57. Rodrigues RS, Bozza FA, Christian PE, et al: Comparison of whole-body PET/CT, dedicated high-resolution head and neck PET/CT, and contrast-enhanced CT in preoperative staging of clinically M0 squamous cell carcinoma of the head and neck. *J Nucl Med* 50:1205-1213, 2009
58. Berthelsen AK, Holm S, Loft A, et al: PET/CT with intravenous contrast can be used for PET attenuation correction in cancer patients. *Eur J Nucl Med Mol Imaging* 32:1167-1175, 2005
59. Yau YY, Chan WS, Tam YM, et al: Application of intravenous contrast in PET/CT: Does it really introduce significant attenuation correction error? *J Nucl Med* 46:283-291, 2005
60. Ciappuccini R, Aide N, Blanchard D, et al: Incremental value of a dedicated head and neck acquisition during 18F-FDG PET/CT in patients with differentiated thyroid cancer. *PLoS One* 11:e0162482, 2016
61. Yamamoto Y, Wong TZ, Turkington TG, et al: Head and neck cancer: Dedicated FDG PET/CT protocol for detection—phantom and initial clinical studies. *Radiology* 244:263-272, 2007
62. Di Martino E, Nowak B, Krombach GA, et al: [Results of pretherapeutic lymph node diagnosis in head and neck tumors. Clinical value of 18-FDG positron emission tomography (PET)]. *Laryngorhinootologie* 79:201-206, 2000
63. Bushberg JT: *The Essential Physics of Medical Imaging*. Lippincott Williams and Wilkins, 2011
64. Berker Y, Franke J, Salomon A, et al: MRI-based attenuation correction for hybrid PET/MRI systems: A 4-class tissue segmentation technique using a combined ultrashort-echo-time/Dixon MRI sequence. *J Nucl Med* 53:796-804, 2012
65. Boellaard R, Quick HH: Current image acquisition options in PET/MR. *Semin Nucl Med* 45:192-200, 2015
66. Dickson JC, O'Meara C, Barnes A: A comparison of CT- and MR-based attenuation correction in neurological PET. *Eur J Nucl Med Mol Imaging* 41:1176-1189, 2014
67. Bailey DL, Pichler BJ, Guckel B, et al: Combined PET/MRI: Multi-modality multi-parametric imaging is here: Summary report of the 4th International Workshop on PET/MR Imaging; February 23-27, 2015, Tubingen, Germany. *Mol Imaging Biol* 17:595-608, 2015