



Original Articles

Pharmacological inhibition of the IKK ϵ /TBK-1 axis potentiates the anti-tumour and anti-metastatic effects of Docetaxel in mouse models of breast cancer

Ryan T. Bishop^a, Silvia Marino^{a,b}, Daniëlle de Ridder^a, Richard J. Allen^a, Diane V. Lefley^a, Andrew H. Sims^c, Ning Wang^a, Penelope D. Ottewell^a, Aymen I. Idris^{a,b,*}

^a Department of Oncology and Metabolism, Medical School, Beech Hill Road, Sheffield, S10 2RX, UK

^b Bone and Cancer Group, Edinburgh Cancer Research Centre, University of Edinburgh, Western General Hospital, Crewe Road, Edinburgh, EH4 2XR, UK

^c Applied Bioinformatics of Cancer, MRC Institute of Genetics and Molecular Medicine, University of Edinburgh, Western General Hospital, Edinburgh, UK



ARTICLE INFO

Keywords:

IKK ϵ
NF κ B
Breast cancer
Combination treatment
Bone metastasis
Osteolysis
Osteoclast
Osteoblast
Bone

ABSTRACT

IKB kinase subunit epsilon (IKK ϵ), a key component of NF κ B and interferon signalling, has been identified as a breast cancer oncogene. Here we report that the IKK ϵ /TBK1 axis plays a role in the initiation and progression of breast cancer osteolytic metastasis. Cancer-specific knockdown of IKK ϵ in the human MDA-MB-231-BT cells and treatment with the verified IKK ϵ /TBK1 inhibitor Amlexanox reduced skeletal tumour growth and osteolysis in mice. In addition, combined administration of Amlexanox with Docetaxel reduced mammary tumour growth of syngeneic 4T1 cells, inhibited metastases and improved survival in mice after removal of the primary tumour. Functional and mechanistic studies in breast cancer cells, osteoclasts and osteoblasts revealed that IKK ϵ inhibition reduces the ability of breast cancer cells to grow, move and enhance osteoclastogenesis by engaging both IRF and NF κ B signalling pathways. Thus, therapeutic targeting of the IKK ϵ /TBK1 axis may be of value in the treatment of advanced triple negative breast cancer.

1. Introduction

The nuclear factor- κ B (NF κ B) signalling is implicated in inflammation and cancer [1–5]. I κ B kinase subunit epsilon (IKK ϵ), a key component of NF κ B signalling, has been identified as a breast cancer oncogene [6]. A number of studies have shown that IKK ϵ is overexpressed in mammary breast tumours and various human breast cancer cell lines [6,7], and its inhibition reduced the development of primary breast cancer in mice [8]. IKK ϵ has also been found to contribute to Tamoxifen resistance in oestrogen receptor positive (ER+) breast cancers [9]. Binding of IKK ϵ to the TANK-binding kinase 1 (TBK1) leads to the activation of NF κ B, interferon regulatory factor (IRF) signalling pathway [10,11] as well as various cancer-driving factors including, protein kinase B (Akt), tumour necrosis factor receptor (TNFR)-associated factor 2 (TRAF2), and inhibition of tumour suppressors such as cyclin-dramatosis tumour suppressor (CYLD) and Forkhead box O 3a (FoxO3a) [9–15]. Although the mechanisms through which IKK ϵ contributes to cellular transformation and primary tumour growth have been established [6], the role of IKK ϵ in breast cancer metastasis is not fully understood.

Breast cancer cells preferentially metastasise to bone and skeletal related events are diagnosed in approximately 70% of advanced breast cancer cases [16]. NF κ B activation enhances excessive osteoclastic activity and causes bone loss, and various studies have implicated IKK ϵ in bone loss associated with oestrogen deficiency, rheumatoid arthritis and Paget's disease [17–20]. However, the role of IKK ϵ in excessive bone loss associated with metastatic breast cancer has not been investigated. Encouraged by a recent report that pharmacological inhibition of IKK ϵ , using the verified IKK ϵ /TBK1 inhibitor Amlexanox [21], inhibited bone loss in a mouse model of osteoporosis [22], we tested the effects of over-expression/knockdown and pharmacological inhibition of IKK ϵ on the initiation, progression and metastasis of breast cancer. Using mouse models of syngeneic and human breast cancer, we showed that cancer-specific knockdown of IKK ϵ in human MDA-MB-231 and administration of the IKK ϵ /TBK1 inhibitor Amlexanox in immune-competent mice inoculated with syngeneic 4T1 cells inhibited skeletal tumour burden and reduced osteolytic bone damage. Administration of Amlexanox alone in mice reduced the growth of the syngeneic 4T1 breast cancer cells in the mammary fat pads and it significantly enhanced the anti-tumour and anti-metastatic properties of

* Corresponding author. Department of Oncology and Metabolism, Medical School, Beech Hill Road, Sheffield, S10 2RX, UK.

E-mail address: aymen.idris@sheffield.ac.uk (A.I. Idris).

<https://doi.org/10.1016/j.canlet.2019.02.032>

Received 21 November 2018; Received in revised form 13 February 2019; Accepted 14 February 2019

0304-3835/© 2019 Elsevier B.V. All rights reserved.

the FDA-approved Docetaxel [23] in mice after removal of the primary tumour. Collectively, these findings suggest that, due to the combined anti-tumour and anti-resorptive effects, IKK ϵ inhibitors, alone and in combination with chemotherapeutic agents, may be of value in the treatment of skeletal and non-skeletal complications associated with advanced triple negative breast cancer.

2. Materials and methods

2.1. Reagents

The IKK ϵ /TBK1 inhibitor Amlexanox was purchased from Abcam (Cambridge, UK). Parental MDA-MB-231, 4T1, Saos-2 osteoblast-like cells and mouse RAW 264.7 macrophages-like cells were purchased from ATCC (Manassas, VA). The human triple negative bone tropic MDA-MB-231 were generated by Dr Nadia Rucci at the University of L'Aquila, Italy. Single cell clones of MDA-MB-231 were expanded and injected into the left ventricle of 4-week-old female BALB/c-nu/nu mice a clone that generated bone only lesions with significant osteolysis in 8-weeks was used [24–26]. Murine 4T1-Luc2 cells were a kind gift from Dr. Munita Muthana (Sheffield, UK) and were transfected to express Luciferase. Tissue culture medium (DMEM and alpha-MEM) was obtained from ThermoFisher (Leicestershire, UK). All primary antibodies were obtained from Cell Signalling Technology (MA, USA) excluding rabbit anti-actin, which was purchased from Sigma-Aldrich (Dorset, UK). Mouse macrophage colony stimulating factor (M-CSF) was obtained from R&D Systems (Abingdon, UK) and receptor activator of NF κ B ligand (RANKL) was a gift from Patrick Mollat (Galapagos SASU, France).

2.2. Cell lines and conditioned medium

Human MDA-MB-231 breast cancer cells, Saos-2 osteoblast-like cells, mouse 4T1 breast cancer and RAW.267 macrophage-like cells were cultured in a standard D-MEM supplemented with 10% fetal calf serum (FCS), glutamine (2 mM), penicillin (100 U/ml), and streptomycin (100 μ g/ml). Human MCF-7, T47D, SK-BR-3, ZR-75-1 were cultured in a standard RPMI supplemented with 10% fetal calf serum (FCS), glutamine (2 mM), penicillin (100 U/ml), and streptomycin (100 μ g/ml).

2.3. Lentiviral infection and generation of stable cell lines

In order to generate IKK ϵ -shRNA expressing lentivirus, HEK 293T packaging cells were transfected with the 5 μ g pLKO.1 lentiviral vector (empty or containing IKK ϵ -shRNA [TRCN-735, -736] (Dharmacon)), psPAX2 vector, pCMV-VSV-G vector and 40 μ l polyethylenimine, Lentiviral supernatant and polybrene (5 μ g/ml; Sigma-Aldrich Dorset, UK) were incubated with the MDA-MB-231-BT cells. Cells were selected with 1 μ g/ml puromycin. To generate IKK ϵ overexpressing cells, a lentiviral delivered CRISPR activation system was used (Santa-Cruz Biotechnology) and cells were transfected according to manufacturer's instructions. The transfected MDA-MB-231 cells were maintained in selection for at least two passages until knockdown/overexpression was confirmed via western blotting.

2.4. Small RNA interference

Human MDA-MB-231-BT cells were transfected with siRNA (5 nM) using Dharmafect 1 reagent (Dharmacon, CO, USA), according to the manufacturer's instructions. A pool of four annealed double-stranded RNA oligonucleotides against TBK1 (M-003788-02-0005), IKK β (M-003503-03-0005), p65 (M-003533-02-0005), IRF3 (M-006875-02-0005) or non-targeting pool no. 1 (D-001206-13-05) siRNAs were purchased from Dharmacon and used according to the manufacturer's instructions. The cells were cultured for 48 h in antibiotic free complete

medium with the transfection reagent. The efficiency of IRF3, TBK1, IKK β and p65 knockdown was assessed by Western blot analysis.

2.5. Assessment of osteoblast differentiation and bone nodule formation

Primary osteoblasts were isolated from the calvarial bones of 2-day-old mice by sequential collagenase digestion as previously described [27]. Osteoblasts were maintained in standard alpha-MEM. For early osteoblast differentiation assays, primary osteoblasts were plated (7×10^3 cells/well). Test compounds and conditioned medium was added the following day. Osteoblast differentiation was measured after 48 h. For bone nodule assays, Saos-2 osteoblast-like cells were seeded into 12-well and treated with L-ascorbic acid (50 μ g/ml), breast cancer cell conditioned medium (20% v/v), compounds at stated doses. The medium was refreshed every 48 h after two treatments, β -glycerophosphate (10 mM) was also added. The cells were cultured for up to 15 days. At the end of the culture period, osteoblast number, differentiation and bone nodule formation were determined by the AlamarBlue assay, alkaline phosphatase (Alk Phos) assay and alizarin red (ALZ) staining [27,28].

2.6. Assessment of osteoclast formation

Osteoclast formation was studied using RANKL and M-CSF generated human and mouse osteoclasts [29] or the macrophage-like cell line RAW.267. For human osteoclasts, CD14⁺ monocytes were isolated by using positive selection beads (Miltenyi Biotec) from peripheral blood mononuclear cells of healthy volunteers following separation using Ficoll-Paque. Following isolation, CD14⁺ monocytes were plated with 25 ng/ml MCSF to generate macrophages. Mouse osteoclasts were generated as previously described [30]. For cultures involving conditioned medium, conditioned medium was prepared as previously described [31] and added to cultures at 10% (v/v) with M-CSF, RANKL and test compounds for up to 9 days with medium refreshed every 48 h. In macrophage-like RAW cell cultures, MCSF was omitted. Osteoclasts were identified using Tartrate-resistant Acid Phosphatase (TRAcP) staining [31].

2.7. Assessment of cell motility

The migration of human MDA-MB-231 cells was assessed by wound healing assay [32]. The confluent cell monolayer was scratched and cellular migration was assessed over time using time-lapse microscopy and T-scratch analysis program. The invasive phenotype of human MDA-MB-231 cells was assayed using the transwell migration assay. Briefly, matrigel (Corning, UK) was diluted to a final concentration of 1.5 mg/ml in serum-free DMEM. A volume of 20 μ l of diluted matrigel was added to 0.8 μ m Transwell inserts (Corning, UK). A cell suspension in serum free DMEM containing 10×10^3 cells and test compounds was added to each insert. Standard DMEM (500 μ l) was used as the chemoattractant. After 72 h, invasive cells on the underside of the membrane were fixed and stained with haematoxylin and eosin. Invasive cells were quantified using ImageJ.

2.8. Western blotting

Western blot analysis was used to detect protein expression and phosphorylation in cultured bone and breast cancer cells. Briefly, cells were seeded in 12 well plates and maintained in standard media until confluency. Prior to stimulation with test agents or vehicle, cells were incubated in serum free tissue culture medium 16 h. Test agents or vehicle were prepared in serum free media and were then added for the desired period of time. Cells were lysed and protein amount was quantified as previously described [33]. Total protein 50–100 μ g was resolved by SDS-PAGE on 12% polyacrylamide SDS gels, transferred onto PVDF membranes (BioRAD, UK) and immunoblotted with

appropriate antibodies according to manufacturer's instructions, using horseradish peroxidase-conjugated secondary antibody (Jackson labs, UK), and then visualised using chemiluminescence (Amersham, UK) on a Biorad ChemiDocMP imaging system.

2.9. Measurement of levels of tumour-derived factors

Level of tumour-derived factors in conditioned medium from human MDA-MB-231 breast cancer cells was determined by Proteome Profiler Human XL Cytokine Array Kit (ARY022, R&D Systems, Abingdon, UK), according to the manufacturer's instructions.

2.10. Animal experiments

All experimental protocols were approved by the Ethics Committee at the University of Edinburgh and Sheffield and were conducted in accordance with the UK Home Office regulations.

2.11. Intra-tibial injection in mice

The effects of cancer-specific inhibition and pharmacological inhibition of IKK ϵ on skeletal tumour growth and osteolysis were investigated through intra-tibial injection of either human MDA-MB-231 or syngeneic 4T1 breast cancer cells. Briefly, 7 four week-old female BALB/c-nu/nu athymic mice were randomly allocated in groups received intra-tibial injection of human MDA-MB-231-BT IKK ϵ -shRNA or IKK ϵ -mock (4×10^3 cells) in the left tibia. In the second intra-tibial experiment, 14 female BALB/c 10-week-old mice received intra-tibial injection of mouse 4T1 breast cancer cells (4×10^3 cells) in the left leg or a sham injection of PBS into their right leg. Animals were sacrificed after 21 days (MDA-MB-231) or 14 days (4T1). Animals were divided into two groups and received intraperitoneal injection of either vehicle (PBS) or Amlexanox (20 mg/kg/daily). Following sacrifice, bones were analyzed by micro-computed tomography (microCT, Skyscan 1172 scanner). Skeletal tumour growth was measured on 2D microCT images using Image J and results were expressed as a percentage of total metaphyseal area (4T1) or on H&E stained sections using the OsteoMeasure software at $10 \times$ magnification.

2.12. Intracardiac injection of murine 4T1 breast cancer cells

We studied the effects of treatment with Amlexanox on the formation of bone metastasis and following intracardiac injection of syngeneic mouse breast cancer cells. Briefly, 8 week-old female BALB/c mice were randomly allocated into groups, anesthetized and single-cell suspension of 1×10^5 4T1-Luc2 cells/100 μ L PBS was injected into the left cardiac ventricle. The following day mice were divided in to groups (8 mice per group) and treated with either daily i.p. injection of vehicle (5% DMSO in PBS) or Amlexanox (35 mg/kg/day). Animals were monitored daily for cachexia (evaluated by body weight) and behaviour. Bone metastases were monitored tri-weekly using the IVIS system and animals were sacrificed by cervical dislocation on day 11.

2.13. Orthotopic injection of murine 4T1 breast cancer cells

We studied the effects of treatment with Amlexanox, Docetaxel or a combination treatment on primary tumour growth of 4T1-Luc2 after orthotopic injection. Briefly, 8 week-old female BALB/c mice were randomly allocated into groups and anesthetized. 4T1-Luc2 (1×10^6 /ml) were re-suspended in 60% PBS, 39% growth factor reduced matrigel (Corning) and 1% trypan blue (Sigma Aldrich). A volume of 100 μ l of the cell suspension was injected into the fat pad of both the left and right inguinal breast (10 mice per group). Animals were monitored daily for cachexia (evaluated by body weight) and behaviour. The size of each tumour was measured externally three times a week by measuring their two axes using a caliper and applying the

formula of the volume of a sphere [$4/3\pi r^3$ where r is the mean of the two axes]. Once tumour volume had reached the maximum volume of 1 cm³, the tumours were resected. The development of metastases was monitored biweekly using the IVIS system. The primary tumours and the lungs, brain, spleen and hind limbs of the animals were removed and *ex vivo* IVIS imaging was performed before formalin fixation.

2.14. Analysis of clinical data

Retrospective analysis of IKK ϵ and TBK1 expression with respect to primary tumour growth and bone relapse was conducted using published cohorts of data. The NKI295 dataset was chosen to assess the association of IKK ϵ bone metastasis, which did not initially probe for TBK1 expression. The CB560 dataset was used for analysis of TBK1 association with bone metastasis. Exhaustive survival analysis was performed by considering all possible cut-points using the surviALL package. *P* values were derived from Wilcoxon test and were two-tailed.

2.15. Statistical analysis

Results were reported as mean \pm standard deviation (SD) unless otherwise stated. A *p*-value value of 0.05 or below was considered statistically significant. All statistical analyses carried out between two groups were students T-test and between three groups analysis of variance (ANOVA) was used. All statistical tests were conducted in GraphPad prism 7.0 for windows. The concentration that produced 50% of response (IC₅₀) was calculated using GraphPad Prism 7 for Windows.

3. Results

3.1. IKK ϵ inhibition reduces breast cancer cell growth *in vitro* and *in vivo*

Previous studies have implicated that IKK ϵ in breast cancer initiation and bone disease [6,17–20]. In our studies, we have observed that IKK ϵ expression is upregulated in highly metastatic human triple-negative breast cancer cell lines including MDA-MB-231, when compared to hormone-sensitive MCF7 and their non-transformed MCF10A control (Fig. S1A). With this in mind and the observation that TBK1 expression is associated with higher risk for bone metastasis (Fig. S2), we tested the effects of the verified IKK ϵ /TBK1 inhibitor Amlexanox on a panel on the viability of a panel of ER+ and ER-breast cancer cell lines with different metastatic abilities to metastasise to the skeleton. As shown in Table 1, Amlexanox (0–100 μ M) significantly reduced the viability of breast cancer cells, however the ER-breast cancer cell cohort had a significantly lower IC₅₀ value. In view of this, we stably knocked down and over-expressed IKK ϵ in the sub-clone MDA-MB-231-BT (Fig. S3), and assessed the growth and metastatic behaviour of these cells *in vitro*

Table 1

Effects of Amlexanox on the viability of a panel of ER+ and ER-breast cancer cell lines with different metastatic abilities to metastasise to the skeleton.

Receptor Status	Breast Cancer Cell line	IC ₅₀ (μ M \pm SD)
ER+	MCF7	159.6 \pm 1.4
	T47D	330.4 \pm 1.6
	ZR-75-1	142.6 \pm 1.3
ER-	SKBR3	98.9 \pm 1.2
	MDA-MB-468	101.9 \pm 1.2
	MDA-MB-231	108.1 \pm 1.1
	MDA-MB-BT1	73.4 \pm 2.5
	4T1-Luc2	30.0 \pm 1.1

Cell viability was measured after 72 h of continuous exposure to Amlexanox by AlamarBlue assay. Calculation of half maximal inhibitory concentrations (IC₅₀) has been performed as described under "Materials and Methods". Values are mean \pm SD and are obtained from 3 independent experiments. BT denotes osteotropic cells.




Fig. 1. IKK ϵ inhibition inhibited breast cancer cell growth *in vitro* and *in vivo*. A-B. *In vitro* cell growth of human MDA-MB-231 (MDA-MB-231-BT) cells overexpressing (IKK ϵ OE) and deficient in (IKK ϵ KD1 and KD2) IKK ϵ or their mock control in the presence and absence of the IKK ϵ /TBK1 inhibitor Amlexanox (10–100 μ M) after 72 h as assessed by AlamarBlue assay. B-C. *In vitro* cell migration of MDA-MB-231-BT described in panel A after 6 h as assessed by wound healing assay. Representative photomicrographs from the experiment described are shown in panel B. D-E. *In vitro* cell invasion of the human MDA-MB-231-BT cells described in panel A after 72 h as assessed by Transwell invasion assay. Representative photomicrographs from the experiment are shown in panel D. F. *In vivo* tumour growth in bone (% tumour area/bone area) after intra-tibial injection of human MDA-MB-231 (left panel) cells deficient in IKK ϵ (IKK ϵ KD2) or their mock control into nu/nu BALB/c mice or mouse 4T1-BT (right panel) cells into immuno-component BALB/c mice treated with the IKK ϵ /TBK1 inhibitor Amlexanox (AmX, 20 mg/kg/daily) (n = 7 group, 14 days). **p < 0.01 from vehicle or mock control.

and *in vivo*. As shown in Fig. 1, over-expression of IKK ϵ in MDA-MB-231-BT enhanced cell migration within 6 h, invasion within 48 h and growth with 72 h, whereas knockdown and pharmacological inhibition of IKK ϵ using the IKK ϵ /TBK1 inhibitor Amlexanox were inhibitory (Fig. 1A–E). *In vivo*, both immuno-deficient mice inoculated with MDA-MB-231-BT cells deficient in IKK ϵ and immuno-competent mice which received the IKK ϵ /TBK1 inhibitor Amlexanox after inoculation with the syngeneic 4T1 cells exhibited significant reduction in the growth of tumour cells in the tibial metaphysis of mice (Fig. 1F).

3.2. IKK ϵ inhibition reduced breast cancer-induced osteolysis *in vivo*

To test if IKK ϵ is implicated in the ability of cells to cause osteolytic bone loss, we assessed bone volume and architecture in mice after intra-tibial injection of breast cancer cells. Detailed micro-computed tomography (microCT) analysis of bone volume at the tibial metaphysis of immuno-deficient mice inoculated with human MDA-MB-231-BT cells deficient in IKK ϵ (Fig. 2A) showed significant increase in total bone volume (BV/TV; Fig. 2B), increased trabecular connectivity (Conn,




Fig. 2. Cancer-specific IKK ϵ inhibition reduced breast cancer-induced osteolysis in nude mice. A. Graphic representation of intra-tibial injection of human MDA-MB-231 deficient in IKK ϵ (IKK ϵ KD2) or their mock control into nu/nu BALB/c mice (n = 5 mock and n = 7 IKK ϵ KD, 21 days). B-D. *In vivo* total trabecular bone volume (B, BV/TV, %), trabecular connectivity (C, Tb.conn, μ m) and porosity (D, Po, %) and thickness (E, Tb. Th, μ m) in the nu/nu BALB/c mice described in panel A. F-G. *In vivo* osteoclastogenesis (F) and osteoblast number (G) in mice from the experiment described in panels A - D. *p < 0.05 **p < 0.01 from vehicle.

Fig. 2C) and reduced porosity (Po, Fig. 2D) yet had no effect on trabecular thickness (Tb.Th, Fig. 2E). Histomorphometric analysis of histological samples from these mice demonstrated that cancer-specific knockdown of IKK ϵ reduced osteoclast number (Fig. 2F), but had no effect on the number of osteoblasts (Fig. 2G). Using the syngeneic model of 4T1-BT (Fig. 3A), we went on to demonstrate that treatment of immuno-component mice with the IKK ϵ /TBK1 inhibitor Amlexanox modestly but significantly increased total bone volume (BV/TV; Fig. 3B) and enhanced trabecular thickness (Fig. 3C), reduced bone porosity (Fig. 3D) connectivity (Fig. 3E). Histomorphometric analysis of histological samples from these mice revealed that administration of the IKK ϵ /TBK1 inhibitor Amlexanox reduced osteoclast number (Fig. 3F) without affecting osteoblast number (Fig. 3G). Similar effects on osteoclast and osteoblast differentiation were obtained in *in vitro* experiments using RANKL-stimulated osteoclasts and calvarial osteoblasts. As shown in Fig. 4, over-expression of IKK ϵ significantly increased the

ability of in the MDA-MB-231-BT to enhance RANKL-induced osteoclast formation, whereas exposure to the IKK ϵ /TBK1 inhibitor Amlexanox inhibited this effect (Fig. 4A, left panels). Consistently, exposure of osteoclast precursors to MDA-MB-231-BT (Fig. S4) or conditioned medium from MDA-MB-231-BT deficient in IKK ϵ or treated with Amlexanox reduced RANKL-induced osteoclast formation *in vitro* (Fig. 4B–C). Exposure of calvarial osteoblasts to conditioned medium from MDA-MB-231-BT deficient in IKK ϵ or treated with Amlexanox had no effects on alkaline phosphatase activity (Fig. 4D) or bone nodule formation (Fig. 4E–F). These results demonstrate that expression of IKK ϵ by breast cancer cells contributes to their osteolytic potential without affecting osteoblast number.

3.3. IKK ϵ regulates MDA-MB-231 behaviour by IRF3/NF κ B activation

IRF3/NF κ B signalling plays a key role in IKK ϵ /TBK1 activity in




Fig. 3. Amlexanox reduced breast cancer-induced osteoclast formation and osteolysis in immuno-competent mice. A. Graphic representation of intra-tibial injection of 4T1-BT cells in immuno-component BALB/c mice treated with vehicle or the IKK ϵ /TBK1 inhibitor Amlexanox (AmX, 20 mg/kg/daily) (n = 7, 12 days). B-E. *In vivo* total trabecular bone volume (B, BV/TV, %) and porosity (D, Po, %) and thickness (E, Tb.Th, μm) in mice from the experiments described in panel A. F-G. *In vivo* osteoclastogenesis (F) and osteoblast number (G) in mice from the experiment described in panels A - D. **p < 0.01 from vehicle.

primary cancer cells [6,34]. To examine whether IRF3/NF κ B signalling contributes to the osteotropic behaviour of breast cancer cells in our models, we knocked down, TBK1, IKK β , p65 or IRF3 in MDA-MB-231-BT cells overexpressing IKK ϵ (Figs. S5A–B), and we studied the metastatic and osteolytic behaviour of these cells *in vitro*. First, we observed that knockdown of TBK1, IKK β , IRF3 or p65 in these cells significantly inhibited IKK ϵ -driven cell migration (Fig. 5A) and reduced the ability of these cells to support RANKL-induced osteoclast formation in bone marrow cell line cultures (Fig. 5B). In light of this, analysis of protein level of human cytokines and chemokines in conditioned medium revealed that IKK ϵ overexpression in the MDA-MB-231-BT cells is associated with upregulation by greater than 2-fold of 11 secreted proteins (Fig. 5C). The identified proteins are common tumour-derived factors that have previously been found to be involved in the regulation of

inflammation, angiogenesis, and tumorigenesis and osteolysis including lipocalin-2 (LPN2) and interleukins -1 (IL-1), -6 (IL-6) and -8 (IL-8) (Fig. 5D). Furthermore, only silencing of p65 reduced the expression of these proteins to below basal levels of MDA-MB-231 mock cells, implicating the NF κ B signalling pathway in the regulation of IKK ϵ /TBK1 driven osteotropic behaviour of the model described. Consistent with this, we also show that knockdown of IKK β or p65 - but not IRF3 or TBK1 - reduced the viability of MDA-MB-231 cells over-expressing IKK ϵ (Fig. S5C).


3.4. Amlexanox reduces breast cancer osteolytic metastasis

Previous research has implicated IKK ϵ in the initiation of primary tumour growth [6,8]. However, the effect of IKK ϵ inhibition on the

initiation and development of breast cancer osteolytic metastasis has not been investigated. As shown in Fig. 6, administration of Amlexanox (35 mg/kg/day) in mice reduced the homing and growth of the syngeneic 4T1 breast cancer cells as evidenced by *in vivo* bioluminescent imaging (Fig. 6A–C).

3.5. Combination of Amlexanox and Docetaxel reduces the growth and metastasis of mammary tumour cells

Previous studies have implicated members of the NFκB pathway in increased resistance to various chemotherapies [9,35–37]. Of all the chemotherapeutic agents tested (Docetaxel, Cyclophosphamide,



(caption on next page)

Fig. 4. IKK ϵ regulates behaviour of breast cancer cells. A-C. *In vitro* osteoclastogenesis of RANKL-stimulated human monocytes (A and B left) or mouse bone marrow cells (B right) after exposure to conditioned medium (20% v/v) from human MDA-MB-231 (MDA-MB-231-BT) cells overexpressing (IKK ϵ OE) and deficient (IKK ϵ KD1 and KD2) in IKK ϵ or their control (MDA-mock) in the presence and absence of the IKK ϵ /TBK1 inhibitor (10 μ M) Amlexanox after 120 h as assessed by TRAcP staining assay. Representative photomicrographs of multinucleated TRAcP positive osteoclasts from the experiment described are shown in panel B left. D. *In vitro* differentiation of primary calvarial osteoblasts after exposure to conditioned medium (20% v/v) from human MDA-MB-231 (MDA-MB-231-BT) cells overexpressing (IKK ϵ OE) and deficient (IKK ϵ KD1 and KD2) in IKK ϵ or their control (MDA-mock) in the presence and absence of the IKK ϵ /TBK1 inhibitor (10 μ M) Amlexanox after 72 h as assessed by alkaline phosphatase (ALP) colourimetric assay. E-F. *In vitro* bone nodule formation in human osteoblast-like cells Saos2 after exposure to conditioned medium (20% v/v) from human MDA-MB-231 (MDA-MB-231-BT) cells in the presence and absence of the IKK ϵ /TBK1 inhibitor (0.01–10 μ M) Amlexanox after 12 days as assessed by Alazarin Red staining assay. Representative photomicrographs of bone nodules from the experiment described are shown in panel F. **p < 0.01 from vehicle or mock control, +p < 0.05 from MDA-MB-231-BT-IKK ϵ OE. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)




Fig. 5. IKK ϵ regulates osteotropic behaviour of MDA-MB-231 by IRF3/NF κ B activation A. (Left) *In vitro* cell migration of human MDA-MB-231 (MDA-MB-231-BT) cells overexpressing (IKK ϵ OE) or the mock control following transfection with targeting IRF3, TBK1, IKK β or p65 or a scrambled control siRNA pools (Scr^{si}) compared to the non-transfected controls after 8 h as assessed by the wound healing assay. B *In vitro* osteoclastogenesis of RANKL-stimulated mouse macrophage-like RAW cells after exposure to conditioned medium (20% v/v) from human MDA-MB-231 (MDA-MB-231-BT) cells overexpressing (IKK ϵ OE) or the mock control following transfection with targeting IRF3, TBK1, IKK β or p65 or a scrambled control siRNA pool (Scr^{si}) compared to the non-transfected controls after 120 h as assessed by TRAcP staining assay. C Catalogue of differentially expressed inflammatory mediators in the conditioned medium from MDA-MB-231 IKK ϵ OE transiently transfected with scrambled siRNA (Scr), IRF3 targeting siRNA (IRF3) or p65 targeting siRNA (p65) compared to MDA-MB-231 Mock cells transiently transfected with scrambled siRNA assessed by Proteome Profiler Human XL Cytokine Array Kit. D Photomicrographs of differentially expressed cytokines assessed by Proteome Profiler Human XL Cytokine Array Kit. *p < 0.05, **p < 0.01 from scrambled siRNA treated Mock control MDA-MB-231-BT, #p < 0.05, ##p < 0.01 ###p < 0.001 from scrambled treated MDA-MB-231-BT-IKK ϵ OE.




Fig. 6. Amlexanox reduces primary and skeletal tumour growth in mice. **A.** Graphic representation of intracardiac injection of parental 4T1 cells (4T1-Luc2) in immuno-component BALB/c mice treated with vehicle (PBS), the IKK ϵ /TBK1 inhibitor Amlexanox (35 mg/kg/daily). **B.** Combined *in vivo* hind limb tumour growth in mice from the experiments described in panel A after 11 days post inoculation of cells. **C.** Representative real-time imaging of bioluminescence of metastatic tumours from the experiment described are shown. **D.** Graphic representation of orthotopic injection of parental 4T1 cells (4T1-Luc2) in immuno-component BALB/c mice treated with vehicle, the IKK ϵ /TBK1 inhibitor Amlexanox (AmX, 35 mg/kg/daily), Docetaxel (Doc, 15 mg/kg/1-time-weekly) or the combination of two agents ($n = 10$ per group, up to 35 days). **E.** *In vivo* mammary tumour growth in mice from the experiments described in panel F. **F.** Real-time imaging of bioluminescence of tumours (bottom) from the experiment described are shown in panels D-E. * $p < 0.05$ from vehicle and + $p < 0.01$ from individual treatment.

Doxorubicin, Rapamycin, 5-Fluorouracil and Tamoxifen; data not shown), only Docetaxel significantly enhanced IKK ϵ expression in parental MDA-MB-231 breast cancer cells (Fig. S6A) and treatment of these cells with Amlexanox enhanced the anti-tumour effects of Docetaxel by up to 50% (Figs. S6B–D, $p < 0.05$). With this in mind, we next sought to determine whether Amlexanox sensitizes the parental and syngeneic 4T1 breast cancer cells to Docetaxel in mammary orthotopic mouse model. As shown in Fig. 6D, combined administration of Amlexanox (35 mg/kg/day) and Docetaxel (15 mg/kg/week) reduced mammary tumour growth (59.7%, $p < 0.001$) in mice inoculated with the 4T1-Luc2 cells. After removal of tumours, Amlexanox and Docetaxel reduced incidence of overt metastases (Fig. 7A–C) and improved survival (Fig. 7D and 1.82 fold, $p < 0.005$) ($p < 0.05$), when compared to vehicle or individual treatments.

4. Discussion

Although previous studies have implicated both the canonical and non-canonical IKK/NF κ B pathways in breast cancer [38,39], IKK ϵ is the only IKK family member that has been deemed a breast cancer oncogene [6]. Here we provide evidence for a previously unknown role of IKK ϵ in the regulation of the initiation and progression of breast cancer osteolytic metastasis. The evidence for this comes from the experiments that showed that cancer-specific knockdown of IKK ϵ in the human triple negative MDA-MB-231 breast cancer cells reduced cell growth and motility *in vitro* and both cancer-specific knockdown and pharmacological inhibition of IKK ϵ suppressed the growth of human MDA-MB-231 and mouse 4T1 cells in the long bones of mice, thereby indicating an inhibitory effects on skeletal tumour burden.

Previous studies reported that the activation of IKK/NF κ B signalling plays a critical role in breast cancer bone metastasis and osteolysis [39–42]. Our current data compliment these studies and demonstrate




Fig. 7. Combination of Amlexanox and Docetaxel reduces overt metastases of parental 4T1 in immunocompetent mice. **A.** Quantification of overt metastasis in mice from the experiments described in Fig. 6 panel D-E. **B** Photomicrographs of real-time bioluminescence imaging following the removal of primary metastasis on day 27 since mammary fat pad inoculation. Representative photomicrographs of metastasis from the experiment described are shown in panel C. **D.** Frequency of metastases in excised organs from the experiments described in panels A – C. **E.** Survival of mice from the experiment described in panels A-C. *p < 0.05 **p < 0.01 from vehicle, #p < 0.05 compared to Amlexanox treated group, \$ p < 0.05 compared to Docetaxel treated group.

that systemic administration of the pharmacological inhibitor of IKKε/TBK1 suppressed the ability of the syngeneic 4T1 breast cancer cells to home, grow and to enhance osteoclast number and osteolysis in mice. These data also expand on previous observations and suggest that the osteoprotective effect associated with IKKε/TBK1 inhibition is mediated not only by the effects on breast cancer cells but also by inhibition of IKKε/TBK1 in bone cells of the tumour microenvironment in the

skeleton. Evidence for this comes from the *in vivo* observations that pharmacological inhibition of IKKε/TBK1 reduced osteoclast number and the *in vitro* data that showed that IKKε/TBK1 in either breast cancer cells or osteoclast precursors significantly reduced the ability to support osteoclast formation.

Pro-inflammatory mediators secreted by breast cancer and bone cells in the tumour microenvironment in the skeleton are essential for

the development of breast cancer related bone disease [1–5,43]. The activation of NF κ B and IRF signal transduction pathways by tumour-derived factors is known to regulate osteoclastogenesis, skeletal tumour growth and bone remodelling and bone loss [9–15,17,44]. Mechanistic studies in human breast cancer cell and osteoclast cultures revealed that the IKK ϵ -driven pro-migratory and pro-osteolytic capacities in our models were abrogated by siRNA-mediated knockdown of TBK1, IRF3 or NF κ B, consistent with the involvement of both the IRF and NF κ B signalling pathways [10,11]. Collectively, these results provide a novel insight into the role of IKK ϵ /TBK1 signalling in the reciprocal interaction between bone and breast cancer cells in the skeleton. When combined with previous studies that reported the role of IKK ϵ in breast cancer initiation and development [8,9] – these results implicate cancer and bone-cell autonomous expression of IKK ϵ in primary and secondary breast cancer to bone.

The IKK ϵ /TBK1 inhibitor Amlexanox enhanced the cytotoxic effects of a panel of chemotherapeutic agents including Docetaxel (Fig. S6 and data not shown), and continuous exposure of human breast cancer cells to Docetaxel enhanced IKK ϵ . Consistent with this we have demonstrated that combined administration of Amlexanox and Docetaxel exerted a synergistic effect characterised by significant reduction of primary tumour growth, overt metastases coupled with approximately a 2-fold improvement in metastasis-free survival when compared to vehicle or either treatment as a single-agent. It is important to note that Amlexanox - as a single agent – also reduced primary tumour growth in these experiments but it had no effect on overt metastases or enhance metastasis-free survival of mice following removal of the primary tumour. Our interpretation of these results is that IKK ϵ /TBK1 inhibition is sufficient to reduce mammary and skeletal tumour growth and osteolysis but failed to reduce general metastasis in the 4T1 models described.

Although the work presented here indicates that IKK ϵ inhibition may be of value in the reduction of breast cancer-induced osteolysis, our studies were restricted to the MDA-MB-231 and 4T1 *in vivo* models. It is also important to note that administration of Amlexanox alone or in combination with Docetaxel had no significant effect on the frequency of bone metastasis (Fig. 5C) and the expression of TBK1 – but not IKK ϵ – is associated with higher risk for bone metastasis in the dataset tested (Fig. S2). Therefore, further studies will be required to determine if inhibition of IKK ϵ , TBK1 or both are effective in reducing the development and progression of bone metastases.

In summary, the results of the present study demonstrate that IKK ϵ plays an essential role in the regulation of primary and skeletal tumour growth, osteolysis and enhanced bone cell activity associated with advanced breast cancer (Fig. S7). These results offer new insight into the role of the NF κ B and IRF signalling pathways in the crosstalk between breast cancer cells and bone cells of the tumour microenvironment, and provide evidence to show that disruption of IKK ϵ /TBK1, alone or in combination with clinically relevant chemotherapies such as Taxanes, may have potential therapeutic efficacy in advanced triple negative breast cancer. The potential benefit of combined administration of IKK ϵ /TBK1 inhibitors and anti-resorptive agents such as bisphosphonates and Denosumab in models of in breast cancer bone metastasis needs to be explored. For that, further *in vivo* studies are warranted.

Conflicts of interest

The authors declare no conflict of interest.

Acknowledgments

These studies were supported in part by Cancer Research UK Development Fund (University of Edinburgh) and funding from Breast Cancer Now - 2014NovPhD390 (University of Sheffield).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.canlet.2019.02.032>.

References

- [1] B. Hoesele, J.A. Schmid, The complexity of NF- κ B signaling in inflammation and cancer, *Mol. Canc.* 2013, pp. 86.
- [2] H. Nakshatri, P. Bhat-Nakshatri, D.A. Martin, R.J. Goulet Jr., G.W. Sledge Jr., Constitutive activation of NF- κ B during progression of breast cancer to hormone-independent growth, *Mol. Cell Biol.* 17 (1997) 3629–3639.
- [3] Y. Zhou, S. Eppenberger-Castori, C. Marx, C. Yau, G.K. Scott, U. Eppenberger, C.C. Benz, Activation of nuclear factor- κ B (NF κ B) identifies a high-risk subset of hormone-dependent breast cancers, *Int. J. Biochem. Cell Biol.* 37 (2005) 1130–1144.
- [4] R.J. Jin, Y. Lho, L. Connelly, Y. Wang, X. Yu, L.S. Jean, T.C. Case, K. Ellwood-Yen, C.L. Sawyers, N.A. Bhowmick, T.S. Blackwell, F.E. Yull, R.J. Matusik, The NF- κ B pathway controls progression of prostate cancer to androgen independent growth, *Cancer Res.* 68 (2008) 6762–6769.
- [5] K. Sakamoto, S. Maeda, Y. Hikiba, H. Nakagawa, Y. Hayakawa, W. Shibata, A. Yanai, K. Ogura, M. Omata, Constitutive NF- κ B activation in colorectal carcinoma plays a key role in angiogenesis, promoting tumor growth, *Clin. Cancer Res. : Official J. Amer. Assoc. Canc. Res.* 15 (2009) 2248–2258.
- [6] J.S. Boehm, J.J. Zhao, J. Yao, S.Y. Kim, R. Firestein, I.F. Dunn, S.K. Sjostrom, L.A. Garraway, S. Weremowicz, A.L. Richardson, H. Greulich, C.J. Stewart, L.A. Mulvey, R.R. Shen, L. Ambrogio, T. Hirozane-Kishikawa, D.E. Hill, M. Vidal, M. Meyerson, J.K. Grenier, G. Hinkle, D.E. Root, T.M. Roberts, E.S. Lander, K. Polyak, W.C. Hahn, Integrative genomic approaches identify IKK ϵ as a breast cancer oncogene, *Cell* 129 (2007) 1065–1079.
- [7] V. Williams, A.A. Grosset, N. Zamorano Cuervo, Y. St-Pierre, M.P. Sylvestre, L. Gaboury, N. Grandvaux, Detection of IKK ϵ by immunohistochemistry in primary breast cancer: association with EGFR expression and absence of lymph node metastasis, *BMC Canc.* 17 (2017).
- [8] T.U. Barbie, G. Alexe, A.R. Aref, S. Li, Z. Zhu, X. Zhang, Y. Imamura, T.C. Thai, Y. Huang, M. Bowden, J. Herndon, T.J. Cohoon, T. Fleming, P. Tamayo, J.P. Mesirov, S. Ogino, K.-K. Wong, M.J. Ellis, W.C. Hahn, D.A. Barbie, W.E. Gillanders, Targeting an IKK ϵ cytokine network impairs triple-negative breast cancer growth, *J. Clin. Invest.* 124 (2014) 5411–5423.
- [9] J.P. Guo, S.K. Shu, N.N. Esposito, D. Coppola, J.M. Koomen, J.Q. Cheng, IKK ϵ phosphorylation of estrogen receptor α Ser-167 and contribution to tamoxifen resistance in breast cancer*, *J. Biol. Chem.* 285 (2010) 3676–3684.
- [10] T.L. Chau, R. Gioia, J.S. Gatot, F. Patrascu, I. Carpentier, J.P. Chapelle, L. O'Neill, R. Beyaert, J. Piette, A. Chariot, Are the IKKs and IKK-related kinases TBK1 and IKK- ϵ similarly activated? *Trends Biochem. Sci.* 33 (2008) 171–180.
- [11] H. Buss, A. Dorrie, M.L. Schmitz, E. Hoffmann, K. Resch, M. Kracht, Constitutive and interleukin-1-inducible phosphorylation of p65 NF- κ B at serine 536 is mediated by multiple protein kinases including I κ B kinase (IKK)- α , IKK- β , IKK- ϵ , TRAF family member-associated (TANK)-binding kinase 1 (TBK1), and an unknown kinase and couples p65 to TATA-binding protein-associated factor II31-mediated interleukin-8 transcription, *J. Biol. Chem.* 279 (2004) 55633–55643.
- [12] J.P. Guo, D. Coppola, J.Q. Cheng, IKK ϵ protein activates Akt independent of phosphatidylinositol 3-kinase/PDK1/mTORC2 and the pleckstrin homology domain to sustain malignant transformation, *J. Biol. Chem.* 286 (2011) 37389–37398.
- [13] R.R. Shen, A.Y. Zhou, E. Kim, E. Lim, H. Habelhah, W.C. Hahn, I κ B kinase ϵ phosphorylates TRAF2 to promote mammary epithelial cell transformation, *Mol. Cell Biol.* 32 (2012) 4756–4768.
- [14] J.E. Hutti, R.R. Shen, D.W. Abbott, A.Y. Zhou, K.M. Spratt, J.M. Asara, W.C. Hahn, L.C. Cantley, Phosphorylation of the tumor suppressor CYLD by the breast cancer oncogene IKK ϵ promotes cell transformation, *Mol. Cell* 34 (2009) 461–472.
- [15] J.P. Guo, W. Tian, S. Shu, Y. Xin, C. Shou, J.Q. Cheng, IKK ϵ phosphorylation and inhibition of FOXO3a: a mechanism of IKK ϵ oncogenic function, *PLoS One* 8 (2013) e63636.
- [16] K. Manders, L.V. van de Poll-Franse, G.J. Creemers, G. Vreugdenhil, M.J. van der Sangen, G.A. Nieuwenhuijzen, R.M. Roumen, A.C. Voogd, Clinical management of women with metastatic breast cancer: a descriptive study according to age group, *BMC Canc.* 6 (2006) 179.
- [17] Q. Sun, B. Sammut, F.M. Wang, N. Kurihara, J.J. Windle, G.D. Roodman, D.L. Galson, TBK1 mediates critical effects of measles virus nucleocapsid protein (MVNP) on pagetic osteoclast formation, *J. Bone Miner. Res. : Official J. Amer. Soc. Bone Miner. Res.* 29 (2014) 90–102.
- [18] D. Hammaker, D.L. Boyle, G.S. Firestein, Synovial innate immune responses: TANK-binding kinase-1 as a potential therapeutic target in rheumatoid arthritis, *Rheumatology (Oxford)* 51 (2012) 610–618.
- [19] M. Corr, D.L. Boyle, L. Ronacher, N. Flores, G.S. Firestein, Synergistic benefit in inflammatory arthritis by targeting I κ B kinase epsilon and interferon beta, *Ann. Rheum. Dis.* 68 (2009) 257–263.
- [20] S.E. Sweeney, L. Mo, G.S. Firestein, Antiviral gene expression in rheumatoid arthritis: role of IKK ϵ and interferon regulatory factor 3, *Arthritis Rheum.* 56 (2007) 743–752.
- [21] S.M. Reilly, S.H. Chiang, S.J. Decker, L. Chang, M. Uhm, M.J. Larsen, J.R. Rubin, J. Mowers, N.M. White, I. Hochberg, M. Downes, R. Yu, C. Liddle, R.M. Evans, D. Oh, P. Li, J.M. Olefsky, A.R. Saltiel, An inhibitor of the protein kinases TBK1/

- IKKe improves obesity-related metabolic dysfunctions, *Nat. Med.* 19 (2013) 313–321.
- [22] Y. Zhang, H. Guan, J. Li, Z. Fang, W. Chen, F. Li, Amlexanox suppresses osteoclastogenesis and prevents ovariectomy-induced bone loss, *Sci. Rep.* 5 (2015) 13575.
- [23] S.B. Zeichner, H. Terawaki, K. Gogineni, A review of systemic treatment in metastatic triple-negative breast cancer, *Breast Canc. (Auckl)* 10 (2016) 25–36.
- [24] N. Rucci, R.E.C. Ficarella, M. Longo, D. Perez, C. Di Giacinto, A. Funari, A. Teti, S. M., In vivo bone metastases, osteoclastogenic ability, and phenotypic characterization of human breast cancer cells, *Bone* 34 (2004) 697–709.
- [25] J.G. Logan, A. Sophocleous, S. Marino, M. Muir, V.G. Brunton, A.I. Idris, Selective tyrosine kinase inhibition of insulin-like growth factor-1 receptor inhibits human and mouse breast cancer-induced bone cell activity, bone remodeling, and osteolysis, *J. Bone Miner. Res. : Official J. Amer. Soc. Bone Miner. Res.* 28 (2013) 1229–1242.
- [26] A.A. Rose, F. Pepin, C. Russo, J.E. Abou Khalil, M. Hallett, P.M. Siegel, Osteoactivin promotes breast cancer metastasis to bone, *Mol. Canc. Res. : MCR* 5 (2007) 1001–1014.
- [27] I.R. Orriss, S.E. Taylor, T.R. Arnett, Rat osteoblast cultures, *Methods Mol. Biol.* 816 (2012) 31–41.
- [28] J.G. Logan, A. Sophocleous, S. Marino, M. Muir, V.G. Brunton, A.I. Idris, Selective tyrosine kinase inhibition of insulin-like growth factor-1 receptor inhibits human and mouse breast cancer-induced bone cell activity, bone remodeling, and osteolysis, *J. Bone Miner. Res.* 28 (2013) 1229–1242.
- [29] S. Marino, J.G. Logan, D. Mellis, M. Capulli, Generation and culture of osteoclasts, *BoneKey Rep.* 3 (2014) 570.
- [30] A.I. Idris, A. Sophocleous, E. Landao-Bassonga, R.J. van't Hof, S.H. Ralston, Regulation of bone mass, osteoclast function, and ovariectomy-induced bone loss by the type 2 cannabinoid receptor, *Endocrinology* 149 (2008) 5619–5626.
- [31] A. Sophocleous, S. Marino, J.G. Logan, P. Mollat, S.H. Ralston, A.I. Idris, Bone cell-autonomous contribution of type 2 cannabinoid receptor to breast cancer-induced osteolysis, *J. Biol. Chem.* 290 (2015).
- [32] A.I. Idris, H. Libouban, H. Nyangoga, E. Landao-Bassonga, D. Chappard, S.H. Ralston, Pharmacologic inhibitors of IkappaB kinase suppress growth and migration of mammary carcinosarcoma cells in vitro and prevent osteolytic bone metastasis in vivo, *Mol. Canc. Ther.* 8 (2009) 2339–2347.
- [33] A. Sophocleous, E. Landao-Bassonga, R.J. Van't Hof, A.I. Idris, S.H. Ralston, The type 2 cannabinoid receptor regulates bone mass and ovariectomy-induced bone loss by affecting osteoblast differentiation and bone formation, *Endocrinology* 152 (2011) 2141–2149.
- [34] C. Korherr, H. Gille, R. Schäfer, K. Koenig-Hoffmann, J. Dixelius, K.A. Eglund, I. Pastan, U. Brinkmann, Identification of proangiogenic genes and pathways by high-throughput functional genomics: TBK1 and the IRF3 pathway, *Proc. Natl. Acad. Sci. U. S. A.* 103 (2006) 4240–4245.
- [35] M.A. Tapia, I. Gonzalez-Navarrete, A. Dalmases, M. Bosch, V. Rodriguez-Fanjul, M. Rolfe, J.S. Ross, J. Mezquita, C. Mezquita, O. Bachs, P. Gascon, F. Rojo, R. Perona, A. Rovira, J. Albanell, Inhibition of the canonical IKK/NF kappa B pathway sensitizes human cancer cells to doxorubicin, *Cell Cycle (Georgetown, Tex.)* 6 (2007) 2284–2292.
- [36] J. Codony-Servat, M. Marin-Aguilera, L. Visa, X. Garcia-Albeniz, E. Pineda, P.L. Fernandez, X. Filella, P. Gascon, B. Mellado, Nuclear factor-kappa B and interleukin-6 related docetaxel resistance in castration-resistant prostate cancer, *Prostate* 73 (2013) 512–521.
- [37] C. Wei, Y. Cao, X. Yang, Z. Zheng, K. Guan, Q. Wang, Y. Tai, Y. Zhang, S. Ma, X. Ge, C. Xu, J. Li, H. Yan, Y. Ling, T. Song, L. Zhu, B. Zhang, Q. Xu, C. Hu, X.W. Bian, X. He, H. Zhong, Elevated expression of TANK-binding kinase 1 enhances tamoxifen resistance in breast cancer, *Proc. Natl. Acad. Sci. U. S. A.* 111 (2014) E601–E610.
- [38] B.K. Park, H. Zhang, Q. Zeng, J. Dai, E.T. Keller, T. Giordano, K. Gu, V. Shah, L. Pei, R.J. Zarbo, L. McCauley, S. Shi, S. Chen, C.Y. Wang, NF-kappaB in breast cancer cells promotes osteolytic bone metastasis by inducing osteoclastogenesis via GM-CSF, *Nat. Med.* 13 (2007) 62–69.
- [39] A.I. Idris, H. Libouban, H. Nyangoga, E. Landao-Bassonga, D. Chappard, S.H. Ralston, Pharmacologic inhibitors of IkappaB kinase suppress growth and migration of mammary carcinosarcoma cells in vitro and prevent osteolytic bone metastasis in vivo, *Mol. Canc. Therapeut.* 8 (2009) 2339–2347.
- [40] P. Peramuhendige, S. Marino, R.T. Bishop, D. de Ridder, A. Khogeer, I. Baldini, M. Capulli, N. Rucci, A.I. Idris, TRAF2 in osteotropic breast cancer cells enhances skeletal tumour growth and promotes osteolysis, *Sci. Rep.* 8 (2018) 39.
- [41] S. Marino, R.T. Bishop, J.G. Logan, P. Mollat, A.I. Idris, Pharmacological evidence for the bone-autonomous contribution of the NFkappaB/beta-catenin axis to breast cancer related osteolysis, *Cancer Lett.* 410 (2017) 180–190.
- [42] D.H. Jones, T. Nakashima, O.H. Sanchez, I. Koziaradzki, S.V. Komarova, I. Sarosi, S. Morony, E. Rubin, R. Sarao, C.V. Hojilla, V. Komnenovic, Y.Y. Kong, M. Schreiber, S.J. Dixon, S.M. Sims, R. Khokha, T. Wada, J.M. Penninger, Regulation of cancer cell migration and bone metastasis by RANKL, *Nature* 440 (2006) 692–696.
- [43] B.S. Parker, J. Rautela, P.J. Hertzog, Antitumour actions of interferons: implications for cancer therapy, *Nat. Rev. Canc.* 16 (2016) 131–144.
- [44] B. Zhao, M. Takami, A. Yamada, X. Wang, T. Koga, X. Hu, T. Tamura, K. Ozato, Y. Choi, L.B. Ivashkiv, H. Takayanagi, R. Kamijo, Interferon regulatory factor-8 regulates bone metabolism by suppressing osteoclastogenesis, *Nat. Med.* 15 (2009) 1066–1071.