



## Original research

# Predicting dynamic balance improvements following 4-weeks of balance training in chronic ankle instability patients



Christopher J. Burcal<sup>a,\*</sup>, Michelle A. Sandrey<sup>b</sup>, Tricia Hubbard-Turner<sup>c</sup>,  
Patrick O. McKeon<sup>d</sup>, Erik A. Wikstrom<sup>e</sup>

<sup>a</sup> University of Nebraska at Omaha, School of Health and Kinesiology, USA

<sup>b</sup> West Virginia University, College of Physical Activity and Sport Sciences, USA

<sup>c</sup> University of North Carolina at Charlotte, Department of Kinesiology, USA

<sup>d</sup> Ithaca College, Department of Exercise and Sport Sciences, USA

<sup>e</sup> University of North Carolina at Chapel Hill, Department of Exercise and Sport Science, USA

## ARTICLE INFO

## Article history:

Received 9 April 2018

Received in revised form 1 November 2018

Accepted 6 November 2018

Available online 15 November 2018

## Keywords:

Clinical prediction rule

Postural control

Therapeutic intervention

Response analysis

## ABSTRACT

**Objectives:** Balance training typically features as a central component of exercise-based rehabilitation programs for patients with lateral ankle sprain and chronic ankle instability (CAI). The purpose of this study was to conduct a responder/non-responder analysis using existing data to identify factors associated with improvements in dynamic balance performance in CAI patients.

**Design:** Secondary data analysis.

**Methods:** Data was used from 73 CAI patients who participated in 6 previous investigations that used the same balance training program. We defined treatment success as a patient exceeding the minimal detectable change score (8.15%) for the posteromedial direction of the Star Excursion Balance Test (SEBT-PM). Baseline measures of participant and injury demographics, patient-reported function, and dynamic balance were entered into a step-wise logistic regression model to determine the best set of predictors of treatment success.

**Results:** Only 28 out of 73 patients (38.4%) demonstrated a successful improvement in SEBT-PM reach after balance training. Of the variables assessed, SEBT-PM reach distance  $\leq 85.18\%$  and self-reported function activities of daily living score  $\leq 92.55\%$  were significant predictors of treatment success ( $p < 0.001$ ). If a patient met both these criteria there was a 70% probability of a successful treatment, indicating a 31.6% increase in the probability of a meaningful balance improvement after completing balance training.

**Conclusions:** Without screening, less than 40% of CAI patients experience a meaningful improvement in SEBT-PM following balance training. Completing a brief pre-treatment assessment of a patient- and clinician-oriented outcome can significantly improve the probability of determining patients with CAI who may improve dynamic balance after balance training.

© 2018 Published by Elsevier Ltd on behalf of Sports Medicine Australia.

## Practical implications

- Not all patients with ankle instability will improve balance after balance training.

- A quick, multifaceted pre-treatment screening can improve the likelihood of a successful improvement in balance after balance training.
- Poor dynamic balance is the strongest individual predictor of a successful improvement in balance.

**Abbreviations:** CAI, chronic ankle instability; SEBT, Star Excursion Balance Test; MDC, minimal detectable change; CPR, clinical prediction rule; FAAM, foot and ankle ability measure; FADI, foot and ankle disability index.

\* Corresponding author.

E-mail addresses: [cburcal@unomaha.edu](mailto:cburcal@unomaha.edu) (C.J. Burcal), [msandrey@mail.wvu.edu](mailto:msandrey@mail.wvu.edu) (M.A. Sandrey), [thubbar1@uncc.edu](mailto:thubbar1@uncc.edu) (T. Hubbard-Turner), [pmckeon@ithaca.edu](mailto:pmckeon@ithaca.edu) (P.O. McKeon), [wikstrom@unc.edu](mailto:wikstrom@unc.edu) (E.A. Wikstrom).

While often considered an innocuous injury, a lateral ankle sprain is one of the most common musculoskeletal injuries, accounting for up to 20% of injuries sustained during organized sports.<sup>1,2</sup> Recent data suggests approximately 40% of individuals who sustain a lateral ankle sprain will develop chronic ankle instability (CAI).<sup>3</sup> CAI is characterized by persistent complaints of the ankle giving way or rolling, as well as recurrent ankle sprains and/or mechanical laxity of the joint.<sup>4</sup> Individuals with CAI often

**Table 1**  
Inclusion and exclusion criteria for included studies and the number of participants from each study used.

Study	Sample Size	Criteria
McKeon et al. <sup>12</sup>	15	Inclusion Exclusion ≥4 answers of “Yes” on All <sup>a</sup> , ≤90% on FADI <sup>b</sup> and FADI-Sport Acute injury (≤6 weeks), lower extremity surgery, history of balance disorders, diabetes, or neuropathy.
Schaefer and Sandrey <sup>13</sup>	9	Inclusion Exclusion ≥4 answers of “Yes” on All No history of ankle sprain or perceived giving way, grade III ankle laxity (anterior drawer, talar tilt, medial subtalar glide), lower extremity surgery, history of balance disorders, diabetes, or neuropathy.
Burcal et al. <sup>14</sup>	12	Inclusion Exclusion ≥4 answers of “Yes” on All, ≤90% on FAAM <sup>c</sup> , ≤80% on FAAM-S <sup>d</sup> Acute injury (≤6 weeks), lower extremity surgery, history of balance disorders, vision problems.
Anguish and Sandrey <sup>20</sup>	11	Inclusion Exclusion ≥4 answers of “Yes” on All No history of lower extremity surgery, history of balance disorders, diabetes, or neuropathy.
Burcal et al. <sup>21</sup>	15	Inclusion Exclusion Score ≥11 on IdFAI <sup>e</sup> Acute injury (≤6 weeks), lower extremity surgery, history of balance disorders, diabetes, neuropathy, concussion.
Gonzales <sup>22</sup>	11	Inclusion Exclusion ≥5 answers of “Yes” on All, ≤90% on FAAM, ≤80% on FAAM-S Acute injury (≤12 weeks), lower extremity surgery, history of concussion, neural impairments, lower limb fractures.

<sup>a</sup> Ankle Instability Instrument.

<sup>b</sup> Foot and Ankle Disability Index.

<sup>c</sup> Foot and Ankle Ability Measure.

<sup>d</sup> Foot and Ankle Ability Measure Sport Scale.

<sup>e</sup> Identification of Functional Ankle Instability.

report lower health-related quality of life<sup>5</sup> and decreased physical activity levels relative to their uninjured peers.<sup>6</sup> Functionally, CAI patients exhibit postural control impairments across a spectrum of tasks including static balance, dynamic balance, and gait.<sup>7,8</sup> These impairments are thought to contribute to the continuum of disability experienced by CAI patients, whereby insufficiencies in postural control limit functional performance, and increase the risk of reinjury, which can further increase the constraints and disability of the patient.<sup>9</sup>

Balance training has been used to improve postural control in CAI patients,<sup>7</sup> and has also been shown to decrease the risk of recurrent ankle sprains.<sup>10</sup> The use of balance training dates back to 1965 when Freeman et al.<sup>11</sup> found that (1) patients who reported recurrent episodes of their ankles giving way had diminished single limb balance and (2) using single limb coordination training significantly reduced the complaints of instability and eliminated the proprioceptive deficit in 11 out of 14 of the original patients. Since then, strong consistent evidence continues to support the use of balance training to improve various patient- (e.g. Foot and Ankle Ability Measure),<sup>12</sup> clinician- (e.g. Star Excursion Balance Test [SEBT]),<sup>13</sup> and laboratory-oriented outcome measures (e.g. time-to-boundary)<sup>7,14</sup> in patients with CAI. However, the literature has focused on whether or not the treatment group's mean improvement from baseline is greater than the control group's mean improvement rather than the proportion of patients who do and do not have favorable outcomes following clinical intervention (i.e. responders and non-responders). Along these lines, 25–50% of CAI patients had a meaningful improvement in balance after receiving 2-weeks of manual therapy treatments while the remaining patients did not.<sup>15</sup> Thus, assuming that all patients within a group will demonstrate a similar level of improvement (i.e. relying solely on group means) may mask a more typical bimodal response and impair clinical decision making.

Clinical prediction rules (CPRs) have the potential to provide clinicians with a practical, evidence-based tool to assist with the identification of patients that are more likely to have a meaningful improvement following a given intervention. CPRs have been developed for treating lower extremity musculoskeletal conditions such as patellofemoral pain with orthotics,<sup>16</sup> acute ankle sprains with manipulative therapy,<sup>17</sup> and CAI with sensory-targeted ankle

rehabilitation strategies (STARS).<sup>15</sup> Considering the consistent recommendations to use balance training in the treatment of lateral ankle sprains and CAI, it is surprising that probability estimates of treatment success following balance training are unknown and that no CPRs yet exist for this treatment.

Therefore the purpose of this study was to conduct a secondary response analysis on data acquired through previous intervention studies which included CAI patients in order to (1) quantify the effectiveness of balance training and (2) identify possible predictors of treatment success following a balance training program. More specifically, we chose the posteromedial reach direction of the SEBT as our primary outcome as it is the most representative of overall SEBT balance performance.<sup>18</sup> Multiple studies have also reported SEBT-PM improvements post balance training<sup>11–13</sup> and evidence suggests that decreased (i.e. worse) SEBT-PM reach scores may double the risk of injury in basketball players.<sup>19</sup> More specifically, we aimed to use CAI patients as an injury model and use patient characteristics (e.g. age, height, weight, etc), injury history (e.g. number of sprains, etc), and baseline assessments of patient- and clinician-oriented outcomes to identify key characteristics in CAI patients who would have the highest probability of meaningful improvements in dynamic balance after completing the 4-week progressive balance training program.

## 2. Methods

We conducted a secondary analysis of data from six previous investigations that all examined the effects of an identical balance training program on those with CAI (Table 1). One investigation was the original randomized controlled trial during the development of this balance training program.<sup>12</sup> Three investigations compared balance training against balance training combined with additional interventions<sup>13,14,20</sup>; therefore only data from the standard balance training groups were used for this investigation. Two of the included investigations are from marked theses and dissertations that utilized the program but are otherwise unpublished or in review.<sup>21,22</sup>

The CAI participants that were randomized to the balance training program were physically active young adults and were tested in research laboratories at three large public universities. Most

**Table 2**  
Univariate comparisons between the successful and unsuccessful groups based on SEBT-PM<sup>a</sup> scores exceeding the minimal detectable change score of 8.15%.

Variable	Successful (n = 28, 38.4%)	Unsuccessful (n = 45, 61.6%)	p Value
Sex: female, n (%)	14 (50%)	28 (62%)	0.317
Age (years)	21.07 ± 3.76	19.93 ± 2.41	0.119
Height (cm)	172.02 ± 11.06	170.64 ± 8.59	0.552
Mass (kg)	73.90 ± 21.22	72.43 ± 12.65	0.711
# of recurrent sprains	3.29 ± 2.87	3.71 ± 4.40	0.651
Self-reported function <sup>b</sup> (%)	83.27 ± 8.97	87.77 ± 7.47	0.023 <sup>g</sup>
Self-reported function: sports <sup>c</sup> (%)	71.03 ± 15.20	73.21 ± 13.66	0.528
SEBT-A <sup>d</sup> (% leg length)	63.87 ± 9.60	75.29 ± 10.67	0.007 <sup>g</sup>
SEBT-PM <sup>a</sup> (% leg length)	75.60 ± 10.61	87.16 ± 13.14	0.001 <sup>g</sup>
SEBT-PL <sup>e</sup> (% leg length)	72.91 ± 17.98	80.40 ± 14.50	0.055 <sup>g</sup>
SEBT-C <sup>f</sup> (% leg length)	72.30 ± 11.53	80.95 ± 11.23	0.002 <sup>g</sup>

<sup>a</sup> Star Excursion Balance Test Posteromedial Reach.

<sup>b</sup> Measured by the Foot and Ankle Ability Measure or Foot and Ankle Disability Index.

<sup>c</sup> Measured by the FAAM-S or Foot and Ankle Disability Index Sport Scale.

<sup>d</sup> Star Excursion Balance Test Anterior Reach.

<sup>e</sup> Star Excursion Balance Test Posterolateral Reach.

<sup>f</sup> Star Excursion Balance Test Composite Score.

<sup>g</sup> Variables are significant between groups of successful and unsuccessful treatment responses  $p < 0.10$  and retained in the regression model.

of these investigations were initiated prior to the International Ankle Consortium's published recommendations for CAI research inclusion criteria,<sup>23–25</sup> however the inclusion criteria for all of the included datasets are comparable to these recommendations. Specific inclusion and exclusion criteria for each study can be seen in Table 1. Participant characteristics can be seen in Table 2.

Each participant included in the analysis completed the 4-week progression-based balance training program developed by McKeon et al.<sup>12</sup> The program consisted of 12, 20-min sessions over a 4-week period that incorporated 5 exercises that progressed in difficulty independent of one another. Full descriptions of the exercises and progression criteria can be seen in Appendix A.

Each investigation collected patient demographics, ankle injury history, dynamic postural control, and patient-reported function at baseline. Patient characteristics common to all studies included demographics such as age, height, mass, and sex. Injury history information common to all studies included the total number of ankle sprains sustained by the individual. Patient-reported function was recorded using either the Foot and Ankle Ability Measure activities of daily living (FAAM) and sports (FAAM-S)<sup>13,14</sup> or Foot and Ankle Disability Index activities of daily living (FADI) and sports (FADI-S)<sup>12</sup> with lower percentages representing more perceived dysfunction during activities of daily living or sports. Dynamic postural control was assessed using the SEBT in all studies. All participants completed the SEBT on the training limb in three directions, anterior (SEBT-A), posteromedial (SEBT-PM), and posterolateral (SEBT-PL). Participants in each investigation were instructed to complete the test by reaching their non-stance limb as far as they could on a flexible tape measure without lifting or moving the stance foot from the center of the testing grid, removing their hands from their hips, or excessive hip rotation. The reach distance of each trial (cm) was normalized to leg length, defined as the distance from the ASIS to the ipsilateral medial malleolus, and the average of 3 trials was used for analysis.<sup>26</sup> The SEBT composite (SEBT-C) score was also calculated, modified from Plisky et al.,<sup>19</sup> where the normalized reach score for all three directions were summed, divided by 3, and multiplied by 100. The SEBT is known to have a learning effect, and it is recommended that participants perform practice trials in order to attribute changes in SEBT performance to the intervention.<sup>19,26,27</sup> Investigators in each of the 6

studies made an effort to accommodate this learning effect by performing between 3 and 4 practice trials in each direction prior to taking the 3-trial average of the test trials.

Participants were dichotomized as having successful or unsuccessful treatments based on the change in the SEBT-PM score from baseline to post-test. A successful treatment was defined as a change in SEBT-PM score that exceeded the minimal detectable change (MDC) for the SEBT-PM (8.15%).<sup>26</sup> The MDC, an estimate of a measure's responsiveness over time, is generated from a non-intervention group over time. Thus, determining treatment success relative to an MDC score is comparable to including a non-intervention arm. For example, McKeon et al.<sup>12</sup> utilized a non-intervention control group, and reported no balance differences from pre- to post-testing. Individual predictor variables were tested for univariate relationship with the reference criterion (e.g. MDC) using independent samples t-tests for continuous variables and  $\chi^2$  tests for categorical variables.<sup>16,17</sup> Variables with a significance level of  $p < 0.10$  were retained as potential predictor variables.<sup>17</sup> This significance level was selected, as previously reported, to reduce the chance that potentially important predictor variables might be overlooked.<sup>15</sup> Potential predictor variables were then entered into a stepwise logistic regression model to determine the most accurate set of variables for prediction of treatment success.<sup>16,17</sup> Variables with a significance level  $p > 0.10$  were removed from the equation to further minimize the risk of excluding variables that could help strengthen the model. Variables retained in the regression model served as the predictors in the CPR.

All potential predictor variables were also submitted to a receiver operator characteristic curve analyses.<sup>16,17</sup> Sensitivity, specificity, positive (+LR) and negative likelihood ratios (–LR), and odds ratios were then calculated for identified cut-off scores for each potential predictor variable. Similarly, diagnostic accuracy and probability of success was calculated for the various combinations of predictor variables. Cut-off scores were identified using Youden's index, or identifying the maximal value of the sum of sensitivity + specificity – 1.<sup>28</sup> This number represents the predictor variable's score with the best diagnostic accuracy,<sup>28</sup> allowing for a CPR to be developed from the cut-off scores with each variable included in the final model.<sup>15</sup> Post-test probability of treatment success was calculated using the +LR and assumed a pre-test probability equal to the proportion of patients who exceeded the SEBT-PM MDC prior to stratification (i.e. prior to being grouped based on potential predictor variables).

### 3. Results

After completing the balance training program, 28 of 73 patients (38.4%) were considered to have a successful treatment with a mean ( $\pm$ SD) SEBT-PM improvement of  $14.8 \pm 4.8\%$  relative to the unsuccessful group (45 of 73 patients [61.6%],  $2.3 \pm 4.3\%$ ). The univariate comparisons of the potential predictor variables between successful and unsuccessful groups sorted by treatment group can be seen in Table 2. Univariate variables that were retained after the step-wise regression for SEBT-PM success and combinations of predictor variables can be seen in Table 3. The variables in the regression analysis included both patient- (Self-reported function) and clinician-oriented (all SEBT directions and SEBT-C) outcomes. As illustrated in Table 3, the SEBT-PM is the individual predictor with the greatest post-test probability (63.5%). In other words, individuals with pre-treatment SEBT-PM reach distances  $\leq 85.2\%$  had a 63.5% chance of a meaningful improvement in SEBT-PM scores. However, the most robust method to predict a meaningful improvement in dynamic postural control following balance training was a SEBT-PM reach distance  $\leq 85.2\%$  combined with a self-reported function score  $\leq 92.5\%$  as this model resulted in a

**Table 3**

Sensitivity, specificity, positive likelihood ratio, post-test probability, and odds ratios for individual predictor variables and predictor variable combinations, when applicable.

	Sensitivity (95% CI)	Specificity (95% CI)	Positive likelihood ratio (95% CI)	Posttest probability	Odds ratio (95% CI)
Individual predictor variables (pre-test probability = 38.4%)					
Self-reported function $\leq 92.5\%^a$	92% (77–98%)	33% (21–48%)	1.39 (1.11–1.75)	46.1%	6.5 (1.36–31.12)
Anterior SEBT reach distance $\leq 78.6\%$	89% (73–96%)	42% (29–57%)	1.55 (1.17–2.05)	48.6%	6.09 (1.6–23.16)
Posteriomedial SEBT reach distance $\leq 85.2\%^a$	82% (64–92%)	71% (57–82%)	2.84 (1.74–4.64)	63.5%	11.32 (3.54, 36.2)
Posteriolateral SEBT reach distance $\leq 78.9\%$	64% (46–79%)	69% (54–80%)	2.07 (1.24–3.46)	55.9%	3.99 (1.47, 10.81)
Composite SEBT reach distance $\leq 81.2\%$	79% (61–90%)	58% (43–71%)	1.86 (1.26–2.76)	53.3%	5.02 (1.71, 14.76)
Predictor variable combinations (pre-test probability = 38.4%)					
Both self-reported function $\leq 92.5\%$ and posteromedial SEBT reach distance $\leq 85.2\%$	79% (61–90%)	80% (66–89%)	3.93 (2.12–7.27)	70.7%	14.67 (4.59, 46.84)

CI: confidence interval, SEBT: Star Excursion Balance Test.

<sup>a</sup> Retained in regression model.

70.7% posttest probability of success. This model also had a positive LR of 3.93, which may indicate a meaningful shift in the posttest odds.

#### 4. Discussion

In this secondary analysis of data from six investigations using identical balance training programs, we identified individual and a combination of variables that could predict which CAI patients are more likely to have a meaningful improvement in dynamic balance following the 4-week intervention. Specifically, we identified that self-reported function during activities of daily living and the SEBT-PM can be used to predict balance improvements following 4-weeks of this particular balance training program.

It is generally accepted that balance improves after balance training in both injured and uninjured populations. This is based on comparing pre- and post-intervention differences in group means. Using this approach, five of the six included studies reported significant improvement in SEBT-PM scores after completing 4-weeks of balance training (except Anguish and Sandrey<sup>20</sup>; Table 1). When combined (n=73), significant pre- to post-intervention mean improvements in all 3 SEBT directions are identified (Hedges'g effect sizes: SEBT-A=0.65, SEBT-PM=0.95, SEBT-PL=1.16). However, our response analysis contradicts these findings as only 38% of participants had a meaningful improvement in dynamic balance (i.e. an SEBT-PM change that exceeded the MDC of 8.15%). Clearly, patients with a drastic improvement in balance scores are skewing the post-intervention means and thus allow for statistical improvements in group means to be noted. Although between-group means and effect sizes are useful for determining treatment efficacy, they may over simplify clinical decision making and downplay the importance of personalized treatment plans based on specific impairments.

The practice of reporting both group means and the percentage of patients responding to a given treatment could certainly improve clinician decision making for patients with musculoskeletal injuries. The individual predictors noted in the current investigation have good to very good diagnostic utility. For instance, using the self-reported function ADL cutoff score of 92.5% there was only an 8% false negative rate (1 – Sensitivity), but there was a 67% false positive rate (1 – Specificity). The higher sensitivity of this measure (92%) suggests that the strength of the self-reported function questionnaire in this model is in identifying patients that are likely to

not improve their balance after balance training. A similar diagnostic relationship is seen with the Ottawa ankle rules, which has a very high sensitivity and allows providers to rule out ankle and foot fractures with a negative test result. However, the test has a poor specificity, meaning that positive tests do not guarantee a fracture is present.<sup>29</sup> In addition, our results suggest that the SEBT-PM is the strongest individual predictor of treatment success. The false negative rate of this measure is moderate, at 18%, however it had the lowest false positive rate out of any of the predictor variables (29%). When participants met this criterion there was a 23% increase in the probability of a successful treatment. Cumulatively, our results support the use of a multi-dimensional evaluation of patient function prior to treatment prescription to improve the likelihood of a successful treatment, as our most profound results are apparent when a patient meets both the self-reported function and SEBT-PM criteria.

The current data would suggest that only 1 in 3 CAI patients would positively respond to a balance training program. Yet our data also indicates that a quick pre-treatment screening may improve this success rate. For example if a clinician had a group of 10 patients with a self-reported function score  $\leq 92.5\%$  and a normalized SEBT-PM reach distance  $\leq 85.2\%$ , ~7 of these patients would have a meaningful improvement in dynamic balance after completing this balance training program. This means that a 15-min screening can increase the probability of treatment success by 30%. Importantly, this combination of predictors had a relatively low false negative (21%) and false positive (20%) rate, suggesting it has an acceptable sensitivity and specificity for identifying CAI patients who respond to balance training. This increase in the probability of a successful treatment for a select group of CAI patients highlights the benefit of an impairment-based model (i.e. treating identified assessments rather than relying on a standard set of treatments for a given pathology).<sup>30</sup>

The results of our response analysis and CPR are consistent with those of Wikstrom and McKeon.<sup>15</sup> Their results demonstrated that ankle joint mobilizations, plantar massage, and calf-stretching resulted in meaningful balance improvements in 25–50% of their sample of CAI patients depending on the intervention in question. However, the presence of specific patient characteristics resulted in post-test probabilities ranging from 75 and 99% among these three common interventions. Similar to the results of the present analysis, Wikstrom and McKeon<sup>15</sup> reported a high likelihood of improving balance (99%) when a patient had a combination of poor

balance and lower scores of self-reported function. This measure of self-reported function was identified as a significant predictor of treatment success for all 3 treatments analyzed.<sup>15</sup> The ability of patient-reported function to identify patients more likely to positively respond to both manual therapies and balance training also highlights the necessity of incorporating patient-reported outcomes into clinical practice.

It is important to note that multiple MDC values exist for the SEBT testing protocol. We chose the more conservative (i.e. higher) value<sup>26</sup> but a lower value (3.36% for SEBT-PM) generated from CAI patients<sup>31</sup> does exist. Using this value, our pre-test probability would be ~65%. Regardless of the MDC value (i.e. measurement error) used, it has not been determined whether or not exceeding the MDC is a *clinically significant* change in a CAI patient's balance. Unfortunately, this is not a new problem as Hertel et al.<sup>18</sup> called for the establishment of a minimal clinically important difference (MCID) for the SEBT in CAI patients over a decade ago. Until an MCID has been established for common clinician-oriented outcomes, we suggest using the standard error of measurement and MDC to complement the traditional between-group effects by reporting the percentage of patients that respond favorably to treatments.

This investigation is not without limitations. While the FADI and FADI Sport are the precursors to the FAAM and FAAM-S, and thus contain nearly identical content and are both reported as a percentage,<sup>32,33</sup> the use of different self-reported function measures must be noted. Further, the best individual predictor of treatment success was the baseline SEBT-PM, which was our criterion for treatment success. Although SEBT-PM reach distance has been shown to be related to overall performance on the SEBT<sup>18</sup> it is possible that since this variable is not independent of our criteria for a successful/non-successful treatment, it could have introduced bias into our analysis. It has been shown that SEBT-A reach performance is related to increased risk of lateral ankle sprain.<sup>19</sup> Furthermore, anterior reach distance during the SEBT has been shown to be related to dorsiflexion range of motion (DFROM).<sup>34</sup> Due to differences among the included investigations, we were unable to control for the influence of DFROM on reach performance in this direction. We could also only use a small number of possible independent variables to serve as potential predictors due to a limited number of consistent variables among the included investigations. While confident in the results, future research should focus on a wider range of measures capable of capturing the full spectrum of impairments present in patients with CAI. Lastly, our investigation is preliminary in nature and has not been validated in a large independent population of CAI patients or other populations (e.g. uninjured controls, ACL-R). Our results indicated that individuals with worse balance at baseline had the greatest improvements, likely due to the fact there was greater room for improving balance. This observation is probably not unique to patients with CAI, and would likely be observed in uninjured individuals as well as patients with other musculoskeletal injuries. Therefore further research should aim to validate the findings of CPRs using treatment outcomes in CAI patients and determine if similar response rates to balance training are present in other pathologic groups (e.g. ACL-R) or uninjured controls who would complete balance training programs as part of an injury prevention program.

## 5. Conclusion

Balance training has been shown to improve dynamic balance and self-reported function at the group level. However, when examining patient responses at the individual level, a bimodal response pattern emerged that affords clinicians the opportunity to identify patients who will most likely benefit from balance training in isolation. Our response analysis showed that both individual

and a combination of patient- and clinician-oriented outcomes can increase the probability that a CAI patient may have a meaningful improvement in dynamic balance. In particular, patients who had a self-reported function score  $\leq 92.5\%$  and a SEBT-PM reach distance  $\leq 85.2\%$  were most likely to have a meaningful improvement in dynamic balance. Because of the clinical implications that a bimodal response has, we recommend that future interventional studies report a response analyses as a clinically-applicable analysis to augment traditional group comparisons.

## Funding sources

The original data analyzed in this manuscript was obtained through support from the NFL Charities, the National Athletic Trainers' Association Research and Education Foundation, and the Mid-Atlantic Athletic Trainers' Association.

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.jsams.2018.11.001>.

## References

- Hootman JM, Dick R, Agel J. Epidemiology of collegiate injuries for 15 sports: summary and recommendations for injury prevention initiatives. *J Athl Train* 2007; 42(2):311–319.
- Swenson DM, Collins CL, Fields SK et al. Epidemiology of U.S. high school sports-related ligamentous ankle injuries, 2005/06–2010/11. *Clin J Sport Med* 2013; 23(3):190–196.
- Doherty C, Bleakley C, Hertel J et al. Recovery from a first-time lateral ankle sprain and the predictors of chronic ankle instability: a prospective cohort analysis. *Am J Sports Med* 2016; 44(4):995–1003.
- Hiller CE, Kilbreath SL, Refshauge KM. Chronic ankle instability: evolution of the model. *J Athl Train* 2011; 46(2):133–141.
- Houston MN, Hoch JM, Hoch MC. Patient-reported outcome measures in individuals with chronic ankle instability: a systematic review. *J Athl Train* 2015; 50(10):1019–1033.
- Hubbard-Turner T, Turner MJ. Physical activity levels in college students with chronic ankle instability. *J Athl Train* 2015; 50(7):742–747.
- Wikstrom EA, Naik S, Lodha N et al. Balance capabilities after lateral ankle trauma and intervention: a meta-analysis. *Med Sci Sports Exerc* 2009; 41(6):1287–1295.
- Hiller CE, Nightingale EJ, Lin CW et al. Characteristics of people with recurrent ankle sprains: a systematic review with meta-analysis. *Br J Sports Med* 2011; 45(8):660–672.
- McKeon PO, Hertel J. The dynamical-systems approach to studying athletic injury. *Athl Ther Today* 2006; 11(1):31–33.
- Schifftan GS, Ross LA, Hahne AJ. The effectiveness of proprioceptive training in preventing ankle sprains in sporting populations: a systematic review and meta-analysis. *J Sci Med Sport* 2015; 18(3):238–244.
- Freeman MA, Dean MR, Hanham IW. The etiology and prevention of functional instability of the foot. *J Bone Joint Surg Br* 1965; 47(4):678–685.
- McKeon PO, Ingersoll CD, Kerrigan DC et al. Balance training improves function and postural control in those with chronic ankle instability. *Med Sci Sports Exerc* 2008; 40(10):1810–1819.
- Schaefer JL, Sandrey MA. Effects of a 4-week dynamic-balance-training program supplemented with graston instrument-assisted soft-tissue mobilization for chronic ankle instability. *J Sport Rehabil* 2012; 21(4):313–326.
- Burcal CJ, Trier AY, Wikstrom EA. Balance training versus balance training with stars in patients with chronic ankle instability: a randomized controlled trial. *J Sport Rehabil* 2017; 26(5):347–357.
- Wikstrom EA, McKeon PO. Predicting balance improvements following STARS treatments in chronic ankle instability participants. *J Sci Med Sport* 2017; 20(4):356–361.
- Vicenzino B, Collins N, Cleland J et al. A clinical prediction rule for identifying patients with patellofemoral pain who are likely to benefit from foot orthoses: a preliminary determination. *Br J Sports Med* 2010; 44:862–866.
- Whitman JM, Cleland JA, Mintken P et al. Predicting short-term response to thrust and nonthrust manipulation and exercise in patients post inversion ankle sprain. *J Orthop Sports Phys Ther* 2009; 39(3):188–200.
- Hertel J, Brahm RA, Hale SA et al. Simplifying the star excursion balance test: analyses of subjects with and without chronic ankle instability. *J Orthop Sport Phys* 2006; 36(3):131–137.
- Plisky PJ, Rauh MJ, Kaminski TW et al. Star excursion balance test as a predictor of lower extremity injury in high school basketball players. *J Orthop Sport Phys* 2006; 36(12):911–919.

20. Anguish B, Sandrey MA. Comparison of the effectiveness of two four-week balance training programs for chronic ankle instability. *J Athl Train* 2018; 53(7):662–671.
21. Burcal CJ, Jeon HJ, Gonzales JM et al. Cortical measures of motor planning and balance training in patients with chronic ankle instability. *J Athl Train* In press.
22. Gonzales JM. *Effects of a 4-week Balance Training and Cognitive Loading Program in Subjects with Chronic Ankle Instability (Vol M.S.)*, Charlotte, University of North Carolina, 2017.
23. Gribble PA, Delahunt E, Bleakley C et al. Selection criteria for patients with chronic ankle instability in controlled research: a position statement of the international ankle consortium. *J Orthop Sports Phys Ther* 2013; 43(8):585–591.
24. Gribble PA, Delahunt E, Bleakley C et al. Selection criteria for patients with chronic ankle instability in controlled research: a position statement of the International Ankle Consortium. *Br J Sports Med* 2014; 48(13):1014–1018.
25. Gribble PA, Delahunt E, Bleakley CM et al. Selection criteria for patients with chronic ankle instability in controlled research: a position statement of the international ankle consortium. *J Athl Train* 2014; 49(1):121–127.
26. Munro AG, Herrington LC. Between-session reliability of the star excursion balance test. *Phys Ther Sport* 2010; 11(4):128–132.
27. Robinson RH, Gribble PA. Support for a reduction in the number of trials needed for the star excursion balance test. *Arch Phys Med Rehabil* 2008; 89(2):364–370.
28. Youden WJ. Index for rating diagnostic tests. *Cancer* 1950; 3(1):32–35.
29. Bachmann LM, Kolb E, Koller MT et al. Accuracy of Ottawa ankle rules to exclude fractures of the ankle and mid-foot: systematic review. *BMJ* 2003; 326(7386):417.
30. Donovan L, Hertel J. A new paradigm for rehabilitation of patients with chronic ankle instability. *Phys Sportsmed* 2012; 40(4):41–51.
31. Hoch MC, Andreatta RD, Mullineaux DR et al. Two-week joint mobilization intervention improves self-reported function, range of motion, and dynamic balance in those with chronic ankle instability. *J Orthop Res* 2012; 30(11):1798–1804.
32. Garcia CR, Martin RL, Drouin JM. Validity of the foot and ankle ability measure in athletes with chronic ankle instability. *J Athl Train* 2008; 43(2):179–183.
33. Hale SA, Hertel J. Reliability and Sensitivity of the foot and ankle disability index in subjects with chronic ankle instability. *J Athl Train* 2005; 40(1):35–40.
34. Gabriner ML, Houston MN, Kirby JL et al. Contributing factors to star excursion balance test performance in individuals with chronic ankle instability. *Gait Posture* 2015; 41(4):912–916.