

## Corrigendum to “Increased serum levels of sFlt-1/PlGF ratio in preeclamptic women with onset at < 32 weeks compared with $\geq 32$ weeks” [Pregnancy Hypertens. 12 (2018) 96–103]


Hirota Suzuki<sup>a,1</sup>, Chikako Hirashima<sup>a,1</sup>, Shiho Nagayama<sup>a</sup>, Kayo Takahashi<sup>a</sup>,  
Tatsuo Yamamoto<sup>b</sup>, Shigeki Matsubara<sup>a</sup>, Akihide Ohkuchi<sup>a,\*</sup>

<sup>a</sup> Department of Obstetrics and Gynecology, Jichi Medical University School of Medicine, Shimotsuke-shi, Tochigi, Japan

<sup>b</sup> Department of Obstetrics and Gynecology, Kasukabe Medical Center, Kasukabe-shi, Saitama, Japan

The authors regret that an error occurred in Fig. 3 of the published article indicated above. The correct Fig. 3 is shown below.

The authors would like to apologise for any inconvenience caused.



DOI of original article: <https://doi.org/10.1016/j.preghy.2018.03.008>

\* Corresponding author at: Department of Obstetrics and Gynecology, Jichi Medical University School of Medicine, 331-1 Yakushiji, Shimotsuke-shi, Tochigi 329-0498, Japan.

E-mail address: [okuchi@jichi.ac.jp](mailto:okuchi@jichi.ac.jp) (A. Ohkuchi).

<sup>1</sup> Both authors equally contributed to this work.

<https://doi.org/10.1016/j.preghy.2019.01.006>

Available online 01 February 2019

2210-7789/ © 2019 International Society for the Study of Hypertension in Pregnancy. Published by Elsevier B.V. All rights reserved.