



Prenatal exposure to antiepileptic drugs and early processing of emotionally relevant sounds

Mari Videman^{a,*}, Susanna Stjerna^b, Valtteri Wikström^c, Taina Nybo^d, Reina Roivainen^d, Sampsa Vanhatalo^b, Minna Huotilainen^{c,e,f}, Eija Gaily^a

^a Department of Pediatric Neurology, New Children's Hospital, University of Helsinki and Helsinki University Hospital, Finland

^b Department of Children's Clinical Neurophysiology, HUS Medical Imaging Center and New Children's Hospital, Helsinki University Hospital and University of Helsinki, Finland

^c Cognitive Brain Research Unit, Department of Psychology and Logopedics, Faculty of Medicine, University of Helsinki, Finland

^d Clinical Neurosciences, University of Helsinki and Helsinki University Hospital, Finland

^e CICERO Learning Network, Faculty of Educational Sciences, University of Helsinki, Finland

^f FinnBrain Birth Cohort Study, Turku Brain and Mind Center, Institute of Clinical Medicine, University of Turku, Finland

ARTICLE INFO

Article history:

Received 10 May 2019

Revised 10 August 2019

Accepted 14 August 2019

Available online 13 September 2019

Keywords:

Evoked potentials

Epilepsy

Newborn

Fetal

Pregnancy

Intrauterine

ABSTRACT

Introduction: Prenatal exposure to antiepileptic drugs (AEDs) is associated with developmental compromises in verbal intelligence and social skills in childhood. Our aim was to evaluate whether a multifeature Mismatch Negativity (MMN) paradigm assessing semantic and emotional components of linguistic and emotional processing would be useful to detect possible alterations in early auditory processing of newborns with prenatal AED exposure.

Material and methods: Data on AED exposure, pregnancy outcome, neuropsychological evaluation of the mothers, information on maternal epilepsy type, and a structured neurological examination of the newborn were collected prospectively. Blinded to AED exposure, we compared a cohort of 36 AED-exposed with 46 control newborns at the age of two weeks by measuring MMN with a multifeature paradigm with six linguistically relevant deviant sounds and three emotionally uttered sounds.

Results: Frontal responses for the emotionally uttered stimulus *Happy* differed significantly in the exposed newborns compared with the control newborns. In addition, responses to sounds with or without emotional component differed in newborns exposed to multiple AEDs compared with control newborns or to newborns exposed to only one AED.

Conclusions: These preliminary findings suggest that prenatal AED exposure may alter early processing of emotionally and linguistically relevant sound information.

© 2019 Elsevier Inc. All rights reserved.

1. Introduction

Approximately 0.3–0.7% of pregnant women in Western countries have a diagnosis of epilepsy. Most of them receive antiepileptic medication during pregnancy [1–3]. The adverse effects of prenatal antiepileptic drug (AED) exposure have been vigorously studied during the last decades. Prenatal exposure to AEDs is associated with both structural and functional teratogenic effects [4–6]. The most prominent effects

have been observed with valproic acid (VPA), which is associated with decreased verbal intelligence [4,7,8], and an elevated risk of autism spectrum disorders [9].

Cortical processing can be measured by event-related potentials (ERPs) extracted from the electroencephalography (EEG) already in neonates. Event-related potential measurement is an economical and non-invasive technique that provides repeatable and quantitative information with high temporal resolution. Auditory processing has been examined by using Mismatch Negativity (MMN) response [10], and these responses can be detected already during neonatal period and also in preterm newborns [11–15]. Altered MMN responses are observed in infants and children suffering from several neural and developmental conditions including dyslexia or autism spectrum disorder [16–22]. More specific alterations in cortical responses have been observed in different age groups. Adults with dyslexia seem to have impaired pitch discrimination [19,23] whereas children with autism spectrum disorders have modified MMN responses for pitch and

Abbreviations: AED, antiepileptic drug; MMN, Mismatch Negativity; ERP, event-related potential; VPA, valproic acid; CBZ, carbamazepine; OXC, oxcarbazepine; LEV, levetiracetam; LTG, lamotrigine; TPM, topiramate; CZP, clonazepam; PWE, pregnant women with epilepsy; EEG, electroencephalography; HNNE, Hammersmith Neonatal Neurological Examination; CA, conceptional age; GA, gestational age.

* Corresponding author at: Child Neurology, New Children's Hospital, P.O.Box 347, FI-00029 HUS, Finland.

E-mail address: mari.videman@hus.fi (M. Videman).

phoneme-category changes [17,18]. Moreover, ERPs measured in term or preterm newborns may already have predictive value for the later development [14,20,21].

In addition to phonetic changes, MMN responses to changes in the emotional components of the sounds have been investigated. Children and adults with autism spectrum disorders are shown to have impaired discrimination of emotional speech prosody [24,25]. Altered auditory processing of emotional information has been detected in young people with conduct disorder symptoms [26]. Orientation to emotionally salient speech may be altered in patients with bipolar disorder and job burnout [27,28]. Recently, a multifeature MMN paradigm to compare semantic and emotional components of linguistic and emotional processing in newborns has been developed [13,29].

Taken together, prior literature suggests that the long-term developmental compromises after prenatal exposure to AEDs are mainly observed in the areas of verbal intelligence and social skills [4,9,30,31]. To test whether early precursors of these developmental compromises could also be detected in newborns after fetal exposure, we used the recently established multifeature MMN paradigm that allows combined assessment of emotional and linguistic processing in the newborn period.

2. Material and methods

2.1. General aspects

The study was conducted at the Helsinki University Hospital in collaboration with Finnish Institute of Occupational Health and University of Helsinki Cognitive Brain Research Unit. The ethics committee of the Helsinki University Hospital has approved the study. All mothers signed a written informed consent during pregnancy. The study protocol followed the Declaration of Helsinki. Recruitment process was prospective and included background information, exposure data, pregnancy outcome data, and mothers' neurocognitive evaluation. The process is represented in detail in a previous publication [32]. The examinations were carried out between April 2010 and May 2014. Examiners were blinded to the exposure status of the newborns.

2.2. Cohort

The original cohort included 56 newborns with prenatal AED exposure and 67 control newborns. Antiepileptic drugs used during pregnancies were carbamazepine (CBZ), clobazam (CLB), clonazepam (CZP), gabapentin (GBP) lamotrigine (LTG), levetiracetam (LEV), oxcarbazepine (OXC), topiramate (TPM), and VPA. Because of newborn-related issues, we were not able to extract ERP epochs from five of the exposed and six of the control newborns. In addition, we missed ERP data because of the lack of reference electrode (eight exposed newborns) or because of inadequate trigger data or artifacts (7 exposed and 15 control newborns). Therefore, a full data set was obtained from 36 exposed and 46 control newborns. Of the exposed newborns, 25 were exposed to only one AED (monotherapy group) and 11 newborns to more than one AED (polytherapy group). Exposure status of the

monotherapy group is shown in Table 1. Polytherapy combinations were as follows (number of newborns in the group): LTG + LEV (1), CBZ + LEV (3), OXC + LEV (1), OXC + GBP (1), LTG + OXC + CZP (1), LTG + LEV + CZP (3), LTG + TPM + CLB (1). In the monotherapy group, mean dosages and mean serum concentrations during the first trimester were as follows: CBZ 733 mg (29 $\mu\text{mol/l}$), LTG 280 mg (7.7 $\mu\text{mol/l}$), OXC 825 mg (27 $\mu\text{mol/l}$), VPA 740 mg (total serum concentration 283 $\mu\text{mol/l}$, free serum concentration 24 $\mu\text{mol/l}$), and LEV 825 mg. Levetiracetam serum levels were not monitored as the monitoring of LEV serum levels is not included in clinical practice in our district.

Of background variables (Table 2), the exposed newborns had lower mean birth weight compared with the unexposed newborns. The difference in Apgar scores was due to one outlier, and vanished if we left this outlier out. Mothers of the AED group tended to have higher educational level and lower performance intelligence quotient (IQ) level than the mothers of the control group. Background information is revealed in detail in Table 2.

Neurological status of the newborns was assessed by Hammersmith Neonatal Neurological Examination (HNNE) [33] at the conceptional age (CA) of 41 to 42 weeks. As observed before when analyzing the entire cohort [32], the exposed newborns of this subcohort had slightly lower limb and axial tone than the control newborns. The mean Compound Optimality Score Tone for exposed newborns of the subcohort was 5.9 and for controls, 7.1 ($p = 0.003$). The mean Total Optimality Score were 23.9 and 25.4 ($p = 0.029$). Neurological findings were reported in detail in our previous publication [32].

2.3. MMN

Electroencephalography signals were collected by using NicOne EEG amplifier at sampling frequency of 250 or 500 Hz and EEG caps with 20–32 sintered Ag/AgCl electrodes positioned according to the International 10–20 standard. For sleep state assessment, we included channels for chin EMG (electromyography), ECG (electrocardiogram), eye movements, and respiratory sensors. The sleep state of trace alternant or quiet sleep was determined visually by using NicoletOne Reader software by consensus agreement (MV and SV) when needed.

Mismatch Negativity recordings were assessed with the multifeature MMN paradigm during either trace alternant or quiet sleep. A standard stimulus (pseudoword /ta-ta/), six types of *linguistically relevant deviant stimuli* (vowel duration, vowel change, intensity changes (± 6 dB), frequency changes (± 25.5 Hz)), and three types of *emotionally uttered stimuli* (happy, angry, sad) were performed as explained in full detail in previous publications from Pakarinen et al. [29] and Kostilainen et al. [13]. The stimuli were presented via a loudspeaker located at a distance of approximately 1 m. The MMN data were collected from the frontal (F3, Fz, F4), central (C3, Cz, and C4), and occipital (Oz) electrodes. The Oz was used as a reference electrode. As the paradigm for emotional stimuli was only introduced after the recruitment had already started, the first 18 recordings lacked the emotional stimuli.

Intervals between 100–200 ms and 300–500 ms were considered descriptive for the brain responses, and the averaged value of the signal within these latency windows was collected. The mean amplitude

Table 1
AED exposure.

	OXC ^a /CBZ ^a N (%)	VPA ^a N (%)	LTG ^a N (%)	LEV ^a N (%)	TPM ^a N (%)	Polytherapy N (%)	All N (%)
All enrolled	10/9 (35)	5 (9)	8 (15)	7 (13)	1 (2)	14 (25)	55 ^b (100)
ERP data available	9/0 (25)	5 (14)	5 (14)	6 (17)	0 (0)	11 (31)	36 (100)

OXC = oxcarbazepine, CBZ = carbamazepine, VPA = valproic acid, LTG = lamotrigine, LEV = levetiracetam, TPM = topiramate.

The standard MMN stimuli were recorded from 36 AED exposed and 46 control newborns. Of these newborns, 18 exposed and 46 control newborns had emotional stimuli.

^a Monotherapy.

^b Exposure status is not available for one newborn.

Table 2
Background information.

	AED (n = 46)	Controls (n = 36)	Sig. (AED vs. controls ^e)
GA ^a (weeks, mean (range, SD))	40.2 (37.9–42.3, ± 1.15)	40.3 (38.4–42.1, ± 1.11)	0.67
CA ^b during ERP (weeks, mean)	42.3 (40.4–44.4, ± 0.94)	42.2 (40.0–43.7, ± 0.86)	0.81
Age during ERP (weeks, mean)	2.1 (0.58–0.40, ± 1.04)	2.0 (0.0–4.85, ± 1.04)	0.65
Enrollment (GA ^a weeks, mean)	7.35 (3–24, ± 3.75)	10.00 (5–15, ± 7.07)	0.36
Educational level of the mother (primary/secondary/tertiary)	34%/57%/9%	64%/30%/7%	0.05
Age of the mother (mean, yrs)	31.5 (24.0–41.0, ± 4.36)	31.7 (21.0–38.0, ± 3.66)	0.82
Smoking during the third trimester (%)	3%	0%	0.53
Neuropsychology of the mothers			
VIQ ^c (mean)	111 (69–137, ± 13)	115 (103–133, ± 9)	0.25 ^f
PIQ ^d (mean)	116 (62–132, ± 13)	123 (102–138, ± 10)	0.05^f
Executive problems (no problems/slight problems)	72%/28%	87%/13%	0.46
Gender (Male %)	64%	69%	0.58
Apgar at 1 min (mean)	9.0 (7–10)	8.5 (2–10)	0.02
Folic acid amount during the 1st trimester (mean, mg)	2.9 (0.3–5.0, ± 1.75)	3.3 (0–8.0, ± 2.29)	0.42
Birth weight (grams)	3480 (2490–4555, ± 465)	3714 (2808–4800, ± 474)	0.03
Hemoglobin of the newborn	167 (127–199, ± 19)	166 (127–214, ± 22)	0.93

^a GA = gestational age (weeks).

^b CA = conceptional age (weeks).

^c Verbal intelligence quotient.

^d Performance intelligence quotient.

^e t-Test, Pearson's Chi-Square test, or Fisher's Exact test. P-values 0.05 or below are marked with bold.

^f After discarding one outlier mother with VIQ 69 and PIQ 62, the p-value for VIQ is 0.42 and for PIQ 0.12 (mean VIQ: AED 112, control 114 and mean PIQ: AED 117, control 122).

values from frontal electrodes were averaged together as electrode area F and from central electrodes as electrode area C. The mean values were calculated for each infant and each stimulus type for both time windows. To obtain standard-subtracted mean amplitudes, amplitude of standard stimulus was subtracted from the mean value in each electrode area (F and C).

Epochs with signal values larger than ± 150 µV at any channel, or response average larger than ± 10 µV at an individual electrode, were considered to be artifactual. Artifactual epochs were rejected, and individual artifactual ERP responses were left out from the averages.

2.4. Statistical analysis

To compare the ERPs of AED exposed and unexposed newborns, the mean standard-subtracted ERP amplitudes of frontal and central signals were computed for the given time windows.

Both latency windows (100–200 ms and 300–500 ms) were tested separately. For group comparisons, we used independent-sample t-test (2-tailed) for continuous variables, and Chi-square test for independence, or if the expected frequency was less than five, Fisher's Exact test, for categorical variables.

Table 3
Standard-subtracted mean amplitudes of AED exposed and control newborns.

Stimulus	AED/100–200 ms µV (min, max, SD)	Control/100–200 ms µV (min, max, SD)	p*	AED/300–500 ms µV (min, max, SD)	Control/300–500 ms µV (min, max, SD)	p*
Emotional variants						
Happy /ta-ta:/						
Frontal	0.72 (− 5.79, 4.74, ± 3.14)	− 0.25 (− 11.25, 10.95, ± 4.69)	0.33	2.08 (− 3.93, 8.32, ± 3.03)	− 0.13 (− 14.93, 9.36, ± 4.35)	0.04
Central	0.51 (− 5.03, 3.67, ± 2.19)	0.21 (− 6.61, 6.88, ± 2.97)	0.65	0.44 (− 6.08, 3.80, ± 2.21)	− 0.24 (− 14.89, 6.90, ± 3.14)	0.32
Sad /ta:-ta:/						
Frontal	− 0.38 (− 7.26, 7.63, ± 4.21)	− 0.59 (− 8.28, 11.45, ± 4.34)	0.86	1.57 (− 6.35, 12.13, ± 4.99)	1.27 (− 11.42, 14.41, ± 5.44)	0.83
Central	0.46 (− 0.39, 7.73, ± 2.76)	0.04 (− 6.05, 7.67, ± 2.63)	0.57	1.01 (− 6.66, 6.18, ± 3.04)	0.30 (− 11.14, 11.50, ± 4.25)	0.45
Angry /ta-ta/						
Frontal	− 0.25 (− 10.05, 7.59, ± 4.49)	− 0.29 (− 7.02, 6.47, ± 3.45)	0.97	0.48, (− 5.16, 4.47, ± 2.77)	1.28 (− 8.55, 21.05, ± 5.01)	0.41
Central	0.45 (− 4.35, 7.08, ± 2.89)	0.15 (− 8.16, 11.26, 3.30)	0.71	0.39 (− 2.43, 7.22, ± 2.14)	1.53 (− 5.11, 18.57, ± 4.07)	0.14
Deviants						
Vowel change /ta-to/						
Frontal	− 0.34 (− 4.00, 3.75, ± 1.88)	0.26 (− 5.33, 7.69, ± 2.28)	0.19	0.03 (− 2.83, 3.12, ± 1.58)	0.34 (− 4.52, 5.91, ± 2.26)	0.47
Central	− 0.24 (− 2.82, 3.03, ± 1.17)	− 0.02 (− 4.93, 4.90, ± 1.88)	0.53	0.27 (− 2.78, 3.98, ± 1.36)	− 0.02 (− 2.34, 3.82, ± 1.28)	0.33
Vowel duration /ta-ta:/						
Frontal	− 0.10 (− 2.42, 5.82, ± 1.71)	0.12 (− 5.36, 7.04, ± 2.58)	0.64	0.07 (− 2.90, 3.73, ± 1.59)	− 0.65 (− 5.01, 3.13, ± 1.90)	0.07
Central	− 0.05 (− 2.62, 3.21, ± 1.36)	− 0.02 (− 4.50, 6.56, ± 2.06)	0.94	0.08 (− 2.18, 2.80, ± 1.15)	− 0.36, (− 4.28, 3.83, ± 1.50)	0.13
Frequency (positive) /ta-ta/						
Frontal	0.22 (− 3.95, 4.81, ± 2.40)	0.83 (− 5.45, 6.21, ± 2.60)	0.27	− 0.11 (− 5.82, 6.21, ± 2.50)	0.12 (− 4.86, 5.61, ± 2.30)	0.67
Central	− 0.21 (− 4.18, 3.30, ± 1.91)	0.45 (− 5.15, 6.07, ± 2.09)	0.13	0.07 (− 2.40, 5.63, ± 1.72)	0.07 (− 5.54, 4.57, ± 2.03)	1.00
Frequency (negative) /ta-ta/						
Frontal	0.33 (− 5.48, 8.40, ± 2.90)	0.29 (− 10.34, 12.20, ± 3.66)	0.95	− 0.18 (− 4.96, 7.59, ± 2.99)	0.15 (− 5.15, 4.58, ± 2.06)	0.57
Central	0.14 (− 3.24, 6.115, ± 1.94)	0.44 (− 7.10, 9.28, ± 2.67)	0.56	− 0.23 (− 3.93, 8.32, ± 3.03)	0.29 (− 3.76, 5.64, ± 1.72)	0.19
Intensity (positive) /ta-ta/						
Frontal	0.30 (− 6.16, 5.58, ± 2.62)	− 0.08 (− 11.35, 8.09, ± 4.05)	0.60	− 0.16 (− 5.94, 5.86, ± 3.07)	− 0.01 (− 5.40, 8.85, ± 2.83)	0.82
Central	− 0.002 (− 5.95, 2.50, ± 1.91)	0.11 (− 9.27, 8.11, ± 3.04)	0.84	0.13 (− 3.10, 4.29, ± 1.93)	0.49, (− 4.44, 7.21, ± 2.13)	0.43
Intensity (negative) /ta-ta/						
Frontal	0.34 (− 5.87, 8.79, ± 3.59)	0.73 (− 4.00, 10.82, ± 3.00)	0.61	1.21 (− 3.84, 6.97, ± 2.33)	0.34 (− 7.10, 6.37, ± 3.02)	0.14
Central	− 0.46 (− 5.14, 3.73, ± 2.13)	0.29 (− 3.54, 5.09, ± 2.06)	0.12	0.47 (− 2.76, 4.20, ± 1.56)	0.09 (− 11.09, 5.61, ± 2.76)	0.43

* p-Values of t-test. Bonferroni correction was applied. P-values below 0.05 are marked with bold.

When evaluating whether gender affected the mean amplitudes, we applied one-way analysis of variance (ANOVA), and when evaluating the effect of age and CA on the brain responses, we applied the analysis of covariance (ANCOVA). In cases with multiple comparisons, we applied Bonferroni correction.

3. Results

3.1. AED group

Standard-subtracted mean amplitudes, both *emotionally uttered stimuli* and *linguistically relevant deviant stimuli*, in the exposed and unexposed newborns are shown in Table 3 and Figs. 1 and 2.

The mean amplitude of the emotional variant *Happy* in the late latency window from the frontal electrode area was found to significantly differ between the groups: the exposed newborns showed positive polarity (2.08 μV , standard deviation (SD): ± 3.03) while the controls showed negative polarity ($-1.30 \mu\text{V}$, SD: ± 4.35 , $p = 0.04$). To evaluate the impact of age and CA on the reaction to the stimulus, we conducted one-way between-groups ANCOVA by using age and CA of the newborn as covariates. Exposure status was the independent variable, and standard-subtracted mean amplitude was the dependent variable. After adjusting for age and CA, there were still no significant differences between AED and control groups for any other stimuli or latency window than emotional variant *Happy* in the late latency window from the frontal electrode area ($p = 0.04$).

3.2. Polytherapy and monotherapy groups

Newborns with exposure to more than just one AED (polytherapy group) had significantly more positive standard-subtracted mean amplitudes than control newborns in the early latency window for emotional variant *Angry* in the frontal area (polytherapy group: 4.26 μV , SD: ± 2.41 ; control group: 0.29 μV , SD: ± 3.45 , $p = 0.01$). The polytherapy group also differed significantly from the control group with regard to early positive frequency deviants in both frontal and in central electrode areas (frontal electrodes: polytherapy group: $-1.02 \mu\text{V}$, SD: ± 0.89 ; control group: 0.83 μV , SD: ± 2.60 , $p < 0.001$, and central electrodes: polytherapy group: $-1.12 \mu\text{V}$, SD: ± 1.97 ; control group 0.45 μV , SD: ± 2.09 , $p = 0.03$). We observed similar statistically significant findings when we compared standard-subtracted mean amplitudes of the polytherapy and monotherapy groups with each other, though monotherapy group as itself did not differ significantly from the control group.

When different monotherapy groups were compared with each other and to the control group, we did not find significant differences in standard-subtracted mean amplitudes (ANOVA).

A summary of the statistically significant findings is outlined in Table 4.

4. Discussion

Our results suggest that prenatal AED exposure may affect early processing of emotionally relevant linguistic information. Earlier studies have shown that auditory-evoked potentials are linked to language functions, and they may specifically predict language impairments [34]. There has been an association between MMN amplitudes and performance in cognitive tests [35], and the intensity of auditory-evoked potentials has been shown to correlate with attention [36]. Atypical processing of the emotional sounds has been observed in patients with

autism spectrum disorders [24] [25]. These together are compatible with the idea that the present findings might reflect an early precursor of developmental challenge caused by AED exposure.

Our study is in line with previous studies showing no major adverse effects on verbal development in association with prenatal OXC, CBZ, LEV, or LTG monotherapy exposure [4,7,8]. Our study was initially motivated by the earlier findings that intrauterine VPA exposure leads to later risk of verbal and social compromise [37]. Patients with epilepsy with VPA medication have shown to have decreased P300 amplitudes when compared with patients with epilepsy with CBZ medication, patients with epilepsy without medication, or healthy controls [38]. Delayed and smaller MMN responses (N270) have been observed in patients with epilepsy compared with the responses of healthy controls [38].

The main limitation of our study was the relatively small number of exposed newborns in each monotherapy group, particularly VPA group (Table 1). Mismatch Negativity data were further limited because of artifact or incomplete electrode settings, and data from emotionally uttered stimuli were limited as the paradigm for emotional stimuli was only introduced after the first 18 recordings of all MMN data. Thus, the number of the newborns exposed to any particular AED was too few to evaluate the impact of an individual medication. We cannot exclude the possibility that individual drugs might have, either adverse or protective, effects on neonatal auditory processing.

There are several strengths in this study. The background information including exposure data was collected prospectively. During all measurements and analyses, the researchers were blinded for the exposure status of the child. Furthermore, MMN recordings were conducted during the same sleeping state (quiet sleep) and during limited time range (conceptual age) thus eliminating the effects of sleeping states and maturation on the quality of the MMN response [39–42].


In our cohort, there was a trend toward lower mean performance IQ in women with epilepsy than in women without epilepsy (Table 2) which could be explained by lower processing speed of the mothers with epilepsy [43]. However, the percentage of the women having either tertiary or secondary education was higher among women with epilepsy. Higher educational level might be due to neuropsychological evaluation included in neurological follow-up thus enabling earlier and/or more intensive educational support or due to higher motivation to show one's capacity despite living a life with a chronic disease. Aspects regarding both strengths and limitations regarding enrollment and background data collection are discussed in more detail in our previous publication [32].

Mismatch Negativity paradigms are applied to several clinical approaches to increase the understanding of disease mechanisms of many neurodevelopmental, neuropsychiatric, and neurological disorders, or to serve as biomarkers for risk of these disorders [44]. In particular, MMN responses have been used to explore the relationship between auditory processing and developmental disorders. It remains seen whether the deviant responses to sad and angry stimuli in frontal areas of the polytherapy group in our study (Table 4) have correlation to later susceptibility to depression or anxiety disorders. Mismatch Negativity might provide a noninvasive bedside method for detecting functional neurotoxicity associated with fetal drug exposure and, thus, offer the possibility of early supportive measures [45,46].

Together with other neuroscience techniques, auditory ERP responses have supported the idea of continuity in the development of language starting from the early precursors of language in the first year of life to full-blown linguistic abilities [47]. Fetal AED exposure has shown to interfere with several developmental events in rodent

Fig. 1. Standard-subtracted mean amplitudes frontal areas. The waveforms of the standard-subtracted mean amplitudes of the newborns exposed to antiepileptic drugs (AED, red line) and control newborns (blue dotted line) from frontal (F) electrode areas. Reactions to each emotional variant and each deviant are illustrated in separate figures: *Happy*, *Sad*, *Angry*, *Vowel change*, *Vowel duration*, *Positive intensity change*, *Negative intensity change*, *Positive frequency change*, and *Negative frequency change*.

F



C




Table 4
Outlines of significant differences in standard-subtracted mean amplitudes.

Comparisons	AED vs. controls		Mono vs. controls		Poly vs. controls		Mono vs. poly	
	100–200 ms	300–500 ms	100–200 ms	300–500 ms	100–200 ms	300–500 ms	100–200 ms	300–500 ms
Emotional variants								
Happy /ta-ta:/								
Frontal	#	0.04	#	#	#	#	#	#
Central	#	#	#	#	3	#	#	0.08
Sad /ta:-ta:/								
Frontal	#	#	#	#	0.06	#	0.02	#
Central	#	#	#	#	#	#	#	#
Angry /ta-ta/								
Frontal	#	#	#	#	0.01	#	0.02	#
Central	#	#	#	#	#	#	0.06	0.04
Deviants								
Vowel change /ta-to/								
Frontal	#	#	#	#	0.06	#	0.01	#
Central	#	#	#	#	#	#	#	#
Vowel duration /ta-ta:/								
Frontal	#	0.07	#	#	#	#	#	#
Central	#	#	#	#	#	#	#	#
Frequency (positive) /ta-ta/								
Frontal	#	#	#	#	<0.001	#	0.005	#
Central	#	#	#	#	0.03	#	0.06	0.10
Frequency (negative) /ta-ta/								
Frontal	#	#	#	#	#	#	#	#
Central	#	#	#	#	#	#	#	#
Intensity (positive) /ta-ta/								
Frontal	#	#	#	#	#	#	#	#
Central	#	#	#	#	#	#	#	#
Intensity (negative) /ta-ta/								
Frontal	#	#	#	#	#	#	#	#
Central	#	#	0.06	#	#	#	#	#

p-Values (t-test) 0.05–0.1 are represented with blue, p-values <0.05 with red, and p-values >0.1 with #. Bonferroni correction was applied.

AED = newborns exposed to antiepileptic drugs.

Mono = newborns exposed to one drug.

Poly = newborns exposed to more than one drug.

brain including apoptosis and myelination [44,48]. These particular developmental phenomena are reflected in evoked potentials [49]. Functional toxicity of fetal AED exposure seen in previous studies [4] may thus have origin in disturbing neonatal developmental “building blocks” [48,50].

By investigating auditory responses already as early as the neonatal time period, we were able to focus on the effects of fetal exposure and minimize the effect of imminent environmental confounders encountered in long-term follow-up studies. Subtle drug-related alterations in auditory processing measured in the newborn period may lead to developmental compromises later in life. This remains unanswered in the present study and will require further follow-up of the cohort.

5. Conclusions

Prenatal exposure to AEDs may affect auditory processing of emotionally relevant information. This may be detected within the first postnatal weeks. These findings are preliminary and require replication in a larger cohort. Clinical relevance and possible applications of evaluating early auditory processing still require further research.

Declaration of competing interest

None of the authors have potential conflicts of interest to be disclosed.

Fig. 2. Standard-subtracted mean amplitudes central areas. The waveforms of the standard-subtracted mean amplitudes of the newborns exposed to antiepileptic drugs (AED, red line) and control newborns (blue dotted line) from central (C) electrode areas. Reactions to each emotional variant and each deviant are illustrated in separate figures: *Happy*, *Sad*, *Angry*, *Vowel change*, *Vowel duration*, *Positive intensity change*, *Negative intensity change*, *Positive frequency change*, and *Negative frequency change*.

Acknowledgments

Funding

The study was supported by Foundation for Pediatric Research, Arvo and Leo Ylppö Foundation, Lastenlinna Foundation, Sigrid Juselius Foundation, and Märta Donner Foundation, Academy of Finland grant 288220 (SV) and Finnish Cultural Foundation.

References

- [1] Viinikainen K, Heinonen S, Eriksson K, Kälviäinen R. Community-based, prospective, controlled study of obstetric and neonatal outcome of 179 pregnancies in women with epilepsy. *Epilepsia* 2006;47:186–92.
- [2] Pennell PB. Use of antiepileptic drugs during pregnancy: evolving concepts. *Neurotherapeutics* 2016;13:811–20.
- [3] Artama M, Gissler M, Malm H, Ritvanen A, Drug and Pregnancy Group. Effects of maternal epilepsy and antiepileptic drug use during pregnancy on perinatal health in offspring: nationwide, retrospective cohort study in Finland. *Drug Saf* 2013;36:359–69.
- [4] Baker GA, Bromley RL, Briggs M, Cheyne CP, Cohen MJ, García-Fiñana M, et al. IQ at 6 years after in utero exposure to antiepileptic drugs: a controlled cohort study. *Neurology* 2015;84:382–90.
- [5] Tomson T, Xue H, Battino D. Major congenital malformations in children of women with epilepsy. *Seizure* 2015;28:46–50.
- [6] Videman M, Stjerna S, Roivainen R, Nybo T, Vanhatalo S, Gaily E, et al. Evidence for spared attention to faces in 7-month-old infants after prenatal exposure to antiepileptic drugs. *Epilepsy Behav* 2016;64:62–8.
- [7] Gaily E, Kantola-Sorsa E, Hiilesmaa V, Isoaho M, Matila R, Kotila M, et al. Normal intelligence in children with prenatal exposure to carbamazepine. *Neurology* 2004;62:28–32.
- [8] Bromley RL, Mawer GE, Briggs M, Cheyne C, Clayton-Smith J, García-Fiñana M, et al. The prevalence of neurodevelopmental disorders in children prenatally exposed to antiepileptic drugs. *J Neurol Neurosurg Psychiatry* 2013;84:637–43.
- [9] Christensen J, Gronborg TK, Sorensen MJ, Schendel D, Parner ET, Pedersen LH, et al. Prenatal valproate exposure and risk of autism spectrum disorders and childhood autism. *JAMA* 2013;309:1696–703.
- [10] Naatanen R, Paavilainen P, Rinne T, Alho K. The mismatch negativity (MMN) in basic research of central auditory processing: a review. *Clin Neurophysiol* 2007;118:2544–90.
- [11] Alho K, Sainio K, Sajaniemi N, Reinikainen K, Näätänen R. Event-related brain potential of human newborns to pitch change of an acoustic stimulus. *Electroencephalogr Clin Neurophysiol* 1990;77:151–5.
- [12] Cheour-Luhtanen M, Alho K, Kujala T, Sainio K, Reinikainen K, Renlund M, et al. Mismatch negativity indicates vowel discrimination in newborns. *Hear Res* 1995;82:53–8.
- [13] Kostilainen K, Wikstrom V, Pakarinen S, Videman M, Karlsson L, Keskinen M, et al. Healthy full-term infants' brain responses to emotionally and linguistically relevant sounds using a multi-feature mismatch negativity (MMN) paradigm. *Neurosci Lett* 2018;670:110–5.
- [14] Depoorter A, Fruh J, Herrmann K, Zanchi D, Weber P, et al. Predicting neurodevelopmental outcome in preterm born infants using auditory event-related potentials: a systematic review. *Neurosci Biobehav Rev* 2018;89:99–110.
- [15] Suppiej A, Mento G, Zanardo V, Franzoi M, Battistella PA, Ermani M, et al. Auditory processing during sleep in preterm infants: an event related potential study. *Early Hum Dev* 2010;86:807–12.
- [16] Kujala T, Lovio R, Lepisto T, Laasonen M, Näätänen R. Evaluation of multi-attribute auditory discrimination in dyslexia with the mismatch negativity. *Clin Neurophysiol* 2006;117:885–93.
- [17] Lepisto T, Kajander M, Vanhala R, Alku P, Huotilainen M, Näätänen R, et al. The perception of invariant speech features in children with autism. *Biol Psychol* 2008;77:25–31.
- [18] Lepisto T, Kujala T, Vanhala R, Alku P, Huotilainen M, Näätänen R. The discrimination of and orienting to speech and non-speech sounds in children with autism. *Brain Res* 2005;1066:147–57.
- [19] Lovio R, Naatanen R, Kujala T. Abnormal pattern of cortical speech feature discrimination in 6-year-old children at risk for dyslexia. *Brain Res* 2010;1335:53–62.
- [20] Guttorm TK, Leppanen PH, Hamalainen JA, Eklund KM, Lyytinen HJ. Newborn event-related potentials predict poorer pre-reading skills in children at risk for dyslexia. *J Learn Disabil* 2010;43:391–401.
- [21] Guttorm TK, Leppanen PH, Poikkeus AM, Eklund KM, Lyytinen P, Lyytinen H. Brain event-related potentials (ERPs) measured at birth predict later language development in children with and without familial risk for dyslexia. *Cortex* 2005;41:291–303.
- [22] Stephen JM, Hill DE, Peters A, Flynn L, Zhang T, Okada Y. Development of auditory evoked responses in normally developing preschool children and children with autism spectrum disorder. *Dev Neurosci* 2017;39:430–41.
- [23] Kujala T, Halmetoja J, Naatanen R, Alku P, Lyytinen H, Sussman E. Speech- and sound-segmentation in dyslexia: evidence for a multiple-level cortical impairment. *Eur J Neurosci* 2006;24:2420–7.
- [24] Fan YT, Cheng Y. Atypical mismatch negativity in response to emotional voices in people with autism spectrum conditions. *PLoS One* 2014;9:e102471.
- [25] Lindstrom R, Lepisto-Paisley T, Vanhala R, Alén R, Kujala T. Impaired neural discrimination of emotional speech prosody in children with autism spectrum disorder and language impairment. *Neurosci Lett* 2016;628:47–51.
- [26] Hung AY, Ahveninen J, Cheng Y. Atypical mismatch negativity to distressful voices associated with conduct disorder symptoms. *J Child Psychol Psychiatry* 2013;54:1016–27.
- [27] Paris M, Mahajan Y, Kim J, Meade T. Emotional speech processing deficits in bipolar disorder: the role of mismatch negativity and P3a. *J Affect Disord* 2018;234:261–9.
- [28] Sokka L, Huotilainen M, Leinikka M, Korpela J, Henelius A, Alain C, et al. Alterations in attention capture to auditory emotional stimuli in job burnout: an event-related potential study. *Int J Psychophysiol* 2014;94:427–36.
- [29] Pakarinen S, Sokka L, Leinikka M, Henelius A, Korpela J, Huotilainen M. Fast determination of MMN and P3a responses to linguistically and emotionally relevant changes in pseudoword stimuli. *Neurosci Lett* 2014;577:28–33.
- [30] Meador KJ, Baker GA, Browning N, Cohen MJ, Bromley RL, Clayton-Smith J, et al. Fetal antiepileptic drug exposure and cognitive outcomes at age 6 years (NEAD study): a prospective observational study. *Lancet Neurol* 2013;12:244–52.
- [31] Meador KJ, Loring DW. Prenatal valproate exposure is associated with autism spectrum disorder and childhood autism. *J Pediatr* 2013;163:924.
- [32] Videman M, Tokariev A, Stjerna S, Roivainen R, Gaily E, Vanhatalo S. Effects of prenatal antiepileptic drug exposure on newborn brain activity. *Epilepsia* 2016;57:252–62.
- [33] Dubowitz L, Dubowitz V, Mercuri E. The neurological assessment of the preterm & full-term newborn infant. London, United Kingdom: Mac Keith Press; 1999.
- [34] Oram Cardy JEF, Flagg EJ, Roberts W, Roberts TPI. Auditory evoked fields predict language ability and impairment in children. *Int J Psychophysiol* 2008;68:6.
- [35] Hovel H, Partanen E, Tideman E, Stjernqvist K, Hellström-Westas L, Huotilainen M, et al. Auditory event-related potentials are related to cognition at preschool age after very preterm birth. *Pediatr Res* 2015;77:570–8.
- [36] Naatanen R, Paavilainen P, Tiitinen H, Jiang D, Alho K. Attention and mismatch negativity. *Psychophysiology* 1993;30:436–50.
- [37] Tomson T, Battino D, Perucca E. Valproic acid after five decades of use in epilepsy: time to reconsider the indications of a time-honoured drug. *Lancet Neurol* 2016;15(2):210–8.
- [38] Sun W, Wang W, Wu X, Wang Y. Antiepileptic drugs and the significance of event-related potentials. *J Clin Neurophysiol* 2007;24:271–6.
- [39] Rector DM, Schei JL, Rojas MJ. Mechanisms underlying state dependent surface-evoked response patterns. *Neuroscience* 2009;159:115–26.
- [40] Zhang Q, Cheng Q, Li H, Dong X, Tu W. Evaluation of auditory perception development in neonates by quantitative electroencephalography and auditory event-related potentials. *PLoS One* 2017;12:e0183728.
- [41] Wunderlich JL, Cone-Wesson BK. Maturation of CAEP in infants and children: a review. *Hear Res* 2006;212:212–23.
- [42] Sousa AC, Didone DD, Sleifer P. Longitudinal comparison of auditory steady-state evoked potentials in preterm and term infants: the maturation process. *Int Arch Otorhinolaryngol* 2017;21:200–5.
- [43] Dow C, Seidenberg M, Hermann B. Relationship between information processing speed in temporal lobe epilepsy and white matter volume. *Epilepsy Behav* 2004;5:919–25.
- [44] Naatanen R, Sussman ES, Salisbury D, Shafer VL. Mismatch negativity (MMN) as an index of cognitive dysfunction. *Brain Topogr* 2014;27:451–66.
- [45] Lyytinen H, Erskine J, Hamalainen J, Torppa M, Ronimus M. Dyslexia-early identification and prevention: highlights from the Jyväskylä longitudinal study of dyslexia. *Curr Dev Disord Rep* 2015;2:330–8.
- [46] Guzzetta F, Conti G, Mercuri E. Auditory processing in infancy: do early abnormalities predict disorders of language and cognitive development? *Dev Med Child Neurol* 2011;53:1085–90.
- [47] Kuhl P, Rivera-Gaxiola M. Neural substrates of language acquisition. *Annu Rev Neurosci* 2008;31:511–34.
- [48] Verrotti A, Scaparrotta A, Cofini M, Chiarelli F, Tiboni GM. Developmental neurotoxicity and anticonvulsant drugs: a possible link. *Reprod Toxicol* 2014;48:72–80.
- [49] Shah SN, Bhargava VK, Johnson RC, CM McKean. Latency changes in brain stem auditory evoked potentials associated with impaired brain myelination. *Exp Neurol* 1978;58:111–8.
- [50] Inoyama K, Meador KJ. Cognitive outcomes of prenatal antiepileptic drug exposure. *Epilepsy Res* 2015;114:89–97.